

GEOLOGI LIARD FLUORSPAR PROJECT (May 20, 1972 to November 10 1972)

> MINERAL CLAIMS TEE, TAM, FIRE, WEST, GEM Included within Gem/72, Tam/72, Coral/72, Bar/72, Fire/72, West/72, North-Tee/72

LIARD MINING DIVISION

Eight miles north of Mile 496. Alaska Highway at latitude 59°32'N, longitude 126°05'W

Department of					
Mines	s and	ana ta c	sien (la cources	
, č ^s		E .	1X 2.7	· ŘÍ	

ЪУ

J. R. Woodcock Consultants Ltd.

ЖČ

Mineral Claims owned by - CONWEST EXPLORATION COMPANY LIMITED JOREX LIMITED and J. R. WOODCOCK

North Vancouver, British Columbia

TABLE OF CONTENTS

Page No.

INTRODUCTION	1
LOCATION AND ACCESS	2
CLAIMS AND OWNERSHIP	2
GENERAL GEOLOGY	3
FLUORSPAR OCCURRENCES	5
Tam Deposit	5
Tee Deposit	6
Coral Deposit	7
Fire Deposit	8
Cliff Prospect	9
Gen A and E Showings	10
Camp Prospect	10
Nick Prospect	11
Henry Prospect	11
Bar Prospect	12
Strap Prospect	13
CONCLUSIONS AND RECOMMENDATIONS	14

APPENDICES

-

Cost Data Survey Data Claim Data

.

FIGURES

		Page No.
井1 1.	Index Map	2 a
#22.	Claims Location Map (scale 1:50,000)	In Pocket
₩33.	General Geology Gem Sheet (scale $1^n = 800$ feet)	In Pocket
#44.	General Geology Tam Sheet (scale 1" = 800 feet)	In Pocket
井5 5.	General Geology Tee Sheet (scale 1" = 800 feet)	In Pocket
#6 6.	General Geology West Sheet (scale $1^{"}$ = 800 feet)	In Pocket
井了 7.	Ground Survey Map (1" = 800 feet)	In Pocket
#§ 8 .	Tam Deposit Geology (scale 1" = 50 feet)	In Pocket
#9 9.	Tam Deposit Transit Survey (scale 1" = 50 feet)	In Pocket
#1010.	Tee Deposit Geology (1" = 50 feet)	In Pocket
# 11.	Tee Deposit Transit Survey (scale 1" = 50 feet)	In Pocket
#12 12.	Coral Deposit Geology (scale $1^n = 50$ feet)	In Pocket
#13 13.	Coral Deposit Transit Survey (scale 1" = 50 feet)	In Pocket
#1414.	Fire Deposit Geology (scale 1" = 50 feet) Contour Map	In Pocket
#15 15.	Fire Deposit Transit Buryey (scale 1" = 50 feet)	In Pocket
#16 16.	Cliff Prospect Geology (scale 1" = 50 feet)	In Pocket
#1717.	Gem Prospects Geology (scale 1" = 100 feet)	In Pocket
#18 18.	Camp Prospect Geology (scale $1^{\nu} = 50$ feet)	In Pocket
#19 19.	Nick Prospect Geology (scale 1" = 50 feet)	In Pocket
#20 20.	Henry Prospect Geology (scale $1^n = 50$ feet)	In Pocket
#21 21.	Bar Prospect Geology (scale 1" = 50 feet)	In Pocket
	Bar Prospect Transit Survey (scale 1" = 50 feet)	In Pocket
#JZ 23.	Strap Prospect Geology (1" = 50 feet)	In Pocket

LIARD FLUORSPAR PROJECT

INTRODUCTION

Regional prospecting in July of 1971 by a four-man crew with helicopter support resulted in the discovery of fluorspar prospects near the Alaska Highway of northern British Columbia. Several very good prospects were discovered north of Liard Hot Springs and a large group of claims was staked to cover the widely spaced fluorspar prospects. The surface exposures of mineralization on the Tam and the Tee prospects were mapped and sampled before a major forest fire interrupted the work in August.

In early September, 1971 the exploration crews returned to the Liard Hot Springs area where they did additional prospecting, found additional showings, and staked numerous more claims. An access road was built to the central part of the claim group and to several of the fluorspar prospects. Bulldozer trenching was done on the Tam, Camp, Coral, and Fire prospects.

In September 1971 a BQ Diamond Drill from D. W. Coates Enterprises Ltd. was employed and a total of 1891 feet of drilling in 14 holes was completed. Most of this drilling (1617 feet) was done on the Tam prospect; the remainder was on the Cliff prospect.

Many of the prospects and some of the general geology were briefly mapped in the 1971 program. In addition, bulk samples were taken from the Tam prospect, the Coral prospect, and the Fire prospect. These bulk samples and the core rejects were shipped to Lakefield Research for metallurgical tests.

The 1971 program was directed by J. R. Woodcock. Mr. K. W. Livingstone looked after the drill program and logged most of the core. Mr. Tom Wilkinson looked after road construction and general management of the camp.

In early 1972, a new company — Liard Fluorspar Mines Ltd. was formed to acquire the claims staked under the direction of J. R. Woodcock Consultants Ltd. in 1971 and also the original Gem mineral claims held for many years by Conwest Exploration Company Ltd.

In early June 1972, exploration crews again returned to the area. Diamond drilling was done by Canadian Longyear Ltd. under the supervision of Mr. Scott Zimmer of Conwest Exploration Company Ltd. Geological mapping of the mineral prospects and the claims was done by Mr. Henry Meixner, Mr. Andre Audet, and Mr. James Brander of J. R. Woodcock Consultants Ltd. J. R. Woodcock consulted on the overall program.

The present report deals with the surface geology of the fluorspar showings and the area of the claims around these showings. The regional mapping is presented on topographical maps, scale $1^{\circ} = 800^{\circ}$, contour interval 100 feet. There are four of these maps and their respective positions are shown on the index on each map. These maps are a compilation of all the mapping done on the mineral showings by the various geologists, and that done on the claims, mainly by Mr. Henry Meixner, geologist.

- 1 -

The fluorspar showings have been mapped on a scale of $1^{*} = 50^{\circ}$ with contours ranging from 10-foot intervals to 25-foot intervals. This mapping has been done by various geologists including James Brander, Henry Meixmer, J. R. Woodcock, K. W. Livingstone, Andre Audet, and Scott Zimmer.

Control for the mapping of the mineral prospects was by transit and stadia or chain done by Mr. Andre Audet, geologist, and his assistant. Each of the mineral showings has been tied to a mineral claim post. In addition a survey carried along the access roads was used to tie in many of the claim posts in the vicinity of the showings. Other claims have been plotted on the 800-scale topographical maps.

LOCATION AND ACCESS

The Liard fluorspar prospects are located near the Alaska Highway at Nile 498, about 200 miles north of Fort Melson, British Columbia (Figure 1). Fort Melson is a small town with population of 3500, serviced by the British Columbia Railway from Vancouver. The Alaska Highway is the main access road to the Yukon and is a maintained gravel surface road.

The fluorspar prospects are located along a stratigraphic contact and are thus quite widespread. The prospects are found over a distance of eleven miles along a north-south contact and six miles along an east-west trending contact.


The main prospects may be reached from the Alaska Highway by the CHT Microwave Tower all-weather gravel access read at Mile 498. About 2000 feet west of the tower on the CHT access read, a forest access read suitable for 4-wheel drive vehicles was extended to the main prospect area. In addition an access read which leaves the highway at Mile 497, extends in a northeasterly direction to the area of the original Gem prospects.

The prospect area is one of gentle to moderate relief, at an average elevation of 3000 feet. The rolling topography reflects underlying bedrock. Limestone ridges and knolls are separated by recessive areas and valleys underlain by shale. Two main crocks, Tester and Mould, crossout the major structural control and bedrecks, and flow obliquely across the prospect area.

During the first phase of work, a thick forest of balsam, lodgepole pine and spruce covered most of the region; outcrops were scarce. The forest fire killed all the timber. Old or re-burned areas are now completely bare; the abundant trees in the newly burned areas are starting to fall forming an inaccessible mass of deadfall.

CLAIMS AND CHIERRENTP

Previous to the discovery of fluorepar mineralization in July, 1971 the only claims in the prospect area were held by Convest Exploration Company Limited. The Convest property included 34 mineral claims and fractional claims (Gen 1-25, Les Fraction and New 1-8). In 1971 an additional 49 Gen mineral claims were staked and recorded for Convest Exploration Company Limited, and for Jorex Limited.



Page 2a

· · ·

In addition to the Gem mineral claims, a total of 678 claims were staked in the area. These included the Tam 1-151, Tee 1-140, Fire 1-26, Fire 35-109, West 1-128, Fall 1-76, Fall 79-106, and Hal 1-54, which were staked for J. R. Woodcock. Another claim group south of the Liard River, the Pet 1-100, were staked for Conwest Exploration Company Limited. The claims are plotted on Figure 2 and the claim data is summarized in the appendix.

Assessment work has been applied at various times in 1972 to many claims included in the appendix. This was applied by Mr. Bruce Pope of Conwest Exploration Company Limited after consulting Mr. J. R. Woodcock on the merits of the various claims. Many of the original claims staked in 1971 did not get assessment work credits and will accordingly lapse in 1972. A forthcoming separate report will outline the new status of the claims.

The mineral claims presently are in the name of J. R. Woodcock, Jorex Limited, or Conwest Exploration Company Limited. Eventually they should be transferred to the new Company, Liard Fluorspar Mines Ltd.

GENERAL GEOLOGY

Station - Antonio

The Liard Hot Springs area is the southernmost extension of the MacKenzie Mountains, where the north-south MacKenzie Mountain structures are crosscut by the dominantly northwest-southeast Rocky Mountain structures.

The Liard Hot Springs fluorspar deposits are located along a contact between limestone and shale of Middle Devonian age. This area is one of several outcrop areas of rocks of similar lithology and age occurring in northeastern British Columbia.

Gabrielse (1963) in his regional mapping in the area did not attempt to attach formational names to the stratigraphic rock units. However, since that time, additional study by Gabrielse (1968), Taylor (1969), and Griffin (1967) have shown that the rocks are lithologically similar to those to the south where formational names have been defined.

Taylor (1969) describes the limestone as Dunedin Formation. Taylor and MacKenzie (1970) in Bulletin 186 give descriptions of the Dunedin The Dunedin Formation is a uniform sequence of argillaceous, as follows: in places siliceous and dolomitic, dark-grey bedded limestones that overlies light-coloured dolomites of the Stone Formation, and that is overlain by the Besa River shales. Dolomite is present in the lowermost strata in parts of this Formation. These lower dolomites are argillaceous, finely crystalline and thinly bedded and they commonly contain dark shale partings, thin, laterally restricted intercalations of shale and scattered sandstone lenses. Dolomite also occurs throughout the Formation as small euhedral crystals sporadically associated with the rock fractures, as partial or complete replacement of fossil remains, in vugs, or as diffuse patches and mottlings in some of the more argillaceous beds. The upper two-thirds of the sequence consists of thick-bedded to massive, finely crystalline dark-grey limestone and interbedded granular limestone. In places the

- 3 -

upper strata are conspicuously siliceous, particularly in the upper 100-foot interval where lenses and nodules of black chert are common. The upper 30 feet of the limestone are siliceous and argillaceous and a dark-grey black colour and very fossiliferous, particularly on the upper surface where many coral colonies are present. There are many remains of brachiopods, gastropods, and crinoids, and scattered chert nodules occur at the base of this 30-foot unit.

In the vicinity of the Liard Hot Springs, mapping has shown that the contact area between the Dunedin Limestone and the overlying Besa Shales is characterized by many coral colonies.

Overlying the limestone is a thick succession of black shale, siltstone, and minor calcareous siltstone and sandstone. In the prospect area there is no faunal evidence to date the shales. Gabrielse found some Upper Devonian and Mississippian fossils in the shales exposed to the west in the Liard River and Smith River areas. However, these shales form a separate fault block in which the contact with limestone is not exposed and thus their exact location in the shale section is not known. To the south and east, the shale overlying the Dunedin Formation is known as the Besa River Formation. The shale at the limestone contact becomes progressively older in age to the north and northwest. North of the prospect area it is correlated with the Funeral Formation (Douglas and Norris, 1961) of early Middle Devonian age (Griffin, 1967).

Taylor (1969) uses the Besa River Formation in his mapping of the area as defined by Kidd (1962, 1963). This is a thick sequence of black shale in northeastern British Columbia which lies between Mississippian cherty limestone and Middle Devonian carbonates. The problem of subdividing the Besa River Formation arose because of the lack of fauna and lithological data at the time the unit was named. Thus the pelitic section in the prospect area is undivided and grouped into Besa River Formation.

There is some doubt as to the nature of the shale - limestone contact in the prospect area. Along the eastern mountain front south of Summit Lake and in the mountain range northeast of Summit Lake (Mile 392 Alaska Highway) the contact appears quite conformable with no conspicuous irregularities to denote an erosional disconformity. However in the vicinity of the fluorspar prospects north of Liard Hot Springs, outcrops are scarce. Surface mapping done in previous years showed great irregularities in the position of the contact and thus indicated a possible disconformity. Subsequent diamond drilling has shown that a large portion of this irregularity is due to tectonic disturbance -mainly caused by thrusting, with consequent folds and thrust faults.

The data obtained thus far is not sufficient to determine the attitude of all folds and thrusts. There appears to be a linear northsouth zone of disturbance which runs through the mineralized area. In places (e.g. upper Teeter Creek) it coincides with the west side of north-south valleys.

To the west of this disturbed zone the thrusting appears to be from the west with the thrust faults dipping westerly.

A prominent shale ridge occurs to the east of the disturbed zone

and extends northward from the Alaska Highway for at least 20 miles. The shale ridge is separated from the mineralized shale - limestone contacts in the disturbed area by valleys that obscure the relationship between the two geological terraines. The shale ridge to the east of the valley is topographically higher than the shale - limestone contact to the west. The structural relationship is not apparent, but one must not overlook the possibility of overthrusting from the east such as mapped on the La Biche sheet only 30 miles to the northnortheast.

FLUORSPAR OCCURRENCES

Tam Deposit

Location:

The Tam deposit occurs on the Tam 2 and Tam 4 mineral claims at the end of the access road. It is exposed on the steep slope just east of the main structural valley. It has been exposed by numerous surface trenches and explored by 22 short drillholes.

Geology:

Mineralization occurs at the contact between the limestone and the overlying Besa River shales; a contact which generally dips easterly. In the zone of mineralization the limestone and the shale at the contact are brecciated. The shale breccia, in many places, consists of chaotic angular blocks. The mineralization occurs in the breccia and in fractures in the overlying shale and underlying limestone.

The mineralized zone, which is of variable thickness, generally also dips easterly and it pinches rapidly to the east where there is no shale breccia. Along the west side, the limestone - shale contact is eroded; and the west boundary of "ore" is quite sharp against barran limestone.

The mineralization extends along the hillside in a northerly direction for 900 feet as indicated by surface exposures and diamond drillholes. It is still open to the north. Widths vary from a minimum of 160 feet to a maximum of 550 feet.

Although over the large part of the length of the mineralized deposit, the control appears quite simple with the best mineralization occurring in the breccis, especially the limestone, some structural complexities occur towards the north. For instance in diamond drillhole 72-52 limestone breccia with good fluorite mineralization occurs above shale breccia with only minor fluorite mineralization. It is quite probable that had the drillhole been continued it would have intersected additional limestone below the shale breecia. The data is inadequate to determine whether this complexity is due to sharp folding or to some thrust faulting. Such a complexity could be attributed to thrusting from the west which has thrust the breccisted limestone over top of the brecciated shale. Such a thrust fault could also pass through the limestone breccia of Hole #72-53 but not be recognized because of the high degree of replacement often found within the limestone breccia. In fact much of the breecis could have originally been formed by fracturing along the thrusts and the precciated nature could have subsequently been amplified by solution of the limestone and collapse of the adjacent rocks. If such suggested thrusting is present and is

- 5 -

important in the ore control, and if impricate thrusting is present then great potential exists in the northern part of this deposit for finding additional somes under either barren limestone or barren shale.

The best mineralization occurs in the limestone broccin and in places the rock is almost completely replaced by fluorite plus barium minerals (witherite, barytocalcite, barite). A few silicocous bands may be cherty layers left from the original limestone. Calcite also occurs as a few limestone remnants. The fluorite is generally purple to black. The black variety appears to have more impurities (e.g. fine-grained quarts) than does the purple variety.

The Tee Deposit

Location:

The Tee prospect is on the Tee 3, 5, 16, and 18 mineral claims, on the steep valley slopes west of Teeter Creek. The mineralized exposures also lie a short distance north of a small stream that enters Teeter Creek from the west. The mineralization is exposed on the steep (about 25°) northeasterly-facing slopes of Teeter Creek and also on the much steeper cliffs and slopes that face southward and form the valley sides for the small stream from the west.

Geology:

The mineralization is generally at the contact of the Dunedin limestone with the overlying Besa River shale. This contact has a general overall dip of about 25° easterly; however numerous little detailed irregularities change this dip locally and also make for discontinuities in thickness of mineralization. Generally the shale and some of the limestone are brecciated in the vicinity of the mineralization.

To facilitate the description, four areas of mineralization labelled zones "A" to "D" inclusive are indicated on the map.

Zone A, the most northwesterly one exposed, consists of remnants of highly mineralized limestone surrounded by and presumably resting on areas of barren limestone. A few shale remnants in the eastern part of this zone indicates that the some was originally at the limestone shale contact. The mineralization in this zone is somewhat unique for the Liard fluorspar area in that most of the fluorite is colourless. Mineralization of this zone has been mapped over an area 300 feet long and up to 220 feet wide, or a surface area of about 50,000 square feet. The high grade exposure in the southwest part of the zone has an exposed thickness up to 15 feet.

Zone B is exposed on the sharp ridge that occurs on the interfluve between Teeter Creek and the above mentioned subsidiary stream from the west. The mineralization in this place is also at the contact of shale and limestone with considerable replacement of shale breccis and limestone breccis. The steep cliffs at the west end of the exposure show that the main "ore" zone is underlain by fractured limestone containing abundant veins and lenses of fluorite. The exposures of good grade mineralization at the west end of Zone B are over a vertical interval of 60 feet. The mineralized area extends easterly down the steep hillside (approximately 30°), for a distance of 700 feet with an average width of about 200 feet. Some prominent east-west fracturing or familting is exposed in places and this might indicate some additional east-west control of this zone. If such is the ease, then the position of greatest thickness would also have an east-west strike. The position of the postulated greatest thickness is unknown. It could occur anywhere within the exposed part of Zone B; it could have occurred to the south of Zone B and have been removed by erosion; or it may have never existed!

Zone C includes the mineralization that occurs in the vicinity of some limestone cliffs trending at 330° aximuth across the property. Most of this mineralization is exposed along the face of the cliffs in discontinuous lenses. Shale remnants are also found along these limestone cliffs. The strike of these limestone cliffs is essentially parallel to the valley side and also almost parallel to the strike of the limestone shale contact. Presumbly it is the major regional strike in this part of the Taeter Creek valley. The exact structure along these cliffs is not evident. However there is some suggestion that the limestone has been thrust from the west over shale and shale breccis. The amount of exposed mineralization is not significant in the overall ore reserve picture.

Zone D includes an area of exposures of shale and shale breccia lying downslope to the east of the limestone cliffs of Zone C. Some of these exposures are mineralized with barium carbonates and/or fluorspar. The exposures of shale and shale breccia occur over an area 250 x 300 feet.

Shale breccia, especially mineralized shale breccia, is very important in that it overlies most of the better mineralization on many of the other deposits. Therefore should one find mineralized shale breccia it constitutes a prime exploration target and drillholes should be collared within this breacia some to explore the underlying contact with the limestone. The shale breccis of Zone D was supposed to be explored by drillholes 62, 63, and 64. However because of difficulty in getting the drills up the steep moss-covered slopes (with underlying parmafrost) the holes were collared east of the exposed exploration target. Hole #62 was a vertical hole drilled to 204 feet and it never got out of the barren shale. Hole #63 drilled northwesterly intersects barren shale to a depth of 181 feet. From 181 to 249 feet the drill intersected fractured limestone containing fluorite mineralization in fractures. The average grade of this 60 feet was 8.8% CaFo. Hole #64 drilled westerly at an angle of 45° intersected shale to a depth of 122 feet and barren limestone from 122 to 208 feet. These drillholes delimit any potential mineralized zone to the east; however they do not explore the possibility of mineralization underlying the mineralized shale breccia. The mineralization encountered in Hole #63 is encouraging; however one must note that this intersected mineralization is over 200 feet vertically below the exposures of mineralized shale breccia.

Coral Deposit

Ĺ

Location:

The Coral deposit is on mineral claims Tam 23 and Tam 24 about 3500 feet southwest of the Tam deposit. An access road has been built across the mineralized zone from the main access road. The prospect has been exposed in numerous transhes and tested with 12 short drillholes.

The exposures are on the gentle south slope of a small limestone hill in an area of forest cover and widespread overburden. Geology:

The fluorspar mineralization is at the limestone - shale contact. This contact dive gently southward from the mineralized exposures. In most cases there is some brecciation of the shale and/or the limestone along this contact and considerable intersections of both the limestone breccia and shale breccia have been logged in the core. However the exposures do not have the chaotic breccia of large blocks that are visible in the exposures on the Tam prospect. Whether or not such chaotic breccia occurs in the core is not evident from the data. Much of the breccia that has been logged as such is broken rock some of which consists of remnant shale fragments in a replacement matrix, without the chaotic orientation of the blocks.

The largest exposure of mineralization occurs at the west end of the prospect on the Tam 24 mineral claim. This mineralization is exposed over a length of 300 feet (in a northwesterly direction) and a width of 150 feet. However the appearance of this mineralization in the trenches is that it is a skin or remnant left on top of the limestone formation and that it has very little vertical extent.

At the main showing (on Tam claim 23) mineralization is exposed in trenches over an east-west distance of 600 feet and a northsouth distance of about 200 feet. Diamond drilling has followed the mineralization under the shale capping for an additional 500 feet to the south-southeast.

The drilling down the central part of this zone has intersected up to 87 feet of good mineralization (DDH 72-43 intersected 87 feet grading 39% CaFp). The best mineralization occurs in holes 72-40, 72-42, and 72-43. In these three better holes the mineralization is largely confined to limestone breccia with lower grade mineralization over small thicknesses in the overlying shale breccia. In other holes to the east and to the west the mineralized zone has pinched, probably quite erratically. Hole 72-40 in the centre area of the mineralized exposures is collared in breccia, mainly limestone breccia. However the first 20 feet of core are relatively barren: thus at this place the mineralized zone is below barren limestone. The north-south section of the three high grade holes shows that the upper part of the mineralized zone in each of the three holes forms a fairly straight line. Two of the points of the upper "ore" contact are at the shale limestone contact and the third point (Hole 72-40) is within the limestone breccia. Thus it is possible that there is some faulting in addition to the contact which has controlled the mineralization, and that this faulting is generally along the contact; but in places it leaves the contact and goes into the adjacent formations.

Fire Deposit

Location:

The mineralization of the Fire prospect is mainly on Fire 46 and Fire 48 mineral claims but extends onto the Fire 45 and Fire 47 mineral claims.

The property can be reached along the east-west access road which extends westward from the main access road (about 6000 feet). The property has been investigated with some bulldozer tranches and with 18 short diamond drillholes.

Geology:

1

In this area the limestones are generally flat-lying; remants of overlying shale are widespread. In places, this overlying shale appears to be breccisted and also mineralized with fluorspar and barium minerals.

At the southeast end of the prospect, a narrow, highly mineralized zone extends at azimuth 330° for 900 feet. It has exposed widths between 100 feet and 200 feet. Diamond drilling has shown that most of the fluorspar mineralization occurs in the shale breccia with some underlying limestone breccia. The thicknesses of the mineralized zones are generally less than 50 feet.

Fluorspar mineralization has been noted in bulldoser trenches for an additional 1600 feet to the northwest. Most of this is in shale or shale breccia with values generally less than 10% CaP₀. However at the extreme northwest end of the drilling (DDH 72-37) mineralization also occurs in limestome breccia below the shale breccia. The lower part of this mineralized limestome breccia grades 37% CaP₂ over a thickness of 65 feet. The closest hole is 600 feet away so the extent of the mineralized limestome breccia is unknown.

Cliff Prospect

Location:

The Cliff prospect is exposed in a northerly-trending zone along the mutual boundaries of the West 55 and 56 claims and the West 57 and 58 claims. It can be reached by the access road which extends westward past the Fire deposit.

In this area exposures are mainly flat-lying limestones which form a series of small benches, generally separated by some linear zone possibly representing a fault or fracture zone. The area is gently undulating and the dense forest cover has been killed by the 1971 forest fire.

The prospect has been investigated with four short drillholes.

Geology:

This linear zone of mineralization is exposed along the face of some low limestone cliffs over a north-south length of 500 feet, and up to 100 feet wide. On the east it is bounded by an upper bench of flatlying barren limestone. On the west it is separated from another hill of flat-lying barren limestones by an overburden-covered north-south pass.

The fluorspar mineralization is unusual for this mining camp in that a horizontal banding occurs in the main exposure. Also much of the fluorite is colourless, although purple varieties also occur. The banded "ore" is quite siliceous and has been mapped as impure quartzite. Possibly some siliceous zones were present in the limestone. However the silice could have been added during the introduction of the fluorspar.

Just south of the mineralized exposures, shale debris and one shale outcrop have been noted. A drillhole in this south area intersected a very thin layer of fluorite mineralization at the contact of the shale and the underlying limestone.

The linear nature of the some and the presence of shale breccia associated with the mineralization indicates a northerly-striking structure, probably fault controlled, into which some brecciated shale fragments have moved. Possibly the solution collapse along this fault some allowed the shale fragments to drop down into their present position.

- 9 -

Gen A and E Showings

Location:

The Gem A and E showings were discovered during the geological mapping in 1954. In 1972 three short holes were placed in the Gem A showing and 14 short holes were placed in the Gem E showing.

11

These showings are north of the Liard Hot Springs Park and can be reached by the access roads that extend northerly from Mile 497. These access roads were built about 20 years ago and rebuilt and extended by the Forest Service during the forest fire of 1971.

Geology:

The Gen A showing is the original discovery of fluorspar in this region. It occurs as a replacement under shales on a very small flat-topped butte which lies in the centre of a small depression. This appears to be an erosion remnant of what was probably a much more extensive deposit. The thickest mineralization occurs on the northeast corner of this butte (about 30 feet thick). However the mineralization appears in outcrop to pinch to the southeast and this has been confirmed in the diamond drillholes of 1972.

The Gen E showing consists of mineralization in shales at the contact with the underlying limestone. Several exposures were mapped over an area about 1000 feet x 900 feet (around the circumference of the shale exposures). Diamond drilling in 1972 explored the contact area beneath this shale. The diamond drilling has shown that the mineralization is quite extensive along the shale - limestone contact; all of the holes encountered some mineralization. However in most places the mineralization was less than 10 feet thick. Three holes (72-12, 72-14, 72-8) encountered somewhat greater widths with the best mineralization in hole 72-12 (50 feet grading 39% CaF₂). From the present data the mineralization appears to be very discontinuous and lensy along this contact. However the drilling has been insufficient to eliminate the possibility of some of these greater thicknesses expanding into small reserves.

The mineralization in the Gem E showing is unusual in that a large portion of it occurs in the "shale breccia" rather than in limestone breccia, and also there is a very high proportion of barium minerals in most of the mineralization. Much of the so-called shale breccia could be angular remnant pieces of shale contained within a replacement matrix of barium minerals and fluorspar.

Camp Prospect

Location:

The Camp prospect is on the Tam 6 and 21 mineral claims 900 feet southwest of the drill camp.

The mineralization is exposed on the northeast slopes of a small hill in an area of gentle topography. Overburden is extensive and the area is well forested; however the trees have all been killed in the 1971 forest fire.

Access roads have been built to this prospect and some bulldozer stripping has been done. However there has been no diamond drilling.

- 10 -

Geology:

The mineralization is again at the limestone - shale contact. The hill to the west and northwest is underlain by limestone and the lower area to the southeast underlain by shale. The mineralized shale breccia occurs over an area about 150 feet in diameter. The underlying limestone is also well mineralized in places close to the exposure of mineralized shale breccia.

Bulldozing for 800 feet to the southeast has exposed barren shale. Also, bulldozer stripping to the east-northeast has exposed barren shale. Much of the overburden in the area 1500 feet eastnortheast of the showing is deep glacial till where trenches did not meet bedrock. However abundant fluorite-bearing float (generally shale breccia) occurs throughout the till in this area (900 feet east-southeast of camp.).

No diamond drilling or sampling has been done on this prospect or in the area of overlying shale to the east of the prospect.

Nick Prospect

Location:

The Nick prospect occurs along the boundary line between the West 87 and West 89 mineral claims. An access road has been built northwestward from the Cliff prospect across Tester Creek to the Nick prospect. Some buildozer stripping has been done on the Nick prospect.

The Nick prospect is in an area of considerable forest cover and very little outerop. It occurs as a small circular hill, about 500 feet in diameter, which projects upward from the floor of a gentle basin. The mineralized exposures occur on the southeast and northwest flanks of this small flat-topped hill.

Geology:

Exposures along the sides of the hill and in trenches consist of shale, generally barren, and a shale breccia which is generally mineralized.

Mineralization consists of fluorspar and barium minerals within the "shale breccia" and within sparse fractures within the shale. The breccia may actually be remnant fragments of shale which occur in the replacing matrix of barium minerals and fluorspar. Samples taken in four places with sample results are shown on the sketch of the prospect made by Mr. Andre Audet.

The closest limestone exposure would be about 1000 feet to the east where very steep limestone cliffs fall away to the Teeter Creek below. The writer suggests that limestone could underlie this mineralized shale at shallow depth and that better grade mineralization will probably occur at the limestone contact. This can only be tested with a few vertical diamond drillholes.

The Henry Prospect

Location:

The Henry prospect occurs in the northeast corner of the Gem 27 mineral claim along the creat of a northerly-trending hill. There is no road access to the prospect. However a helicopter pad has been built near the southern end of the mineralized rocks.

Geology:

The mineralization occurs near the crest of a pronounced anticline which is exhibited in the limestone cliffs along strike to the south. A pronounced northerly-trending trough occurs near the crest of this anticline and can be followed on air photos for several miles. At first appearance this appears to be the loci of axial fracturing and has in the past been interpreted as such. However the mapping by Mr. J. Brander shows that the limestones on both sides of the trough dip gently to the east or southeast and that the structure is actually on the upper part of the eastern limb of the anticline.

Mineralization consists of fluorite, witherite, and calcite replacements in flat-lying limestone. The best exposure occurs in the bottom of the trough in the northern part of the mapped area. In this exposure the limestone is almost completely replaced over an area which has a diameter of about 100 feet.

A second exposure occurs about 500 feet further to the southsouthwest and this exposure is about 150 feet west of the trough. At the second exposure, flat-lying beds of limestone are replaced by the fluorite, calcite, and witherite over an area of about 70 feet x 40 feet. It appears that, although good grades can be obtained over thicknesses of about 20 feet, the lateral extent of the mineralization is very limited. There are also additional smaller zones of mineralization near this second southerly exposure.

Bar Prospect

Location:

The Bar prospect is on the Tam 43 mineral claim on the steep northeast flank of a sharp limestone hill. It is 1000 feet southeast of Mould Creek and 1500 feet south of the side access road that extends from the main access road down to Mould Creek.

Geology:

Mineralization consisting of barite, fluorspar, etc. occurs as a replacement in massive limestone beds that dip about 45° east. The mineralization occurs as massive pods, veins and disseminations.

Mr. Andre Audet took chip samples across three areas of the better mineralization and these returned values of 35%, 51% and 31% CaF₂. The grade of the more massive limestone containing pods and veins of fluorite is not known.

The writer suspects that this mineralization is of a very discontinuous poddy nature and will have no economic importance. However it does occur near the shale - limestone contact and it lies about halfway between the Gem showings and the Tam showings. It is important in that it demonstrates additional mineralization in the contact area between these two widely spaced centres of mineralization and it indicates that the contact lying between the Bar prospect and the Tam prospect will eventually merit investigation.

Strap Prospect

Location:

The Strap prospect occurs near the crest of some gentlyrolling limestone hills on the Tee 126 mineral claim. There is no road access to the property. There is, however, a small swampy area suitable for helicopter immediately southeast of the mineralized exposure.

Geology:

Most of the rocks in this area are barren, flat-lying limestones. The mineralized exposures occur along a zone which trends 330° azimuth. The mineralization has been followed intermittently for 600 feet. However the northern exposures are in areas of overburden and the total length and total width of the zone are not known. The maximum exposed width is about 40 feet.

Mineralization is fluorite and barium minerals, mainly barite. A chip panel sample on the southernmost exposure over an area 20 feet x 25 feet assayed 28.6% CaF₂. In addition to the exposures of fluorite and barium minerals, there is one exposure, 40 feet long, consisting of massive limonite, mainly goethite.

The lateral extent and the depth extent of the mineralization are unknown. The presence of one small shale remnant on the southernmost exposure indicates that its original limestone - shale contact would not be far removed.

The writer suggests a northwesterly-striking fault zone which has controlled the mineralization. Possibly the shale fragment was brought into its present position by fault movement or by solution and collapse. The exposed mineralization does not appear particularly interesting from an economic standpoint. However the zone to the northwest is completely covered by overburden and may have potential for more extensive mineralization.

CONCLUSIONS AND RECOMMENDATIONS

- 1. Fluorspar mineralization occurs along a north-south zone for over ten miles, and over an east-west distance of up to three miles. All of these occurrences lie in close proximity to the contact between Dunedin Limestone and overlying Besa River Shale and the best mineralization occurs at the contact.
- 2. The exact nature of this contact is not known. There are many irregularities in it many of which are caused by structural disturbances; however some could be due to an erosional surface or disconformity. The upper part of the Dunedin Limestone, just below the overlying shales, is characterized by highly fossiliferous rock, much of which is coral reef.
- 3. The mineralization appears to be controlled by structural complexities, especially a north-south zone of faulting and overthrusting from the west. The best mineralization occurs within a breccia some of which consists of a chaotic jumble of shale blocks. How much of this breccia has been caused by the faulting and how much of it is due to solution collapse is not known.
- 4. Mineralization consists of fluorspar, witherite, barvtocalcite, with minor barite and silica. Both black and purple varieties of fluorite are present, especially in the replacement of the brecciated limestone. Calcite, some of the silica, powdery hydrocarbons, and H₂S gas are probably part of the original carbonate sequence.

The best grade material occurs within the limestone breccia at the contact. However in some prospects the shale breccia has fair values in fluorspar. The adjacent or overlying shale and shale breccia and the underlying fractured limestone carry veinlets and pods of fluorspar and barium minerals.

5. The Tam deposit is the best one investigated thus far. This has been investigated by 23 short drill holes. The southern portion has better fluorspar grades than most other prospects. This southern portion has been fairly well delimited with the drilling.

The northern portion, which contains lower grade mineralization, is a much more complex structure and has not been fully investigated. Additional drill holes are needed to delimit the mineralization to the north and to determine its structure and grade.

6. The Tee deposit also has some good grade mineralization exposed at the surface. Mineralization occurs over a large area. However only a few rough panel chip samples were taken from some surface exposures. The mineralization dips easterly along the contact and may extend under some shale breccia which occurs near the foot of the hill. The mineralization is possibly open to the north where the hill is completely covered by overburden. Diamond drilling is needed in at least three of the zones of this showing to determine the grade and vertical extent.

- 7. The Coral deposit has also received sparse diamond drilling on 200-foot centres. This may have indicated a central zone of material grading over 30% CaP₂ which extends in a south-southeasterly direction under the shale capping. Adjacent to the possible central higher grade zone is lower grade material (between 15% and 20% CaP₂) which has more erratic nature (grades and thicknesses). This zone is still undelimited to the south and southeast. Further drilling is necessary to establish the grade and continuity of the possible central narrow higher grade zone, to determine the extent of the lower grade adjacent mineralization, and to determine if additional relatively high grade pods occur in the general vicinity.
- 8. The Camp prospect at present appears to be too limited to be of economic importance. However it does occur between the Coral and the Tam prospects and the till overburden in this general area contains considerable float of well mineralized breccia. Thus this complete zone, extending eastward from the Coral prospect for more than 1500 feet will eventually need investigation.

9.

The Fire deposit has well mineralized shale breccia continuous over a length of about 900 feet and a width of about 100 feet. The depth of mineralization is quite limited as shown by the drill holes placed in the main showing.

In addition, mineralized shale or brecciated shale occurs for at least 1600 feet to the northwest. The grade in this material is very low. However the most westerly hole intersected good grade mineralized limestone breccia under the shale breccia.

- 10. The Cliff prospect has good grade mineralization exposed over a length of about 500 feet. This prospect has been investigated by four short drill holes. The results indicate a very restricted zone.
- 11. Mineralization on the original Gem claims occurs in several places and has been mapped as showings Gem A to G inclusive. The highest grade material was originally exposed in the Gem A showing. A small amount of drilling to the southeast in 1972 failed to extend this mineralization enough to make it significant.
- 12. The Gem E showing was also tested by drill holes placed on 200-foot spacing. The contact between the limestone and the overlying shale is extensively mineralized but in most cases the mineralized section is very thin. Two good intersections were obtained and further drilling will be needed to determine whether these thicker zones have any significant extent.
- 13. The Nick prospect lies west of Teeter Creek and consists of fluorspar and barium mineralization around the small hill. Mineralization is largely in shale breccia; there are no limestones exposed. Diamond drill holes would be needed to determine if significant mineralization occurs at the underlying limestone shale contact.
- 14. Several other prospects were found and mapped but are not of significance as far as ore reserve goes. These would include the

-15-

Strap, Bar, and Henry prospects.

- 15. The most important untested zone of mineralization (in addition to extensions of the Coral, Tam, and Tee deposits) lies in the area of widespread overburden along the main north-south zone. The more favourable areas would include that between the Bar prospect and the Tam deposit, especially the area south of the Tam deposit and east of the Coral deposit. Another favourable area that would need testing is that immediately northwest of the Tee deposit.
- 16. The diamond drilling done in 1972 failed to prove up tonnage and grade needed for an economic operation. However the amount of drilling, even on some of the prospects, has not been sufficient to delimit the zones and there is great potential for finding additional reserves of similar grade to that already found. Until the economics of the fluorspar production in this locality improve (higher prices for fluorspar and/or cheaper transportation) the geologically favourable areas should be kept for future investigation.

J. R. Woodcock

November 8, 1972

VAGES

8 M. C. 4 4

J. Brander: June 3 to 8 6 days 11 10 to 20 11 11 12 July 10 to 20 28 days Less 4 days applied July 27, 1972 \$ 2,400,00 = 24 days @ \$100/day H. Meixner: June 1 to July 20 50 days Less 2 days applied July 27, 1972 4,080.00 = 48 days @ \$85/day A. Audet: May 23 to 26 4 days 21 May 29 to July 20 53 11 July 23 to Aug. 2 11 11 September 25 1 H October 6 1 70 days Less 5 days applied July 27, 1972 5,525.00 = 65 days @ \$85/day M. Currey: June 4 to 8 days 18 July 3 to August 2 36 days Less 2 days applied July 27, 1972 \$ 1,700.00 = 34 days @ \$50/day J. R. Woodcock: June 5, $6(\frac{1}{2})$, $7(\frac{1}{2})$, 11, $13(\frac{1}{2})$ 31 days 82 " July 4,5,6,7,9,10,12,13,31(1) 13 August $4(\frac{1}{2})$, $10(\frac{1}{2})$ 1 88 September $21(\frac{1}{2}), 26(\frac{1}{2})$ 1 October $11(\frac{1}{2})$, $16, 18(\frac{1}{2}), 30(\frac{1}{2}), 31(\frac{1}{2})$ 11 3 11 1 November 3 18 days Less 1 day applied July 27, 1972 2,550.00 = 17 days @ \$150/day \$ M. Virsumen: 45 days June 8 to July 22 Less 3 days applied July 27, 1972 688.52 = 42 days @ \$500/mo.

WAGES (cont'd)

B. Groat: June 5 to August 12 = 69 days @ \$750/mo. \$ 1,696.72 T. Drews: 1 day July 31 August 1,2,4,5,7,8,9,10,11, 14,15,17,28,29,30 15 days Sept. 7, 18, 21, 22, 25 5 days Oct. 10,11,12,13,16,17,18,30,31 9 days Nov. 1, 2, 3, 6, 7, 8, 9, 10 8 days 38 days -- 38 days @ \$44/day \$ 1,672.00 \$ 20,318.24 Total Wages

CONTINGENT HELICOPTER COSTS

June	5	70 m				
	7	7 0				
	8	00	1			
	11	~~				
	12		1			
	13	40	1			
	12 13 15 18	32	17			
July	9		н.			
•	10	160	17			
	11	100	t 1		•	
	12	100	1			
	13 14	<i>7</i> 0	ur			
	14	100	••			
	16		11			
	17	120	1F			
	18		11			
	19	100	11			
	20 27	. 🗩 🛛	11			
	~ (inutes = 33	hra. 55 min.	_	
		say 34	hours @ \$14	0/hr.	\$	\$ 4,760.00
		Less \$	93.00 applie	d on July 27/	72 _	<u>(93.00)</u>
			Tota	1 Contingent	Helico	pter Costs \$ 4,667.00
ACCO	ACOM	TION IN	FIELD			
		ander ixner	24 days 48 "	Versumen Groat	42 69	days days

Brander Meixner Audet Currey	24 48 63 34	days 	0	Versumen Froat Pilot Engineer	42 69 10	days days "		
Woodcock	10	*1		9				
Total	days	31	10 e	\$10/day	••••		 . \$	3,100.00

TRANSPORTATION

.

1

ï

-

Rental of	G.H.C. Jin	uy	\$	1,398.00	
Road gas	(approx.)		·	200,00	
		Total Transportation			\$ 1,598.00
TOTAL Wages,		Helicopter Costs, Accom and Transportation .			\$ 29,677.24

LIARD FLUCRITE GROUED SURVEY

INTRODUCTION

Critical claim posts and major fluorite showings were accurately located by a chain and transit ground survey and by triangulation. The most promising fluorite showings were surveyed by the stadia method, leaving permanent stations in the ground. Diamond drill holes on the Tam and Coral showings were tied to the main project grid by chain and transit surveys. An index of elevations and co-ordinates is appended. Those stations located by chain and transit are marked by an asterisk.

CHAIN AND TRANSIT SURVEYS

Beginning at the 1971 diamond drill hole No. I on the Tam showing, the survey follows the main access road south, past the Camp and Coral showings to its junction with a road leading west to the Fire and Cliff showings. The survey then approximately follows this road past the Fire showing, terminating at the west side of the Cliff showing. Elevations were carried by both the vertical angle and stadia methods. The loop was closed using stadia and triangulated distances. A closing accuracy of 1:1200 was obtained. It should be noted that this represents a lower limit for accuracy that is probably much better within the chained portion of the loop.

Diamond drill holes located by chain and transit include all those drilled on the Tam showing during 1971 and all holes drilled on the Coral showing during 1972.

Triangulation was used to the distant claim posts, as well as the Nik and Bar showings. This was done from known points using short accurately chained baselines (\pm 0.01 feet) with all angles read by the repetition method to the instrument limitation of \pm 5 seconds of arc.

STADIA SURVEYS

Showings were surveyed by the stadia method to provide more or less evenly spaced stations of known elevation that would also serve to simplify geological mapping. These showings include the Tam, Camp, Coral, Fire, Cliff, Nik, Bar and Tee. Only the Tee showing is not tied to the main survey.

TOPOGRAPHIC CONTOURING

Topographic contour maps of surveyed showings were made in the field using spot station elevations and a clinometre. This method produced reliable contours without requiring a more detailed survey.

1- Cendel

A. J. Audet

October 10, 1972

.

-

		CO-ORDINATES		
STA	ELEVATION	NORTH	WEST	
1*	3625.5	79603.1	19323.2	
2*	3616.1	79603.1	19230.1	
3*	3609.8	79416.7	19051.0	
ц +	3600.9	79241.2	18 980.3	
5*	3587.3	79001. 5	18922.2	
6*	3576.9	78757.2	18973.4	
Claim Post •	3583.9			
7•	3587.2	78607.4	19222.7	
8•	3590.2	78462.3	19191.6	
9•	3623.4	78172.8	19363.5	
10*	3679.1	77960.4	1 9751. 9	
11•	3693.1	77779.5	19894.4	
11-1	3677.8			
11-2	3659.2			
11-3	36 68.6			
11-4	3645.8			
11-5	3633.4			
11-6	3664.7			
11-7	3668.9			
11-8	3674.5			
12*	3691.3	77417.5	20012.8	
13*	3671.6	77188.7	20285.7	
14•	3679.9	76969.3	20350.9	
15*	3690.9	76826.1	20393.5	
15-1	3695.5			
15-2	3695.7			
15-3	3694.7			
15-4	368 7.9			
16*	3685.9	76974.3	20637.6	
17	3684.8	76953.1	20880.6	
18 *	3680.3	77182.5	21030,4	
18-11	3685.7			
18 -1 2	3704.9			
18-13	3725.2			
18-14	3677.6			
18-15	3664.0			

COMMENTS

		CO-ORDI	MATES			
STA	RENVATION	HURTH	KEST	2	XNMENTS	
19	3697.4	77233.7	21277.1			
20	3681.1	77116.0	21 519.0			
21	3680,5	77131.0	21627.0			
22	3668.7	77230.3	21900. 5			
22-1 22-2	3665.8 3693.3					
22- 3	3670.7					
22-4	3694.4					
23	3700.8	77354.5	21914.7			
24	3726.2	77464.5	21823.3			
24-1	3714.1			•		
24-2	3712.8					
24-3	3738.1					
2 5	3755.7	77629.2	21647.4			
26	3746.0	77737.1	21555.1		·	
27*	3824.6	79572.9	20141.3			
28*	3821.6	79601.3	20119.1			
29•	3804.6	79604.9	20066.1			
30+	3862.9	79754.5	20142.9			
31*	3860.8	79860.8	20111.0			
31-1	383 3. 9					
31-2*	3825.0					
31-3*	3806.9			D.D.H.#	1971 - 4,5	
31-4	3772.6					
31-5	3 728. 6					
31- ó	3 6 69.6					
32*	3833.3	80000.0	20000.0	D.D.H.#	1971 - 1	
32 A •	3769.4					
32B •	3761.9			D.D.H.#	1971 - 6, 7	
320 •	3729.5					
32D •	3732.3			D.D.H.#	1971 - 8	
32E •	3680.0					
32-11*	3839.7					

ł

::

., ...

BL BR STOR

•

• •••

		TES	COMMENTS	
STA	<u>ELEVATION</u>	NORTH	WEST	
32-12 •	3826.1			D.D.H.# 1971-9,10,11,12
32-13	3826.1			
32-14	3771.6			
32- 15	3780.2			
32-16	3789.1			
32-17	3843.5			
33 •	3697.7	79453.6	19869.3	
33-1	3669.8			
33-2	3648.8	79 758, 5	19481.9	
34 *	3669.7			
34-1	3656.6			
34-2	3641.6			
35 •	3647.7	79583. 5	19664.0	
36•	3682.6			
36-1	3713.3			
36 -2	3686.7	· · ·		
37	3594.4			
38	3618.5			
39	3651.4			
40	3674.5			
41	3686.8			
42	3694.0			
43	3701.7			
44	3708.5			
44-1	3727.7			
44-2	3722.6			
45 •	3690.9	76513.4	20656.8	
46*	3675.3	76396.2	20949.2	
47•	3664.6	76300.7	21189.6	
48 *	3632.0	76004.7	21206.0	
49 •	3617.9	758 2 9.6	21380.7	
50.•	3600.6	75689.4	21272.5	•
51 *	3575-4	75524.8	21092.6	
52 •	3559.1	75398.4	21292.8	
53 •	3518.4	75280.1	21557.9	
54 *	3487.1	75112.6	21796.1	

÷.,

t

STA	ELEVATION	CO-ORDIN NORTH	ATES Wes <u>t</u>	COMPETTS
55*	3464.9	74952.6	21709.3	
56•	3433.2	74798.6	21432.3	
57•	3398.1	74489.3	21249.4	
58+	3369.9	74210.4	21297.7	,
59	2820.6			
60	2785.6			
61	2729.2			
62	2784.8			
63	2748.2			
64	2768.8			
65	2690.7			
66	2673.0			
67	2697.3			
68	2776.0			
69	2751.7			
70	2778.2		•	
71	272 8.7			
72	2737.3			
73	2805,2			
74	3622.4			
75	3634.2			
76	3634.9			
77	3679.6			
78	3688.8			
79*	2449.8	75504.5	31919.1	
80*	2438.6	75621.5	32141.0	
81 *	2442.0	75 623. 6	32252.0	
82	2420.3			
83	2403.1			
84	2357.2			
DDH#71-13	2422.1			
85	2358.6			
86	2337.7			
86-1	2340.1		DDH #71-14	

.

۰.

		NATES		
STA	ELEVATION	NORTH	WEST	COMMENTS
86-2	2363.3			
87	2311.9			
88	2347.1			
89	2326.4			
89-1	2318.8			
90•	2314.1			
90-1	5588° 8			
91•	2475.3	75299.4	31701.9	
9 2 •	2506.8	74956.3	31495.3	
93+	2566.0	74711.1	30793.2	
94 *	2578.5	74721.7	30504.7	
95•	2637.6	74925.3	29943.3	
96•	2711.0	74526.1	29403.3	
97•	2751.4	74294.9	28401.3	
98 •	2828.5	74225.2	27445.7	
98-1	2756.1			
98-2	2844, 3			
9 8- 3	2820.5			
08-4	2862.9			
09 •	2873.8	73682.8	26781.8	
99-1	2887.7			
) 9 - 2	2832.0			
99-3	2786.6			
99-4	2828.4			
99- 5	2866. 9			
100*	2 987.3	73878,8	26226.7	
101•	3104.7	74147.4	25 786, 9	
102*	3158.3	74334.1	24964.5	
103*	3199.3	74396.8	24422.8	
104•	3246.9	74412.7	23758.7	
105*	3265.5	74410.1	23238.7	
106•	3272.4	74349.4	22445.7	
107*	3302.5	74244.8	2 1 93 2.2	
108*	3299.2	73998.1	21663.2	
109•	3327.5	73462.7	21331,9	

.

.

· .

.

.

ŝ

ł

•	STA	ELEVATION	CO-ORDINA NORTH	TES WEST	CONNENTS
	110 •	3328.3	73876.1	21361.4	
	133	3681.8			
	133-1	364 1.3			
	133-2	3680,6			
	133-3	3678. 9			
	134	3699.4			
	135	3700.2			
	135-1	3687.5			
	135-2	36 71.9			
	135-3	3671.6			
	135-4	3702.0			
	135-5	3700.7			
	136	3691.7			
	136-1	3675.0			
	136-2	3646.7			
	137	3574.6			
-	137-1	3567.6			
	137 -2	3535.5			
	138	3544.3			
	139	3519.3			
	139-1	3 523. 8			
	140	3527.0			
	141	3499.1			
	142	3466.7			
	142-1	3434.2			
	143	3540.1			
	144	3560.9			
	145	3577.2			
	146	2824.6			
	146-1	2832.1			
	146-2	2851.6			
	146-3	2839.0			
	146-4	2814.6			
	147	285 7. 8			
	147-1	2846.1			

,

-- -----

		CO-ORDI	NATES	' 3
STA	ELEVATION	NORTH	WEST	COMMENTS
147-2	2880, 3			
147-3	2905.6			
147-4	2924.5			
148	2977.4			
148-1	2963.0			
148-2	2961.6		· ·	
149	2978.9			
149-1	2982.9			
149-2	2945.4			
150	3682.8			
150-1	3684.2			
151	3677.1			
152	3744.1			Triangulation sta NW of Coral Prospect
153	3741.6	77518.2	21998.5	
154	3716.7			
155	3665.1			
156	3703.5			
157	3702.4			
158	3694.8			a
158-1	3682.5			
159		64765.4	25527.9	Triangulation point 980'SE of AP#1
160				Triangulation Sta. above Bar Show-
161	2838.9	62521.4	21139.9	Triangulation Sta.at the Bar Show-
161-1	2842.7			ing
161-2	2852.0			
161-3	2794.8			
161-4	2787.0			· · · · · · · · · · · · · · · · · · ·
161-5	2816.2			
161-6	2817.7			
161-7	2812.4			
161-8	2 452.6			
161-9	2871.4	-		
1.62		66372.4	22271.9	Triangulation Sta.at sharp turn in
163				150' SW of 162 Bar Rd.
164				260' west of 162
165		۰.		Triangulation Sta. north of Sta.96

. . .

.

-

and an and a second second

ъ .

STA	ELEVATION	CO-ORDIN <u>NORTH</u>	ATES WEST	COMMENTS
166	2572.8	77588.1	33851.3	Triangulation Sta. at Nik Showing
166-1	2524.2			
166-2	2546.3			·
166-3	2547.0			
166-4	2560.6			
166-5	2558.1			
167	2 578.9	776 99. 1	34005.3	,
168	2566.9	77572.1	34022.3	
168-1	2554.0			
168-2	2547.5			
169	2571.9	77750.1	34058.3	
169-1	2548.6			
169-2	2552.7			
170	2546.9	77471.1	33873.3	
171	2490.3	77183.1	341.36.3	
171-1	2477.8			
171-2	2506.8			
172	2782.7			
172-1	2791.6			
172-2	2778.1			
172-3	27 59.6			
173	2913.4			
173-1	2898.9			
173-2	2936.0	т. Г		
173-3	2955.6			
173 - 4	2920.1			
173-5	2903.1			
173-6	2906.2			
173-7	2883.2			
174	2745.1			
175	2720.5			
175-1	2727.0			
175-2	2726.9			
175-3	2763.4			

and the second

.

•

STA	ELEVATION	CO-ORDIN <u>Horth</u>	ATES WEST	COMPANY
A.P. #1		65153.7	26447.4	
J-7	3691.4			
J-9	3676.9			
J-10	3672.6			
J-11	3676.6			
J-13	3688.9			
J-14	368 7.8			
J-15	3685.9			·
J-16	3692.5			
J-1 8	3697.3			
J-19	3697.4			
J- 2 0	3701.1			
T 1	2750.0			
T 2	2798.6			
T 3	2843.9			
T 3-1	2866.6			
T 3-2	2901.6			
T 3-3	2915.2			
T 4	2942.0			
T 4-1	2986.2			
T 4-2	2985.4			
T 5	2976.1			
T 5-1	2981.8			
T 5-2	3044.1			
T 5-3	3079.6			
T 6	3032.3			
T 6-1	3065.1			
T 7	3098.6			
т 8	3165.9			
т 8-1	3176.2			
т 8-2	3113.8			
T 8-3	3 22 8.8			
T 9	3222.4			
T 9-1	318 2.0			
T 9-2	3152.7			

•

17 W

CO-ORDINATES

<u>STA</u>	ELEVATION	NORTH	MEST	CORRECTS
T 9-3	3160.3			
T 9-4	3244.5			
T 10	3271.7			
T 10-1	3280.1			
T 10-2	3314.6			
т 10-3	3318. 2			
T 11	3310.9			
T 11-1	3339.1			
T 11-2	3366.4			
T 11-3	3384.2			
T 12	3367.1			
Ŧ 12-1	3407.6			
T 13	3410.8			
T 13-1	3403.5			
T 13-2	3376.4			
т 14	3330.1			
T 14-1	3315.2			
т 14-2	3259.1			
T 15	3325.6			
T 15-1	3284.6			
Ŧ 15-2	3267.7			
T 1 5-3	3270.2			
T 16	3247.3			
T 16-1	3207.9			
T 17	3201.5			
T 17-1	3142.8			
T 18	3203.0			
T 18-1	3159. 2			
T 18-2	3160.3			
T 18-3	3157.9			
T 19	3150.7			
T 19-1	3110.9			

CLAI	<u>M NAME</u>	RECORD NO. or CROWN GRANT NO. *	DATE RECORDED
GEM	# 1	6700 CG	
**	2	6701 CG	
**	3	6 702 C G	
**	5	6703 CG	
**	6	67 04 C G	
**	7	6 705 C G	
GEM	# 2 Fr.	6 706 CG	
Ħ	3 Fr.	6707 C G	
**	4 Fr.	6708 CG	
•1	8	6709 C G	
GEM	# 5 Fr.	6710 CG	
71	4	6711 CG	
GEM	# 1 Fr.	6712 CG	
•1	9	6713 G	
n	14	67 1 4 o g	
• 1	15	6715 C	
**	11	6716 C G	
11	12	6717 CC	
9†	13	6718 CG	
11	19	6719 C G	
**	16	6720 CG	
11	17	67 21 C G	
*1	25	67 22 C G	
**	23	6723 CG	
f#	24	6724 CG	
*1	2 5	6725 CG	
11	21	67 2 6 C G	
Lea	Fr.	67 27 C G	
Mem to	# 1	49 392 to	September 15, 1970
Mem	# 8	49399	September 15, 1970
GEM	# 2 6	51875	July 23, 1971
to GEM	# 41	to 51890	July 23, 1971
GEM	# 42	55318	September 2, 1971
to GEM	# 58	to 55 33 4	September 2, 1971
		7703 ⁴	way of a second se

·

. . . .

Page 2

· • •

CIAIM	NA	ME		RECORD NO.	or CR	WN GRANT NO. *	DATE RECORDED
	#	68			56544		October 7, 1971
to GEM	#	76			to 565 52		October 7, 1971
	#	77			56766 to		October 7, 1971
to GEM	#	83			567 72		October 7, 1971
GEM to	#	59			57798 to		December 20, 1971
	#	67			5 780 6		December 20, 1971
Pet to	#	1			56953 to		October 20, 1971
	#	100			57052		October 20, 1971
Fall Fall	##	ן 26	to		56 482 56 507	to	October 7, 1971 October 7, 1971
Fall Fall	4 #	27 32	to		565 22 565 27	to	October 7, 1971 October 7, 1971
Fall Fall	##	33 46	to		56508 56521	to	October 7, 1971 October 7, 1971
Fall Fall	# #	47 52	to		565 2 8 565 33	to	October 7, 1971 October 7, 1971
Fall Fall		53 56	to		56 534 565 37	to	October 7, 1971 October 7, 1971
Fall Fall	##	57 6 2	to		565 3 8 56543	to	October 7, 1971 October 7, 1971
Fall Fall	# #		to		57618 57631	to	October 25, 1971 October 25, 1971
Fall Fall			to		576 32 57647		October 25, 1971 October 25, 1971
Fall Fall	# #	95 97	to		57713 57715		October 28, 1971 October 28, 1971
Fall Fall			to		57716 577 22	to	October 28, 1971 October 28, 1971
Pall	#	10 5			57723		October 28, 1971
Fall					577 2 4		October 28, 1971
Fire Fire			to		56746 56765	to	October 7, 1971 October 7, 1971
Fire Fire			to		56609 56614		October 7, 1971 October 7, 1971
Fire Fire			to		56661 56682		October 7, 1971 October 7, 1971
Fire Fire	#	57	to		56683 566 91		October 7, 1971 October 7, 1971
Fire		-			56615		October 7, 1971
Fire					56616		October 7, 1971

OG refers to Crown Grant

.....**S**

Page 3

CLAIM	NAM	Ē		RECORD NO. or CROWN GRANT NO. *	DATE RECORDED
Fire	# (58		56617	October 7, 1971
Fire	# (59		56618	October 7, 1971
Fire Fire			to	566 92 to 56697	October 7, 1971 October 7, 1971
Fire	# 7	76		56698	October 7, 1971
Fire Fire	# 1 # 1	17 t 19	0	566 99 to 567 01	October 7, 1971 October 7, 1971
Fire Fire			to	56702 to 56722	October 7, 1971 October 7, 1971
Fire Fire			to	56723 to 56731	October 7, 1971 October 7, 1971
	# #	1 2 6	to	57075 to 57100	October 19, 1971 October 19, 1971
	# #	27 54	to	57534 to 57561	October 25, 1971 October 25, 1971
	# #	1 4	to	517 3 9 to 517 42	July 26, 1971 July 26, 1971
Tam Tam	# #	5 32	to	51743 to 51770	July 23, 1971 July 23, 1971
	# #	33 84	to	51771 to 518 22	July 23, 1971 July 23, 1971
Tam Tam	# { # :	35 151	to	55 33 5 to 55401	September 2, 1971 September 2, 1971
Tee Tee	# #	1 52	to	518 23 to 51874	July 23, 1971 July 23, 1971
Tee Tee	# #	53 70	to	56643 to 56660	October 7, 1971 October 7, 1971
Tee Tee	##	7 1 84	to	567 32 to 56745	October 7, 1971 October 7, 1971
T ee T ee	# #	85 140	to	57562 to 57617	October 25, 1971 October 25, 1971
West West	# #	1 24	to	56619 to 56642	October 7, 1971 October 7, 1971

*CG refers to Crown Grant

Page 4

· · · ,

. . .


ł


.


CLAIN	NA	ME		RECORD NO. or CROWN GRANT NO. *	DATE J	RECORDED
West West	##	25 80	to	56553 to 56608	Octob er ' Octob er '	
West West	##	81 96	to	57101 to 5 711 6	October : October :	
West West	# #	97 102	to	57528 to 57533	October 2 October 2	.,
West West	# #	103 128	to	57663 to 57688	October : October :	

.













.














A 155

A 46

A 47

& 48

44



------ 76,000 N -------

- 80,000 N -------


----- 72,000 N-----

∆95

FIRE

.

A 92



70 71 FIRE

- 68,000 N----

<u>A</u> 164 163 A 162

•

A AP-1

A 159

------ 64,000 N -------

.

• • . -·· -----. •

.


-

. •

To accompany Report by J. R. Woodcock (November 1972)

•


on Liard Fluorspar Project Tee, Tam, Fire, West, Gem Mineral Claims Liard Mining Division



· · · · ·

.

. Department of



.

∆108 — Survey station - Wooden Hubs marked with numbered aluminum tags


LEGEND


🛛 — Claim post


.


.


LIARD FLUORSPAR MINES LTD.












4.5 12 27,000 Weg To accompany Report by Z. R. Hoodcock (Revention: 1972) on Mard Finorepur Project Saw, Dan, Fire, West, Gen Mineral Ciaina Itari Mining Division


CONWEST EXPLORATION COMPANY LTD.

LIARD RIVER FLUORSPAR PROJECT P-1021

<u>> ₩</u>

Success on joint surfaces :





LIARD FLUORITE

10.0

14

.

.

100

1

SHOWS A, B, D, E, G

1.00

122

G SHOWING D

.

111

1.1

SCALE 1'-100' CONTOUR INTERVAL 10 Fr

LEGEND

C DRE-(a) in outerop, Wassamed (are has ever 60% combined fiverity witherity and parts)

B SLATE & HAGILLITE () IN SURCESH, (Dessamed

A LINTESTONE W = outcrop, Dassames

Cliff edge

Roch contains small Jenses b stringers of are


Float of ore F Foreil Locality

P Rillings of Bedding


Stripped Reea


Grid Ling sugarage


- Grid Ling supering level ar al industried 2- high page














To accordingly Debout J. R. Moodcools (Hovestver 1972)


#1

100

A.S.

2

Lined Finnesper Project Toe, Tun, Fire, Vest, Gen Miseral Claims Mard Hising Division



.

.

.

.

\~

							· · · · · · · · · · · · · · · · · · ·
	••••••••••••••••••••••••••••••••••••••		-	<u> </u>	·		
· · · · · · · · · · · · · · · · · · ·							•
•							
					•	· -	
							•
							•
		-				W	
							•

• . • •


· · -

• • . .

•

٠ -· · · •


•




•

-

•





Tee, Tam, Fire, West, Gem Mineral Claims Liard Mining Division

