

5509

GEOLOGICAL & GEOCHEMICAL REPORT

by

T.D. Pearse, B. Sc.

104G/14W

on surveys completed during the period

July-August 1974

on the

KIT Claim Group

situated on

Winter Creek, Telegraph Creek area

in the

Liard Mining Division

(57°55'N 131°25'W)

(NTS 104G/14W)

and owned by

ECSTALL MINING LTD.

October 1974

Department of

Vancouver, B.C.

Mines and Petroleum Resources

ASSESSMENT REPORT

NO. 5509 MAP

TABLE OF CONTENTS

INTRODUCTION AND SUMMARY

PREVIOUS WORK

CLAIMS AND OWNERSHIP

LOCATION AND ACCESS

PHYSIOGRAPHY

GEOLOGICAL SETTING

DETAILED GEOLOGY

Lithologic Units
Alteration
Mineralization
Structure

GEOCHEMISTRY

Discussion of Field Methods
Laboratory Determination Method
Discussion of Results

APPENDIX A: Geochemical Data

APPENDIX B: Cost Statement

APPENDIX C: Statement of Qualifications

LIST OF ILLUSTRATIONS

- | | | |
|---|-------------------------------------|------------|
| 1 | Index Map, 1: 50,000 (LOCATION MAP) | after p. 3 |
| 2 | Geological Map, 1" = 400' | in pocket |
| 3 | Geochemical Map, 1" = 400' | in pocket |
| 4 | Topographic Map, 1" = 400' | in pocket |

INTRODUCTION & SUMMARY

The Winter Creek property, in the Stikine River area of northwestern B.C., consists of 26 claims overlying a steeply-dipping, undifferentiated sequence of upper Triassic volcanic rocks. Copper mineralization occurs sparsely but widely distributed within augite/feldspar porphyries. Occurrences are of two types: blebs and smears of chalcopyrite along serpentized shear surfaces, and; massive chalcopyrite with pyrrhotite in two small lenticular sulphide replacement bodies. Subsequent to staking, work consisted of prospecting, geologic mapping (1" = 400 scale), and a talus fines sampling program. The geochemistry indicates a high background for copper in the order of 200 - 300 ppm. with some local values running to 2,800 ppm. Molybdenum concentrations range from a background of 2-3 ppm. to a high of 27 ppm. No molybdenum mineralization was observed on the property. Geochemistry and geology indicate a broad, undefined zone of sparse copper-sulphide mineralization throughout the western half of the property. The eastern half of the property contains no significant copper showings. It is the writer's belief that observed modes and concentrations of copper occurrences could account for the high geochemical values obtained. It appears that no concentrating mechanism has been active in concentrating sulphide mineralization into economically interesting amounts and thus the property's economic potential is low. It is recommended that no further work be accorded this claim group.

PREVIOUS WORK

In 1917, a discovery of copper mineralization was made on the south-facing slope of the Winter Creek Valley at an elevation of about 5,000'. This consisted of a small massive sulphide body containing varying amounts of pyrrhotite and chalcopyrite, a sample of which assayed 0.12 oz/ton Au, 0.92 oz/ton Ag., and 5.8% Cu. A description of this occurrence is given by J.D. Mandy in G.S.C. Memoir 246, Lower Stikine and Western Iskut Areas, British Columbia, p. 75. Claims covering the area have been staked and abandoned several times since and include, for example, the Glenora and King Groups of 1929, and the NP Group of 1962. An extremely recent campsite on the small tarn north of the ridge indicates that an exploration team was active in the area probably in the 1973 field season --- apparently no claims were staked at this time. On the 6th. of July, 1974, four men employed by Ecstall Mining Ltd. staked KIT Nos. 1 to 22; KIT NOS. 23 to 26 were added August 26th., 1974. Subsequent to the initial staking,

two men spent approximately forty-two man-days actively engaged in a geologic assessment of the area's copper-bearing potential.

CLAIMS & OWNERSHIP

The property consists of twenty-six contiguous mineral claims in the Liard Mining Division of B.C. The claims are owned by Ecstall Mining Ltd., and are listed as follows:

<u>CLAIMS</u>	<u>TAG NOS.</u>	<u>RECORDING DATES</u>
KIT #1 to 22	352701 M - 352722 M	July 15th., 1974
KIT #23 to 26	352724 M - 352727 M	Sept. 25th., 1974

LOCATION & ACCESS

The Winter Creek property is located approximately 10 miles west of Telegraph Creek, B.C., at 57° 55' N lat. and 131° 25' W. long. (see Index Map). It is accessible by foot along packtrails from Telegraph Creek, or by helicopter from Iskut B.C., approximately 55 miles to the east on the Stewart-Cassiar highway.

PHYSIOGRAPHY

The KIT Group lies along the eastern contact of the Coast Mountain Area and the Central Plateau-Mountain Area of the Interior system of the Canadian Cordillera. It is located on the east flank of the Coast Mountains at elevations of 3,500' to 6,300' and consists of a single, rugged, east-west trending ridge bounded to the north and south by two trunk valleys opening to the east, and to the west by Grass Mountain. To the south of the property, Winter Creek flows steeply east to its confluence with the Stikine River; a small unnamed tributary of Dodjatin Creek drains the north half of the map-area. The floors of Winter Creek Valley are forested with balsam, spruce and other alpine scrub species to about 3,800' grading into buckbrush slide alder, heather and grasses higher up on the valley slopes. Ninety-five percent of the outcrop is above 4,500' and this marked by a pronounced steepening in slope from the vegetated slopes below. Both the north and south faces of the ridge are deeply incised by erosional gulleys and characterized by precipitous topography that makes detailed traversing hazardous.

GEOLOGIC SETTING

The Winter Creek property is underlain by a sequence of undifferentiated intermediate volcanic rocks of upper Triassic age. These are predominantly augite andesite breccia, conglomerate, and volcanic sandstone, but include thick sections of greywacke, siltstone, tuff and minor shale. This sequence of rocks lies along the north flank of the northeast trending Stikine Arch, a crystalline complex that remained relatively positive throughout Mesozoic time and which greatly influenced the style of deposition of great thicknesses of volcanic and clastic sedimentary rocks during this time. The main axis of the Arch lies approximately 20-30 miles to the south of the property.

Immediately to the north of the claim group a small, double-lobed intrusion of quartz-deficient, intermediate to basic rocks of post upper Triassic age outcrops. The main axis of this body is approximately six miles in length and parallels the Stikine Arch Axis. There are several intrusive phases which exhibit gradational or complex migmatitic contacts with each other. Contacts with the intruded volcanic rocks, however, are sharp with relatively little contact metamorphism.

DETAILED GEOLOGY

Lithologic Units

The bulk of the rock types underlying the Winter Creek property comprise intermediate volcanic porphyries of varying compositions, textures, and relative age. Minor nonporphyritic lithologies include rhyolitic-dacitic tuffs, massive pyritiferous andesite, syenite, and monzonite. The rocks are described below in order of their relative abundance:

aP:

Augite Porphyry: a medium-dark grey porphyritic andesite with euhedral augite phenocrysts (up to several cm. in length) in an aphanitic matrix. Crowded porphyry textures with a concomitant decrease in phenocryst size are common. Minor concentrations of euhedral plagioclase phenocrysts or rare blebs of intergranular pyrite may be present. With an increase in pyrite content this rock grades into massive, nonporphyritic andesite.

FP:

Feldspar Porphyry: a medium-dark grey porphyritic andesite containing varying concentrations of fine-medium grained, euhedral plagioclase phenocrysts in a dark, aphanitic matrix. Phenocrysts may range up to 3 mm. in length and locally may exhibit trachytoid texture. Augite phenocrysts may be present in minor quantities.

H:


Hybrid Porphyry: a transitional member between aP and fP characterized by either approximately equal concentrations of plagioclase and augite phenocrysts in the same groundmass or by poorly defined and/or complex zones of mixing between the two rock types. The augite/feldspar porphyries are gradational and have been distinguished in the field on the basis of which phenocrysts are predominant. Unit H, therefore, represents a hybrid zone where distinction between aP and fP was difficult or not practicable.

RT:

Rhyolitic Tuff: an aphanitic felsic rock generally intensely fractured in surface exposures. It contains fine grains of pyrite which oxidize extensively to limonite and thus well-developed stain zones earmark the occurrences of this unit. Classification as a rhyolite is tentative at best because of the difficulty of obtaining fresh rock surfaces. On the few such surfaces observed, however, this unit exhibited poorly-defined, thin banding; this coupled with its well-bedded occurrences suggests it represents a pyroclastic layer of siliceous composition. On the north-east side of the property, this unit occurs in several bands that are wide-ranging and continuous in extent and which may reach thicknesses of several tens of feet. Towards the west end of the property these rocks are restricted in occurrence to small crescentic lenses and "stringers" and are commonly fracture-filled with a cemented quartz-carbonate matrix. It is with these latter occurrences that the two small massive sulphide lenses exist. Locally, quartz-eye phenocrysts have been distinguished.

dT:

Dacitic Tuff: an aphanitic felsic rock of minor occurrence, less siliceous and darker in colour than the rhyolitic tuff.



Locally, it is distinctly more clastic in texture than rT and disseminated pyrite was commonly observed. Epidote in fractures is present to a lesser degree.

M:

Monzonite (plus diorite): a fine-grained equigranular rock with less than 10% quartz, 10-15% hornblende, and 75-80% consisting of an even mixture of K-feldspar and plagioclase. Locally, this rock grades into: a) diorite, wherein the abundance of plagioclase increases significantly and; b) mafic syenite, wherein the concentration of K-feldspar and hornblende increase to the point at which plagioclase and quartz comprise minor constituents of the rock. Generally, the monzonite is fresh in appearance, although the K-feldspars may be bleached on the weathered surfaces. Alteration is extremely weak or absent (minor epidote and K-feldspar along hairline fractures or in patches) and the rocks are not strongly fractured or deformed. Accessory minerals include apatite and magnetite.

S:

Syenite: a light pink to grey, fine-grained equigranular syenite which occurs as dyke-like bodies within the monzonite and the volcanic units. These bodies are all extensively fractured and pervasively altered through quartz-carbonate flooding --- in very few instances is a primary magmatic character distinguishable. Mafic content is low; the rock is composed primarily of K-feldspar with minor quantities of quartz.

QC:

Quartz-carbonate rock: small lenticular occurrences of highly fractured, Fe-stained rock in which the only recognizable constituents are limonite, quartz, and carbonate. This assemblage apparently occurs with both the felsic volcanics and the syenite and associations are generally difficult to make in the field. These bodies range from 10' - 100' long and several feet thick.

Alteration

Alteration types present in the Winter Creek rocks include, in order of relative predominance, chloritization, serpentization, quartz-carbonate alteration, and minor occurrences of epidote and gypsum. Chloritization and

serpentinization are the most widespread; the former occurs both as propylitization of primary pyroxenes in the augite porphyries and with serpentine along shear surfaces. Serpentine development is restricted to shear surface coatings and slickensides in weakly to intensely sheared porphyritic rocks. Commonly the most fractured and altered zones are shot full of fine felsic stringers. Pods of quartz-carbonate rock are present in several locations usually in conjunction with felsic volcanics (at least in the eastern end of the property), although this relationship is not always distinguishable. These bodies of intensely-altered rock have been more substantially described under Section (1) above as they were mapped as a separate rock unit. Other alteration minerals include rare epidote, hematite, gypsum and actinolite-tremolite, all located along hairline fractures in the volcanics and none with any significant continuity. Small, uncommon veinlets of epidote were also observed in the monzonite/diorite rocks to the north.

Alteration in the intrusive rocks along the north edge of the claim group is restricted to limited development of pink, K-feldspathized veinlets and patches, and the thin veinlets of epidote as mentioned above. Generally, the intrusive rocks are unaltered and extremely fresh-looking on the broken surface.

Chloritization and serpentinization are ubiquitous alteration modes in the volcanics; other alterations occur locally and are generally not intensive. There appears to be no significant alteration pattern associated with any of the processes.

Mineralization

Sulphide minerals observed on the property include pyrite, chalcopyrite, and pyrrhotite. Economic mineralization occurs in two distinct forms: 1) fracture-fillings of chalcopyrite in the volcanics, and; 2) pyrrhotite and chalcopyrite in two small massive sulphide lenses. Approximately one dozen small localities within the west half of the claim group carry minor quantities of blebby chalcopyrite along heavily serpentinized shear surfaces in both the porphyritic and pyritiferous, massive andesites. These are all localized in extent (several feet along strike, one or two inches thick) and mineralization is extremely erratic and inconsistent throughout. Moderately well-developed malachite staining results from these sparse sulphide occurrences rendering them easily observable

from a distance. These showings are apparently related to a late-stage fracturing/shear system parallel or semi-parallel the regional trend of the volcanic units.

Chalcopyrite in association with pyrrhotite occurs also in two thin (upto 6" wide) massive sulphide lenses which occurs along the upper contact zone of the massive andesite and the porphyritic member. The western showing occurs along a well-developed shear zone and strikes southerly with a vertical dip: exposure is erratic and offset over a length of approximately 100'. The eastern showing strikes southerly, dipping 65° East, and is exposed for approximately 40' with widths up to 6". Two rock-chip samples (WCTP-5,6) were taken on the western massive sulphide exposure and these assayed:

- 1) Au, 0.20 oz/ton; Ag, 0.41 oz/ton; Cu, 4.25%
- 2) Au, 0.06 oz/ton; Ag, 0.06 oz/ton; Cu, 0.83%

Structure

The volcanic rocks underlying the KIT group apparently comprise a thick sequence of steeply-dipping, interbedded rhyolites and andesites, striking approximately ENE and dipping to the south. Some kind of major structural element breaks the stratigraphic continuity between the east half of the property and the west. To the west the rocks consist of thick, undifferentiated augite/feldspar porphyries with a thick (up to 200') interbedded, pyritiferous unit which is commonly massive and non-porphyrific. The oxidation of pyrite in this unit weathers to yield a distinctive stain zone across the south slope of the ridge. Bedding attitudes were impossible to measure in the field due to the extensive fracturing and shearing that has occurred. However, a best-fit approximation of stratigraphy has been derived from the outcrop pattern of the pyritiferous-member. In addition, a prominent S1 surface is developed in outcrop which parallels the suggested stratigraphy and lends support to the model. The pyritiferous horizon is offset by faulting in several locations and the fault zones are marked by gouge and brecciation over several tens of feet. One of these shear zones contains the western-most massive sulphide occurrence which is apparently associated with felsic volcanics. Smaller fault surfaces with brecciation and slickensides are common throughout the entire sequence. It is these faults and prominent fractures that contain the observed copper mineralization.

The major faults appear to offset slightly from the strike of the stratigraphy, but exhibit the same steep dip to the south. In the most southwesterly outcrops a series of segmented and disoriented dykes are exposed. These are all vertically dipping and consist of augite porphyries, some with large quartz lenses up to six inches and locally comprising 50% of the rock by volume. Also within the pyritiferous horizon an interbedded rhyolitic member is exposed; its extent is unknown. One rock geochemical sample from this locality was taken (WCTP4), yielding 178 ppm. Cu and 1 ppm. Mo. Along the ridgetop several small bodies of felsitic volcanics and one small syenite dyke are exposed, contact relations with the andesites are obscured and indeterminate.

In the northeast corner of the map area stratigraphy is more obvious due to prominent and well-defined interbedded layers of felsitic volcanics and intermediate volcanics. At least ten distinctive bands of rhyolitic tuff ranging up to 40' thick are exposed on the north face. These exhibit strikes to the east and east-north-east and moderate dips to the south. Outcrop patterns and bedding suggest this sequence has been gently folded around a north-south fold axis plunging 20° - 40° to the south. Some small faults and a prominent ENE fracture set were observed. No copper-sulphide mineralization was observed on the east half of the claim group.

To the north of the property the volcanics are intruded by the monzonitic intrusion. This pluton is apparently concordant with the regional structure of the country rocks. Contacts are abrupt with relatively little contact effect, but may exhibit complex textures where intermixing of volcanic and plutonic rocks has occurred. For example, dyking of both rock types into the other is evident; small zones of augite porphyry with monzonitic inclusions and thin intrusive dykes with fragments of volcanic wallrock can be distinguished. This small stock comprises three differentiated phases; a basic dioritic original phase, a monzonitic younger phase, and a late phase of syenite dyking. Late stage volcanic dyking was apparently going on during the initial plutonic intrusive phases. The plutonic rocks, however, are all structurally undeformed and insignificantly altered. They are barren of sulphide mineralization and it is difficult to envision them as source potential for the volcanic copper occurrences.

GEOCHEMISTRY

Discussion of Field Methods

Geochemical work to date consists of 89 samples of talus fine material and 7 rock chip samples. Two lines of talus sampling were run horizontally across the south face at elevations of approximately 5,400' and 4,500'. The lower line was located topographically below the pyritiferous andesite horizon and the upper line above it. Sample stations were chained and established every 200' by flagged markers. An altimeter was used to maintain vertical control. In addition, a shorter line of 9 samples was run, before the claims were staked, between the two lines described above. Also, two samples were subsequently taken north of the main ridge. Approximately four tablespoons of the finest talus material were collected at each site and placed in numbered, 3½" x 6-1/8" ' Open End ' Kraft envelopes. All samples were analyzed for copper and molybdenum in the Bondar-Clegg Laboratory located at 1500 Pemberton Ave., N. Vancouver, B.C.

Seven rock chip samples were secured by taking 20-25 small chips in a randomly distributed manner over a width of approximately 20'. These samples were collected in plastic bags and sent to Bondar-Clegg for Cu. and Mo. analysis; the results are listed in Appendix A.

Laboratory Determination Method

The samples were first separated to a -80 mesh fraction. This process required crushing in the case of rock samples. Combined metal was extracted from a weighed sample of this fraction with Le Fort aqua regia. The resulting solutions were bulked to a 20% acid concentration and analyzed by atomic absorption spectrophotometry, in constant comparison with both synthetic and matrix standards. Results are expressed in parts per million contained metal.

Discussion of Results

Values for copper in the talus fines sampling range from 115 ppm. to 2,800 ppm. with a calculated arithmetic mean at approximately 504 ppm., and median at 370 ppm. Cu. Molybdenum values range from nil to 27 ppm. with a background of 2-3 ppm. Only a very general correlation between Cu and Mo values is revealed; i.e. high Mo concentrations were

generally associated with higher Cu concentrations, but exceptions are common with correlations being erratic and inconsistent.

Observed mineralization could account for the anomalous and high background concentrations of copper. The anomalous Mo values are more difficult to explain as Mo mineralization was unobserved on the property. The results do shown that this sequence of volcanic rocks does have an interestingly high copper background, but that no real pattern of concentration is revealed.

T. J. Pearce
M. J. Maxwell

GEOCHEMICAL DATA SHEET — SOIL SAMPLING

SAMPLER IC/GRP

NTS 1046/14W

DATE 11 June 1974

PROJECT Winter Creek (08)

LINE _____

AIR PHOTO No. TK 5158-165

SAMPLE No.	LOCATION	DEPTH	HORIZ.	DESCRIPTION				SLOPE	VEG.	ADDITIONAL OBSERVATIONS OR REMARKS	ASSAYS					
				COLOUR	PART. SIZE	% ORG.	PH				Cu	Mo	Zn	Pb		
3348		6"	na.	reddish	coarse	—		35°	none	near west edge of one stain	317	3				
49		"	"	"	"	—		"	"	100' east	520	2				
50		"	"	"	"	—		"	"	200' "	450	4				
51		"	"	dk. brn.	"	5%?		"	grass	300' "	415	5				
52		"	"	"	"	—		"	"	400' "	225	6				
53		"	"	reddish	"	—		"	none	500' "	575	6				
54		"	"	"	"	—		"	"	600' "	1700	25				
55		"	"	"	"	—		"	"	700' "	330	2				
3356		"	"	"	"	—		"	"	800' "	1700	27				
										taken ± on 5000' contour, 100' spacing; mostly talus fines.						

GEOCHEMICAL DATA SHEET — SOIL SAMPLING

SAMPLER Pearse/CooperNTS 104G/14WDATE 4 July, 1974PROJECT Winter Creek (08)

LINE _____

AIR PHOTO No. BC 5358-165

SAMPLE No.	LOCATION	DEPTH	HORIZ	DESCRIPTION				SLOPE	VEG.	ADDITIONAL OBSERVATIONS OR REMARKS	ASSAYS				
				COLOUR	PART. SIZE	% ORG.	PH				Cu	Mo	Zn	Pb	
K3063	Cairn above camp	4"	C	m-dk brn	soil	low		mod	grass	Samples on 4500' contour	361	1			
64	200' W.	"	"	"	"	"		steep	willow	andes. tuff bedrock	629	7			
65	400' "	5ft	"	"	"	mod		mod	"	" " "	405	4			
66	600' "	"	"	"	"	low		steep	bk-brn	andes. fldsp. porph.	545	2			
67	800' "	"	"	"	rk chips	nil		"	"	" " "	382	1			
68	1000' W	"	"	"	clips of soil	nil		"	heather	" " "	300	3			
69	1200' "	"	"	"	"	"		"	"	" " "	296	3			
70	1400' "	"	"	lt. brn	"	"		"	nil	" " "	250	2			
71	1600' "	"	"	"	"	"		"	grass	" " "	560	3			
72	1800' "	"	"	"	"	"		"	nil	" ? "	540	6			
73	2000' "	"	"	"	"	"		"	"	" ? "	635	4			
74	2200' "	"	"	"	"	"		"	"	" " "	645	3			
75	2400' "	"	"	"	"	"		"	"	" " "	1880	4			
76	2600' "	"	"	"	"	"		"	"	Samples on 5400' contour	130	2			
77	900' vert above 76	"	"	red brn.	chips	nil		"	"	gossan w. malachite	960	1			
78	700' E of 77	"	"	"	"	"		"	"	some stain, just above gossan	910	7			
79	400' E	"	"	lt. brn.	"	"		"	"	augite porph	140	13			
80	600' E	"	"	"	"	"		"	"	" " "	650	12			
81	800' E	"	"	"	"	"		"	"	" " "	775	6			
K3082	1000' E	"	"	"	"	"		"	"	" " "	615	5			

GEOCHEMICAL DATA SHEET - SOIL SAMPLING

SAMPLER Pearse/CooperNTS 1046/14WDATE 4-5 July, 1974PROJECT Winter Creek (08)

LINE _____

AIR PHOTO No. 8C 5158-165

SAMPLE No.	LOCATION	DEPTH	HORIZ	DESCRIPTION				SLOPE	VEG.	ADDITIONAL OBSERVATIONS OR REMARKS	ASSAYS					
				COLOUR	PART. SIZE	% ORG.	PH				Cu	Mo	Zn	Pb		
K3083	1700' E of 17	sfc	C	lt. brn.	chips	nil		steep	grass	Samples @ 5500' contour.	480	4				
84	1400'	"	"	"	"	"		"	nil	flds. porph. b/r.	659	3				
85	1600'	"	"	"	chips + soil	"		"	grass	" " "	300	3				
86	1800'	"	"	"	"	"		"	nil	gully	371	4				
87	2000'	"	"	"	"	"		"	"	flds. porph.	810	7				
88	2200'	"	"	"	"	"		"	grass	aug porph.	585	6				
89	2400'	"	"	"	"	"		"	nil	"	1350	19				
90	2600'	"	"	"	"	"		"	"	flds. porph. (cpy in flt 100' W)	270	3				
91	2800'	"	"	"	"	"		"	grass	"	350	9				
92	3000'	"	"	"	"	"		"	"	"	320	3				
93	3700'	"	"	"	"	"		"	shrubs	elev. 5520'	308	14				
July 6 94	200' E of K3083	"	"	"	"	low		"	alder	" 4440' (as cairn @ K3083)	430	4				
95	400' E "	"	"	"	"	nil		"	nil	" 4400'	383	3				
96	600' "	"	"	"	"	"		"	"	"	700	3				
97	800' "	"	"	rd. brn.	"	"		"	"	" 4470' (in gessen)	815	3				
98	1000' "	"	"	"	"	low		"	leather alder	"	545	5				
99	1200' "	"	"	"	"	"		"	"	" 4380'	130	2				
K3100	1400' "	"	"	"	"	"		"	grass	possible contamination *	2800	10				
01	1600' "	"	"	lt. gry.	chips	"		"	+ leather	elev 4400 (grass covered talus)	370	2				
02	1800' "	"	"	lt. brn.	"	nil		"	"	"	270	3				

* with what? G.R.P.

GEOCHEMICAL DATA SHEET — SOIL SAMPLING

SAMPLER Pearse / CooperNTS 1046/14WDATE 5 July, 1974PROJECT Winter Creek (08)

LINE _____

AIR PHOTO No. RC 4358-165

SAMPLE No.	LOCATION	DEPTH	HORIZ.	DESCRIPTION				SLOPE	VEG.	ADDITIONAL OBSERVATIONS OR REMARKS	ASSAYS					
				COLOUR	PART. SIZE	% ORG.	PH				Cu	Mo	Zn	Pb		
K3103	2000' E. of 3063	5fc	C	H. brn	clips of soil	low		steep	alder	elev. 4410'	215	2				
04	2200'	"	"	"	"	nil		"	grass		120	2				
05	2400'	"	"	"	"	"		"	"	4360'	675	2				
06	2600'	"	"	"	"	low		"	"		400	3				
07	2800'	"	"	"	"	"		"	shrub		254	1				
08	3000'	"	"	"	"	"		"	"	4310'	280	ND				
09	3200'	"	"	"	"	nil		"	"		375	1				
10	3400'	"	"	"	"	"		"	alder & Rucker	4280'	365	1				
11	3600'	"	"	"	"	low		"	"		375	1				
12	3800'	"	"	"	"	"		"	"	4300'	500	1				
13	4000'	"	"	"	"	"		"	"	flds. purpl. alder.	550	2				
14	4200'	"	"	"	"	"		"	"	4200'	300	1				
15	4400'	"	"	red brn	"	"		"	"	gossan	530	5				
16	4600'	"	"	"	"	"		"	"	4230'	892	7				
17	4800'	"	"	lt. brn.	"	"		"	"		425	2				
18	5000'	"	"	m. brn.	"	"		"	grass	4260'	360	1				
19	5200'	"	"	"	"	"		"	shrub		640	5				
20	5400'	"	"	red brn.	"	"		"	"	4210' gossan.	960	3				
K3121	5600'	"	"	m. brn.	"	"		"	"		660	4				
	End of lower line.															

GEOCHEMICAL DATA SHEET — SOIL SAMPLING

SAMPLER Pearse / CooperNTS 1046/14WPROJECT Winter Creek (08)

LINE _____

DATE 5 July, 1974AIR PHOTO No. EC 5358-165

SAMPLE No.	LOCATION	DEPTH	HORIZ.	DESCRIPTION				SLOPE	VEG.	ADDITIONAL OBSERVATIONS OR REMARKS	ASSAYS			
				COLOUR	PART. SIZE	% ORG.	Ph				Cu	Mo	Zn	Pb
K3122	cairn at el. 5500'	stc	C	red brn.	clips soil	low		mod.	grass	gossan; pyritic volc.	190	ND		
23	200' W 3m	"	"	m. "	"	"		"	"		150	ND		
24	400' "	"	"	" "	"	"		"	"	cl. 5400'	190	ND		
25	600' "	"	"	red. "	"	"		steep	"		170	ND		
26	800' "	"	"	m. "	"	"		"	"	" 5450'	285	ND		
27	1000' "	"	"	red "	"	nil		"	nil	gossan.	160	ND		
29	1200' "	"	"	" gry.	clips	low		"	grass	" 5420'	150	1		
30	1400' "	"	"	" brn	"	"		"	"		190	ND		
31	1600' "	"	"	" "	"	"		"	"	" 5450'	175	ND		
32	1800' "	"	"	lt. "	soil	"		"	"		219	1		
33	2000' "	"	"	m. "	"	"		"	"	" 5500'	270	1		
34	2200' "	"	"	lt. "	clips	"		"	"		115	1		
35	2400' "	"	"	" "	"	nil		"	"	" 5520'	200	3		
36	2600' "	"	"	" "	"	low		"	"		180	2		
37	2800' "	"	"	m. "	soil	"		mod	"	" 5540'	270	3		
38	3000' "	"	"	lt. "	clips	"		steep	"		305	5		
39	3200' "	"	"	" "	"	"		"	"	" 5510'	350	2		
40	3400' "	"	"	red "	"	nil		"	"		365	4		
41	3450' "	"	"	lt. "	"	"		"	"	" 5400' End of Line	240	5		

To: Texas Gulf, Inc.

REPORT No 751

PAGE No. 1

BONDAR-CLEGG & COMPANY LTD.

DATE: Oct. 4, 1974


701 - 1281 West Georgia Street
Vancouver, B. C. V6E 3J7
Attn: Mr. G. Peatfield

CERTIFICATE OF ASSAY

Samples submitted: Sept. 24, 1974
Results completed: Oct. 4, 1974
PROJECT: 08

I hereby certify that the following are the results of assays made by us upon the herein described pulp samples.

MARKED	GOLD		SILVER	Cu							TOTAL VALUE PER TON (2000 LBS.)
	Ounces per Ton	Value per Ton	Ounces per Ton	Percent	Percent	Percent	Percent	Percent	Percent		
WCIP - 5	0.20		0.41	4.25							
WCIP - 6	0.06		0.06	0.83							


 Registered Assayer, Province of British Columbia

APPENDIX B

STATEMENT OF EXPENDITURES

APPENDIX B

STATEMENT OF EXPENDITURES
KIT CLAIMS; WINTER CREEK

Salaries & Fringe Benefits

T.D.Pearse, Geologist	7 - 12 July, 1974		
	15 - 29 Aug., 1974		
	21 days @ \$50	-	\$1050
M.F.J.Cooper, Field Assistant	7 - 12 July, 1974		
	15 - 29 Aug., 1974		
	21 days @ \$25	-	\$ 525
			\$ 1575

Room & Board

	42 man days @ \$10		\$ 420
--	--------------------	--	--------

Equipment Rental

	Traeger radio		\$ 120
--	---------------	--	--------

Preparation of Topographic Base Map

	Photogrammetry by McElhanney Engineering Ltd.		\$ 520
--	--	--	--------

Travel & Helicopter Support

	6 hours Bell 206B helicopter @ \$210		\$ 1260
	Travel for crew		\$ 250

Report Preparation & Supervision

J.M.Newell, P.Eng.	1 day @ \$120	-	\$ 120
T.D.Pearse	Sept. 12 - 20 7 days @ \$50	-	\$ 350
Drafting, reproductions etc		\$ 125	\$ 595

Total Expenditure			<u>\$ 4740</u>
-------------------	--	--	----------------

Total expenditure claimed in Application for Certificates of Work:			\$ 4400
---	--	--	---------

Continued. . . .

N.B. The results of a geochemical survey are included in the supporting report. This work was completed prior to staking and the costs involved are NOT included in the above statement of expenditures.

Presented before me at the *City*
Vancouver, in the
Province of British Columbia, this *8*
day of *July* 1975, A.D.



John Turner
A Commissioner for Mining Affairs in the Province of British Columbia,
A Notary Public in and for the Province of British Columbia.

Sub-mining Recorder.

APPENDIX C.

STATEMENT OF QUALIFICATIONS

T.D. Pearse - obtained his B.Sc. degree from the University of British Columbia in 1971. He worked for 3 years as a staff geologist for Noranda Exploration Co., Ltd. in northern B.C., and assumed a temporary position as field geologist for Texasgulf, Inc. for the 1974 field season.

M.F.J. Cooper - is currently completing his B.Sc. degree at the University of Western Ontario and, as a field assistant, is thoroughly competent in all the tasks assigned to him during this project

A handwritten signature in cursive script, appearing to read "Maxwell", is located in the lower right quadrant of the page.

Department of
Mines and Petroleum Resources
ASSESSMENT REPORT
NO. 5509 MAP 2

To accompany Geological and Geo-chemical Report:
KIT Claims
Liard Mining Division
by T.D. Pearce October 1974

Texasgulf Inc.
FIG. C. Geological Map 5509
KIT GROUP MAP 2
104 G/14 W
WORK BY: TP, GRP
DRAWN BY: K.M.M.
DATE: JANUARY, 1975
DRAWING NO.

LEGEND			
	Syenite		Rock Geochem Station.
	Monzonite		x ^m Malachite
	Rhyolitic / Dacitic Tuff		py Pyrite
	augite / feldspar Porphyry Hybrid.		Limit of Gosson
	Quartz - Carbonate rock		
	Outcrop Pattern		
	Contact		X 5812 Elevation
	Fault		
	Shear		
	Bedding		
	Jointing		



Department of
Mines and Petroleum Resources
ASSESSMENT REPORT
MAP 3
NO. 5509



To accompany Geological and Geochemical Report:
KIT Claims
Liard Mining Division
BY T.D. Pearse October 1974

SCALE ONE INCH=400 FT

[Handwritten signature]



Texasgulf inc.		
FIG. D.	Geochemical Map	5509
	KIT GROUP	MAP 3
	IO4 G/14 W	
TP, GRP	K.M.M	JANUARY, 1975

o Total fines sample, ppm Cu/Mo




Department of
Mines and Petroleum Resources
ASSESSMENT REPORT
NO. 5509 MAP 4

To accompany Geological and Geo-chemical Report:

KIT Claims
Liard Mining Division
by T.D. Pearse October 1974

Scale and elevation datum based on a limited ground control resulting in good relative, but uncertain absolute map accuracy. Compiled from aerial photography at an approximate scale of 1 inch equals 1/600 feet flown in 1960.

TSXASGULLF INC. 5509 MAP 4
WINTER CREEK - STIKINE RIVER
PRELIMINARY RECONNAISSANCE TYPE MAPPING

 McElhanney Surveying & Engineering Ltd. 1200 West Pender Street, Vancouver B.C. Canada	Scale	1"=400'
	Contour Interval	50'
	Date	Aug 1974
	Job No.	06116-0
	Sheet No.	1 of 1