	LOG NO: 1127 RD.
Date received report back from amendments.	CTION:
FILE NO:	FILE NO: 87-812 16419

9/88

ASSESSMENT REPORT

THE HAT GROUP OF MINERAL CLAIMS

SKEENA MINING DIVISION

STEWART, B. C.

FILMED

NTS MAP SHEET 104B/8

LATITUDE: 56 DEGREES 20 MINUTES NORTH

LONGITUDE: 130 DEGREES 20 MINUTES WEST

GEOLOGICAL BRANCH ASSESSMENT REPORT

16,479

September 17, 1987

N. L. Tribe, P. Eng.

SUB-PECONDER
RECEIVED
NOV 24 1987
M.R. # ______ \$_____
VANCOUVER, B.C.

ASSESSMENT REPORT THE HAT GROUP OF MINERAL CLAIMS SKEENA MINING DIVISION

STEWART, B. C.

NTS MAP SHEET 104B/8

LATITUDE: 56 DEGREES 20 MINUTES NORTH

LONGITUDE: 130 DEGREES 20 MINUTES WEST

CONTENTS

	rage
INTRODUCTION	1
LOCATION	2
GEOLOGY	3
ECONOMIC GEOLOGY	5
1987 FROGRAM	6

LIST OF APPENDICES

APPENDIX I

ROCK DESCRIPTION AND ASSAYS

APPENDIX II

AUTHORS QUALIFICATIONS

APPENDIX III

STATEMENT OF COSTS

LIST OF ILLUSTRATIONS

Figure	#1	Prop	perty Location Map	1:5,000,000
Figure	#16	4 Cla	aim Map	1:100,000
Figure	#2	"The	at" Claim Map, Index Map	1:100,000
Figure	#26	4 " Th	at" Claim Map with Geology	1:50,000
Figure	#3	#13	Showing Geology and Assays	1:200
Figure	#4	#26	Showing Geology and Assays	1:500
Figure	#5	a.	Airphoto Overlay B.C. 5504-	-260
		b.	Airphoto Overlay B.C. 5504-	-261
		c.	Airphoto Overlay B.C. 5504-	-259
		d.	Airphoto Overlay B.C. 5504-	-258

ASSESSMENT REPORT THE HAT GROUP OF MINERAL CLAIMS SKEENA MINING DIVISION

STEWART, B. C.

NTS MAP SHEET 104B/8

LATITUDE: 56 DEGREES 20 MINUTES NORTH

LONGITUDE: 130 DEGREES 20 MINUTES WEST

INTRODUCTION

This report will cover work done on the Hat Mineral Claim Group during 1987. This work includes prospecting and geological field mapping.

LOCATION

The Hat Group of mineral claims is located just east of Mount Ketchum at the head of the Frankmackie Glacier. The claims abutt the Newhawk and Brinco claims at the headwaters of Sulphurets Creek and extend southward about 8 miles to that point of land where the two north arms of the Frankmackie Glacier come together. Sulphurets Creek and Mount Ketchum are located approximately 60 km northwest of Stewart, B. C. or approximately 900 km. north-northwest of Vancouver. Access to Stewart is by Highway 37, turning off Highway 16 at Kitwanga, about half way between Smithers and Terrace. Highway 37 is now paved through to Stewart from Kitwanga.

Access to the property is by helicopter either from Stewart (about one hour flight) or by helicopter from the Tide Lake airstrip (about 10 minutes flight). Good all-weather road connects Stewart and the Tide Lake airstrip which is used extensively as a staging area for airlifts into the Sulphurets. Vancouver Island Helicopters has several Bell 206B helicopters and a Bell 204 helicopter stationed in Stewart. The property requires helicopter support as road building into the property is not feasible in the initial stages. Moving about on the property is by foot with helicopter support being required for any sort of aggressive field program. The terrain is severe, with ice fields and glacier cover and very little vegetation other than a little

heather, moss, lichens and an occasional fireweed over most of the claim block. A small portion of the northwest corner of the claim block extends down into the Ted Morris Glacier where there is sparse alpine scrub hemlock, etc.

GEOLOGY

The geology of the Stewart area is typified by moderately folded intermediate volcanics and sediments intruded by a succession of plutons. Those areas around many of the mines are washed by a distinctive red iron alteration and broad zones of sericite alteration in which the numerous showings occur.

The area of the claims contains two volcanic formations, the Unuk River Formation and the Betty Creek Formation, and one sedimentary formation, the Salmon River Formation. Red iron alteration and intense sericitization prevails over most of the claims. The lowermost of these formations, the Unuk River Formation, is composed of dark green volcanoclastic rocks generally termed medium grained matric supported lapilli tuffs. The Unuk River Formation may be as much as 10,000 feet thick and forms the host rock for most of the deposits in the Stewart camp.

Above the Unuk River Formation lies the Betty Creek Formation of andesite pillow lavas. The Betty Creek Formation varies in thickness and may pinch out altogether. The Betty Creek Formation is often mineralized with massive pods of pyrite or pyrrhotite but is usually barren of economic mineralization.

Above the Betty Creek Formation is the Salmon River Formation of siltstones, greywackes and other fine to

medium-grained epiclastic or pyroclastic rocks. Relatively little work has been done in the Salmon River Formation rocks as the sediments were noted early on for their inability to maintain vein type structures that host most of the mineralization in the Stewart Camp.

The That Claims are ideally situated within the sericite and iron alteration zone and near the contact between the Salmon River Formation and the Unuk River Formation and centered on the Sulphurets Syncline. The Betty Creek Formation outcrops between the Salmon River Formation and the Unuk River Formation on the east side of the syncline but does not appear on the western side.

The Sulphurets Syncline forms a major feature in the area trending north-northwest and lunging to the north. The keel of the syncline should outcrop under the idefields on the southern cortion of the That Claims. Outcrops are sparse to this area but new rock is exposed every year as the idefields redeed.

The axis of the syncline forms a ridge of high ground with occasional outcrops and the remainder is covered with ice. However, the ice is expected to be relatively thin compared with the glaciers in the lower parts of the topography.

Tectonically, the Bowser Lake regional crustal cross feature passes along the southern portion of the claims. The cuartz-monzonite divelbliss stock outcrops just to the north of this cross-feature and makes a center of upwelling of magmatic material marking a probable center of volcanic activity.

It is felt that much of the icecap is not sufficiently

thick to mask airborne E.M. responses over good conductive formations.

ECONOMIC GEOLOGY

The most significant vein-type mineral deposit in the Stewart-Sulphurets is the gold-silver deposits at the Premier Mine. The Premier produced 1,820,048 oz. gold and 41,031,673 oz. silver from 4,713,167 tons over a 50-year period from 1918 - 1968. The host rocks at the Premier were sericitized iron-stained andesite lapilli tuffs near the top of the Unuk River Formation and a similarly altered plagioclase prophyry intrusive called the Premier Porphyry.

In a very similar setting the Newcana J.V. Sulphurets property has drilled off mineral reserves of 1,584,145 tons of 0.336 oz. Au/ton and 22.86 oz. Ag/ton also in as intensely sericitized andesite lapilli tuff near the spot of the Unuk River Formation and in an intensely sericitized plagioclase porphyry syenite intrusive.

Further to the north Skyline's Stonehouse Gold Deposit contains 746,000 tons of 0.62 oz. Au/ton and 0.94 oz. Ag/ton in intensely sericitized andesite lapilli tuff.

There are numerous other vein prospects in the immediate area of the claims in the early stages of development. Some of these include: Teuton's Tennyson Gold Zone, Teuton's Delta Gold Zone, Brinco's Kerr deposit, Newhawk's Sulphurats Breccia Zone and Iron Cap Zone, Magna Venture's Doc deposit, Soucie's East Gold Mine, Kerrisdale Resources' Red Bluff, Zone 21, 22, Teuton's

Treaty Central and Copper King Zones, Royal Scot's Scottie Mine and Tide Lake J.V.'s Bend showings.

1987 PROGRAM

The 1987 program consisted of four days of geological field mapping and approximately 10 days of prospecting. The geological mapping was reconnaissance in nature and very limited in scope. The prospecting was also very limited and considerable time was lost due to poor weather.

The typical Upper Unuk River package prevails through the southern portion of the claims with quartz sericite schists, cherts, andesite agglomerates and andesite tuffs intruded by syenite stocks.

Good sericite alteration with pyrite exists in broad halos around the syenites and in regional zones through the claims. An abundance of veining consisting of quartz and quartz-carbonate can be found throughout the ridge which forms the spine between the two arms of the Frankmackie Glacier. Occasionally these veins form into stockworks and occasionally the stockworks carry other sulphides, such as pyrite, pyrrhotite, arsenopyrite, sphalerite, galena. One such area is the Hat #13 discovery which is located near the northern end of the THAT 1 claim and on the top of the western slope of the spine. The outcrop shows a stockwork zone about 13 meters wide by 30 meters long in which the best vein assays: 0.915 oz. Au/ton/1.6 m.,

0.364 oz. Au/ton/0.4 m..

0.006 oz. Au/ton/1.2 m.

A second vein, approximately 30 meters from the first, from which a grab sample was taken assayed 0.825 oz. Au/ton. This stockwork is on the southern edge of the upper ice sheet and is covered with overburden to the south and ice to the north.

A second area of interest running along the western edge of the eastern arm of the Frankmackie Glacier was sketch mapped and sampled with 6 samples (Figure 4).

Summary of the assay results are as follows:

Sample No.

3805	.003	grab
3806	.003	4.0 m.
3807	.005	5.2 m.
38 08	.004	7.0 m.
3809	.003	grab
3810	.004	grab

Work on the northern claims was limited to defining the sediment volcanic boundary in extremely rough terrain.

Several days were lost here due to bad weather and only two traverses were made from the ridge to the ice on western slope on claim THAT #6. The rock was mainly quartz, sericite, schists of undetermined origin. A tentative meta-andesite meta-sedimentary contact was marked but will be checked further in the future.

N. L. Tribe, F. Eng.

APPENDIX I

ROCK DESCRIPTIONS AND ASSAYS

SAMPLE DESCRIPTIONS

LOCATION	SAMPLE NO	DESCRIPTION
i	2326	Chert pebble conglomerate (sed) moderately sericitized weakly silicified
2	2327	Quartz vein with moderate sericitic wallrock alteration
3	2328	Chert intensely silicified
4	2329	Small quartz in a cluster
5	2330	Quartz vein 45cm - 60cm with sulphides sphalerite ?
6	2331	Quartz boulder
7	2331	Quartz vein with minor pyrite
8	2333	Quartz vein sugary texture with minor sulphides - pyrite.
9	2334	Quartz vein in sedimentary rocks
10	2335	Quartz veins - zone of tensional veining
11	2336	Complex vein system in phyllitic sediments
12	2337	Complex vein system in phyllitic sediments
13	2338	Quatrz vein 60cm wide with arsenopyrite in phyllitic sedimentary rocks
14	2339	Complex vein system in cherty sediments
15	2340	Complex vein system in cherty sediments minor sulphides moderatly silicified
16	2341	Quartz vein in cherty sediments stained brown
17	2342	Quartz eye rhyolite
18	2343	Small quartz vein in cherts
19	2346	Stockwork of quartz carbonate veining 15m x 30m
20	2347	Stockwork of quartz carbonate veining 10m x 30m
13	2348	Quartz arsenopyrite vein
13	2349	Footwall rock phyllitic tuffs
13	2350	Hangingwall rock of phyllitic tuffs
13	3801	Stockwork in phyllitic tuffs
13	3802 3803	Arsenopyrite chips in a frost heave
21 22	3803 3804	Pale pink quartz veining in a sericitic andesite tuff
22 26	3805	Quartz carbonate vein stockwork in cherty tuffs Cherty tuffs intensely silicified with limonite staining
26 26	3806	Andesite lapilli tuffs intensely silicified intensely pyritized
26 26	3807	Andesite tuff intensely silicified
26	380B	Stockwork quartz and carbonate in andesite tuffs
26	3809	Intense stockwork quartz carbonate and sulphide - py po
26	3810	Andesite tuff intensely silicified
23	3811	Large kaolin altered zone in andesite tuff
24	3812	•
25	3813	Quartz sericite schist
27	3814	Chert with Pyrite
28	3815	Chert with Pyrite
29	3816	Andesite agglomerate cemented with quartz carbonate and sulphides py po
29	3817	Andesite agglomerate cemented with quartz carbonate and sulphides py po
30	3818	Quartz vein with pyrite
31	3819	Quartz vein with pyrite
32	3820	Silicified shear vein with pyrite
33	3821	Quartz vein with pyrite
34	32526	Quartz vein with pyrite
35	32527	Quartz vein with pyrite
35	32528	Quartz vein with pyrite
36		Quartz vein
37		Quartz vein breccia
38		Quartz vein
39	32532	Quartz vein with pyrite

NEWHAWK GOLD MINES LTD.

Certificate of Assay

all and				Project No.		-
то	Jantni	Resources	Inc	Date	Sept	7/87

SAMPLE No.	WT. A.T.	Dore	Au oz/ton	Ag oz/ton	Remarks
3801			0.009	tr	
3802			0.825	0-13	
3803			0.006	tr.	
3804			0-005	4	
3805			0.003	4	
3806			0.003	br	
3807			0.005	6	
3808			0.004	fr	
3809			0.003	4	
3810			0,004	4	
3811			0.004	4	
3812			0.005	x	
			0.006	t	
3813 3814			0.004	4	
3815			0.006	tr	
3816			F00.0	tr.	
3817			0.005	4	
3818			0.005	. 6	
3819			0.010	0.09	
3820			0.004	₹	
3821			0,003	4	
2345			0.007	4	
2346			0.006	4	
2347			0,007	fr	
23 48			0.364	0.09	
2349			0.006	tr	
2350			0.195	1.39,	
(Statement)					
		1			

CERTIFIED BY:



MAIN OFFICE 1521 PEMBERTON AVE. NORTH VANCOUVER, B.C. V7P 2S3 (604) 986-5211 TELEX: 04-352578

1630 PANDORA ST. VANCOUVER, B.C. V5L 1L6 (604) 251-5656

SSAY ANALYTICAL REPORT

CLIENT: Jantri Resources Inc.

DATE: August 31 1987

ADDRESS: Suite 6-117 East 15th Street : North Vancouver, B.C.

REPORTO: 871218 AA

: V7L 2P7

JOB#: 871218

PROJECT#: Sulphurets

INVOICE#: 871218 NA

SAMPLES ARRIVED: August 29 1987

TOTAL SAMPLES: 7

REPORT COMPLETED: August 31 1987

ANALYSED FOR: Ag Au

REJECTS/PULPS: 90 DAYS/1 YR

SAMPLE TYPE: 7 Rock

SAMPLES FROM: Jantri Resources Inc. COPY SENT TO: Jantri Resources Inc.

PREPARED FOR: Jantri Resources Inc.

ANALYSED BY: David Chiu

SIGNED:

Registered Provincial Assayer



MAIN OFFICE
1521 PEMBERTON AVE.
NORTH VANCOUVER, B.C. V7P 2S3
(604) 986-5211 TELEX: 04-352578

BRANCH OFFICE 1630 PANDORA ST. VANCOUVER, B.C. V5L 1L6 (604) 251-5656

REPORT NUMBER: 871218 AA	JOB MUMBER: 871218	Jantri Resources Inc.	PAGE 1 OF 1
SAMPLE #	Ag oz/st	Au oz/st	
32526	.77	<.005	
32527	1.14	<.005	
32528	1.51	.006	·
32529	.04	<.005	
32530	.05	<.005	
32531	.32	.005	
32532	.10	.005	

DETECTION LIMIT
1 Troy oz/short ton = 34.28 pps

.01 1 pps = 0.00012

pa parts per million

< = loss than

signed:

MAIN OFFICE: 1521 PEMBERTON AVE. N. VANCOLE ... & B.C. V7P 283 PH: (604)986-5211 TELEX: 04-352578 BRANCH OFFICE: 1630 PANDORA ST. VANCOLIVER B.C. VSL 1L6 PH: (604)251-5656

ICAP GEOCHEMICAL ANALYSIS

A .5 GRAM SAMPLE IS DIGESTED NITH 5 ML OF 3:1:2 HCL TO HNO3 TO H20 AT 95 DEG. C FOR 90 MINUTES AND IS DILUTED TO 10 ML NITH WATER. THIS LEACH IS PARTIAL FOR SM, MM,FE,CA,P,CR,MG,BA,PB,AL,MA,K,W,PT AND SR. AU AND PD DETECTION IS 3 PPM.

IS= INSUFFICIENT SAMPLE, MB= MOT DETECTED, -= NOT ANALYZED

COMPANY: JANTRI RESOURCES ATTENTION: PROJECT: SULPHURETS REPORT#: 871218PA JOB#: 871218 INVOICE#: 871218NA DATE RECEIVED: 87/08/29 DATE COMPLETED: 87/09/2 COPY SENT TO:

ANALYST W. Teaces

PAGE 1 DF 1

SAMPLE NAME	AG PP#	AL Z	AS PPN	AU PPM	BA PPH	BI PPM	CA Z	CB PPH	CO PPH	CR PPH	CU PPM	FE 1	X I	ng Z	in PPH	PO PPH	NA I	NI PPK	P	Poil	PB PPM	PT PPH	SB PP#	SN	SR PPN	U PPH	y PPN	ZN PPN
32525 32527 32528 32529 32530	29.4 40.7 53.5 .1	.14 .96 .11 1.46 1.48	243 403 84 42 276	ND 5 5 7 ND	8 2 18 5 13	ND ND 12 ND ND	.02 2.32 2.22 2.49 .27	2.0 33.2 >1000 9.3 1.6	2 42 13 29 18	16 53 40 88 11	626	5.20 21.88 2.42 24.04 7.64	.05 .13 .05 .14	.01 .58 1.15 1.60	40 565 700 290 394	11 10 10 13 3	.25 1.78 21.78 .75 .28	188 27 86 5	.01 .03 .01 .01	219 799 1143 22 12	110 110 110 110 110	EN E	56 60 221 11 6	ND 7 10 10 10	15 17 75 15 7	EM EX CK CK CM	ND ON ON ON	426 2554 63652 640 321
32531 32532	.: .:	1.97 1.23	17 54	ND NB	46 5	4 NB	.07 .04	.2 .1	5 5	27 :2	:57 28	3.54 !6.57	.01 .10	.52 .75	270 337	1	.11	5 3	.02	2 25	ND ND	MD WP	ИD 10	ND NB	3	10 10	GN QY	89 87
DETECTION LIMIT	.1	.01	3	3	1	3	.01	.1	1	1	1	.01	.01	.01	1	1	.01	1	.01	2	3	5	2	2	1	5	3	1



·Ł

MAIN OFFICE
1521 PEMBERTON AVE.
NORTH VANCOUVER, B.C. V7P 2S3
(604) 986-5211 TELEX: 04-352578

BRANCH OFFICE 1630 PANDORA ST. VANCOUVER, B.C. V5L 1L6 (604) 251-5656

ASSAY ANALYTICAL REPORT

CLIENT: JANTRI RES. LTD.

ADDRESS: Suite 6-117 East 15th St.

: North Vancouver, B.C.

: V7L 2P7

DATE: Sept 18 1987

REPORT#: 871331 AA ~

JOB#: 971331

PROJECT#: STEWART

SAMPLES ARRIVED: Sept 14 1987

REPORT COMPLETED: Sept 18 1987

ANALYSED FOR: Ag Au ICP

INVOICE#: 871331 NA

TOTAL SAMPLES: 5

REJECTS/PULPS: 90 DAYS/1 YR

SAMPLE TYPE: 5 Pulps

SAMPLES FROM: Mr. Harry Jang COPY SENT TO: JANTRI RES. LTD.

PREPARED FOR: JANTRI RES. LTD.

ANALYSED BY: David Chiu

SIGNED:

Registered Provincial Assayer

GENERAL REMARK: None



MAIN OFFICE 1521 PEMBERTON AVE. NORTH VANCOUVER, B.C. V7P 2S3 (604) 988-5211 TELEX: 04-352578 BRANCH OFFICE 1630 PANDORA ST. VANCOUVER, B.C. V5L 1L6 (604) 251-5656

REPORT HUNBER: 871331 AA	JOB MUNBER: 871231	JANTRE RES. LTD.	PAGE	1	OF	1
SAMPLE #	Ag oz/st	Au oz/st				
23 48M	5.69	. 425				
2350M	.19	.145				
3801M	. 58	.021	•			
3802M	.54	. 502				
381 9 M	<.01	<.005				

DETECTION LIMIT
1 Trey oz/short ten = 34.28 ppn

1 ppa = 0.00012

,005 mm = parts per million

a loce that

signed:

MAIN OFFICE: 1521 PEMBERTON AVE. N. VAN R B.C. V7P 2S3 PH: (604)986-5211 TELEX: 04-352578 BRANCH OFFICE: 1630 PANDORA ST. VANCOUVER B.C. V5L 1L6 PH: (604)251-5656

ICAP GEOCHEMICAL ANALYSIS

A .5 GRAM SAMPLE IS DIGESTED NITH 5 ML OF 3:1:2 HCL TO HNO3 TO H2O AT 95 DEG. C FOR 90 MINUTES AND IS DILUTED TO 10 ML WITH WATER. THIS LEACH IS PARTIAL FOR SM,MN,FE,CA,P,CR,MG,BA,PD,AL,MA,K,N,PT AND SR. AU AND PD DETECTION IS 3 PPN.

1S= INSUFFICIENT SAMPLE, ND= NOT DETECTED, -= NOT ANALYZED

COMPANY: JANTRI RES.LTD. ATTENTION:

PROJECT: STEWARD

REPORT#: 871331PA JOB#: 871331 INVOICE#: 871331NA DATE RECEIVED: 87/09/14
DATE COMPLETED: 87/09/28

COPY SENT TO:

ANALYST W Junes

PAGE 1 OF 1

SAMPLE NAME	AG PPM	AL Z	AS PPH	AU PPH	BA PPM	BI PPM	CA Z	CD PP#	CO PPM	CR PPM	CU PPM	FE I	K Z	MG Z	HN PPH	NO PPM	NA Z	NI PPM	P .	PB PPM	PD PPN	PT PPM	SB PPM	SN PPN	SR PPM	U PPM	u PPM	ZN PPM	
2348M 2350M 3801M 3802M 3819M	4.1 7.0 .1 7.0 5.5	.09 .31 1.90 .23 .10	9991 397 >101	10 18 ND 28 ND	13 33 41 17 19	3 5 3 ND ND	.08 .08 .64 .04	.1	203 34 10 246 11	10 5 6 7 5	164 49 16 343 10	6.53 5.79 4.69 19.94 6.73	.06 .08 .07 .20	.12 .10 .79 .08	177 226 780 231 49	3 1 3 5 36	.13 .09 .11 .32	27 10 13 21 10	.01 .06 .12 .03	88 11 5 ND 7	ND ND ND ND	ND ND ND ND	26 8 5 139 45	MD MD MD MD	3 7 12 1 ND	ND ND ND ND	NO ND ND ND	96 16 108 8 2	
DETECTION LIMIT	.1	.01	3	3	1	3	.01	.1	1	1	1	.01	.01	.01	.1	1	.01	1	.01	2	3	5	2	2	1	5	3	1	

,

)

÷

APPENDIX II

AUTHORS QUALIFICATIONS

<u>Authors Qualifications</u>

The author is a graduate of the University of British Columbia (1964) with a B.A.Sc. degree in Geological Engineering. The author has 24 years experience in mineral exploration and mine geology. The author has nine years experience in the Sulphurets and Stikine areas. The author has been registered as a Professional Engineer for the past ten years.

APPENDIX III

STATEMENT OF COSTS

JANTRI RESOURCES INC.


COST STATEMENT OF THE 1987 EXPLORATION PROGRAM "THAT" CLAIMS

DRAFTING - John Winfield	\$75 . 00
ASSAYS - Vangeochem	\$331.00
- Northair Group	\$331.00
HELICOPTER - V.I.H.	\$3,193.65
GEOLOGY - N. Tribe	\$6,030.46
- D. Rotherham	\$2,107.68
PROSPECTOR - M. McCrory	\$7,500.00
TOTAL	\$19,568.79

Of these amounts \$3,500 prospecting costs is applied to claims That \$1 to That \$4.

Of these amounts \$3,800 of geological costs is applied to claims That #5 to That #8.

FIGURES



100

200

100

MILES 0

Km O

200

300 MILES

400 Km


FIGURE I.


PROPERTY LOCATION MAP

HAT GROUP


SKEENA MINING DIVISION STEWART AREA, B.C.

TO ACCOMPANY A REPORT BY N. TRIBE & ASSOCIATES LTD. JULY, 1985








N. TRIBE & ASSOCIATES LTD. - 2611 SPRINGFIELD ROAD, KELOWNA, B.C. CANADA V1X 1B9 - TELEPHONE (604) 860-7661





N. TRIBE & ASSOCIATES LTD. - 2611 SPRINGFIELD ROAD, KELOWNA, B.C. CANADA V1X 1B9 - TELEPHONE (604) 860-7661














JANTRI RESOURCES INC.

> THAT CLAIMS GEOLOGY

AIRPHOTO OYELLAY £. E14.50 BC 5504 261