

LOG NO:	AUG 03 1993 RD.
ACTION:	
FILE NO:	

**GEOLOGICAL AND GEOCHEMICAL REPORT
ON THE DEAFIES CREEK PROPERTY**

Located Claims:

HAZ 5 259221(7)
OK 259255(9)

Vernon Mining Division

N.T.S. 82 L/6, 82 L/7

50° 18.6' N., 118° 59' W.


Owner:

ZICTON GOLD LIMITED

504-455 Granville Street
Vancouver, British Columbia

V6C 1T1

FILMED



**John Ostler, M.Sc.P., Geo.
Consulting Geologist**

June 28, 1993

**GEOLOGICAL BRANCH
ASSESSMENT REPORT**

22,954

CONTENTS

SUMMARY	iii
1.0 INTRODUCTION	1
1.1 Terms of Reference	1
1.2 Location and Access	1
1.3 Terrain and Vegetation	2
1.4 Property	3
1.5 History of Development around the Lumby Gold Mine	4
1.6 Previous Work on the Deafies Creek Property	6
1.7 Summary of Present Work	8
1.8 Claims Worked On	9
2.0 GEOLOGY AND GEOPHYSICS	9
2.1 Regional Geology	9
2.2 Regional Geophysics	13
2.3 Economic Mineralization at the Lumby Gold Mine	14
2.4 Property Geology	15
2.4(i) Stratigraphy.	15
2.4(ii) Deformation and Metamorphism of Sicamous Formation Rocks	16
2.4(iii) Exposed Economic Mineralization	17
3.0 SOIL GEOCHEMISTRY	18
3.1 1993 Soil Survey	18
3.2 Interpretation of Soil-metal Distribution Curves	21
3.3 Interpretation of 1993 Soil Survey Results	22
4.0 CONCLUSIONS AND RECOMMENDATIONS	22
4.1 Conclusions	22
4.1(i) Geology	22
4.1(ii) Soil Survey	23
4.2 Recommendations	23
5.0 REFERENCES.	24
6.0 ITEMIZED COST STATEMENT FOR THE MAY-JUNE, 1993 EXPLORATION PROGRAM.	26
APPENDICES	
A. Methods of Soil Analysis and Rock Assay	After text
B. Rock Assays	"
C. Soil Analyses	"
D. Certificate of Qualification	"

CONTENTS
Continued

FIGURES

1.	General Location	After text
2.	Location and Terrain	"
3.	Airborne Electromagnetic Survey	"
4.	Regional Geology from G.S.C. Open File 637	"
4A.	Legend to Figure 4	"
5.	Table of Geological Events and Lithological Units in the Lumby Area.	"
6.	Aeromagnetism: E.M.R. Map 8502G	"
7.	Geology North of Deafies Creek	In pocket
8.	Copper Distribution Curve: Central Area	After text
9.	Arsenic Distribution Curve: Central Area	"
10.	Lead Distribution Curve: Central Area	"
11.	Zinc Distribution Curve: Central Area	"
12.	Copper, Arsenic, Lead and Zinc in Soils: Central Area	In pocket

GEOLOGICAL AND GEOCHEMICAL REPORT ON THE DEAFIES CREEK PROPERTY

SUMMARY

The writer was retained by Zicton Gold Limited of Vancouver, B.C. through Cassiar East Yukon Expediting Ltd. to conduct an exploration program of geological mapping and soil-geochemical survey on the Deafies Creek Property from May to June, 1993.

The Deafies Creek Property is located on the eastern flank of the highland that separates White Valley to the east and Okanagan Valley to the west. This rolling upland area forms part of the southern part of the Shuswap Highland of south-central British Columbia. It comprises two located claims which are owned 100% by Zicton Gold Limited. These claims cover 30 claim-units; about 700 ha (1680 A) after deducting areas of overlapping claims. They adjoin the Lumby Gold Mine Property to the northwest.

The property is centred on 50° 18.6' north and 118° 59' west in the Vernon Mining Division of B.C.

The Village of Lumby is the closest settlement to the Deafies Creek Property. It is 22 km (13.4 mi) east of Vernon and 5.5 km (3.4 mi) south of the southern boundary of the Deafies Creek Property. Lumby is about 504 km (307 mi) from the port of Vancouver. Access to the central part of the Deafies Creek Property from Lumby is via the Shuswap Falls, Trinity Creek and Deafies Creek roads. Access to the central area containing the 1993 soil grid and the main trench is via a branch road that diverges uphill from the Deafies Creek road about 2.8 km in from the Trinity Creek road. Four-wheel drive is required only during periods of poor weather.

During the 1993 program, the crew stayed in a motel in Vernon and drove to the property each day.

Elevations on the property range from about 725 m (2380 ft) where Deafies Creek leaves the easterly boundary of the claims to 1220 m (4000 ft) along the western boundary of the property (Figure 2).

Soils in the Deafies Creek valley developed beneath a fir, spruce pine and hemlock forest. About half of the property has been logged within the last 20 years. Generally, soil profiles are sufficiently mature to have distinct undisturbed horizons amenable to meaningful soil survey results.

Zicton Gold Limited is the recorded owner of 100% of the Deafies Creek Property.

The Deafies Creek Property adjoins the Lumby Gold Mine Property to the northwest. Exploration on the claims has been focused on finding similar gold-bearing structures to those found at the mine. Most recent encouragement has been from the discovery of an extensive airborne electromagnetic anomaly that extends from the Quinto Mining Corp.'s property onto the eastern part of the Deafies Creek Property.

The Deafies Creek Property is underlain by a sequence of Triassic-age turbidites, pelites and high-level diorite intrusives that form part of the Slocan Group. These rocks were complexly deformed, metamorphosed and intruded by granitic rocks in the Lumby region. Deformation and metamorphism began with the folding of the Kootenay Arc at the western margin of proto-North America during the Middle Jurassic Period and culminated during the extension of the Shuswap metamorphic terrane during the Late Cretaceous Period.

During the Pleistocene Stage, glacial erosion moulded the current topography of the property-area. Minor Holocene uplift and erosion has exposed some of the older stratigraphy on the property.

Only one sulphide mineral showing is known to exist on the Deafies Creek Property. It is located in the main trench at the centre of the 1993 soil survey. Two greywacke outcrops exposed in the trench contain white quartz veins. A small pile of massive pyrrhotite cobbles is located

near the vein outcrops. Reportedly, these cobbles are cleaved from the veins.

No economic mineralization was found at this showing.

Soils were sampled in the north-central part of the Deafies Creek Property. The 1993 soil grid was centred on the main trench which has been the focal point of past surface exploration and drilling on the property.

Samples were analyzed for copper, arsenic, silver, gold, lead, zinc, bismuth, mercury, antimony and molybdenum.

Two northwest-southeast trending fault-like structures are indicated in the grid-area by the distributions of copper, arsenic, lead and zinc in soils. These structures seem not to be highly mineralized in the grid-area. It is very unlikely that these structures are the cause of the airborne electromagnetic anomaly discovered by Quinto Mining Corp. to be trending onto the eastern part of the property.

GEOLOGICAL AND GEOCHEMICAL REPORT ON THE DEAFIES CREEK PROPERTY

1.0 INTRODUCTION

1.1 Terms of Reference

The writer was retained by Zicton Gold Limited of Vancouver, British Columbia through Cassiar East Yukon Expediting Ltd. to explore and report on the Deafies Creek Property. This exploration was part the Lumby project during which, work was conducted on the Saddle Mountain and Deafies Creek properties.

Consequently, itemized costs were reported herein for the whole project and apportioned to the two properties.

This report is of work conducted on the Deafies Creek Property. Work on the Saddle Mountain Property is contained in another report (Ostler, 1993).

Exploration was conducted on the Saddle Mountain and Deafies Creek properties from May 26 until June 10, 1993. Historical review, data compilation and processing was conducted from May 12 until June 28, 1993.

1.2 Location and Access

The Deafies Creek Property is located on the eastern flank of the highland that separates White Valley to the east and Okanagan Valley to the west. This rolling upland area forms part of the southern part of the Shuswap Highland of south-central British Columbia (Figure 1). The property comprises two located claims which are owned 100% by Zicton Gold Limited. These claims cover 30 claim-units; about 700 ha (1680 A) after deducting areas of overlapping claims. This property adjoins the Lumby Gold Mine Property to the northwest.

The claims are centred on 50° 18.6' north and 118° 59' west in the Vernon Mining Division of B.C. (Figure 2).

The City of Vernon is the closest major supply and service centre to the Lumby area and the Deafies Creek Property. Vernon is located near the northern end of Okanagan Lake in Okanagan Valley. It services strong

local agricultural and logging industries. Most services required for property exploration and development can be found there. The Village of Lumby is the closest settlement to the Deafies Creek Property. It is 22 km (13.4 mi) east of Vernon and 5.5 km (3.4 mi) south of the southern boundary of the Deafies Creek Property. Lumby is about 504 km (307 mi) from the port of Vancouver. Access to the central part of the Deafies Creek Property from Lumby is via the Shuswap Falls, Trinity Creek and Deafies Creek roads. Access to the central area containing the 1993 soil grid and the main trench is via a branch road that diverges up hill from the Deafies Creek road about 2.8 km in from the Trinity Creek road. Four-wheel drive is required only during periods of poor weather.

During the 1993 program, the crew stayed in a motel in Vernon and drove to the property each day.

1.3 Terrain and Vegetation

The Deafies Creek Property is located on the eastern flank of the highland that separates White Valley to the east and Okanagan Valley to the west. This rolling upland area forms part of the southern part of the Shuswap Highland of south-central British Columbia (Holland, 1976).

Holland's description of the terrain of Shuswap Highland containing the area around the Deafies Creek Property is as follows:

The Shuswap Highland ... extends southward from Mahood and Murtle Lakes to the Coldstream Valley east of Vernon, and lies between the Thompson Plateau on the west and the Monashee Mountains on the east. It is 140 miles long and 50 miles wide ...

The Shuswap Highland consists of gentle or moderate sloping plateau areas rising from about 5,000 to over 7,000 feet, dissected by the Clearwater, North Thompson, Adams and Shuswap Rivers and their tributaries into large polygonal upland tracts. The valley sides are commonly steep because of glacial erosion, and the total relief in the uplands is moderate.

... The high points are progressively lower to the south. Most ridges and summits are rounded, and despite the height of much of the terrain the country lacks the jagged sawtooth profiles of the mountains to the east.

The effects of glaciation in the region were largely to soften and reduce the upland relief while steepening and deepening the valleys. Cirque glaciation on northeastern exposures was a minor feature.

Numerous large lakes, such as Murtle, Adams, Shuswap and Mabel Lakes, occupy some of the major valleys of the area, and these as well as the rather extraordinary pattern of drainage, quite

obviously diverted from its preglacial flow are legacies of pleistocene ice occupation.

Holland, S.S.; 1976: pp. 73-74.

The Deafies Creek Property occupies most of the upper part of the Deafies Creek valley (Figure 2).

White Valley contains Bessette Creek, a northeasterly flowing tributary of Shuswap River which enters Mabel Lake about 17 km (10.4 mi) northeast of the claims. Adequate water for mining purposes exists in the property-area; however, use may have to be negotiated with agricultural users located there.

Elevations on the property range from about 725 m (2380 ft) where Deafies Creek leaves the easterly boundary of the claims to 1220 m (4000 ft) along the western boundary of the property (Figure 2).

Soils in the Deafies Creek valley developed beneath a fir, spruce pine and hemlock forest. About half of the property has been logged within the last 20 years. Original forest on the southern part of the claims will be logged soon judging by the fresh timber cruising and new haulage roads located there.

Soils are developed on a relatively thin layer of till and decomposed local bedrock. Presently, the 1993 grid-area is covered with young second growth forest. Off skidder trails, soil profiles are sufficiently mature to have distinct undisturbed horizons amenable to meaningful soil survey results.

Soon, there will not be sufficient available timber on the Deafies Creek Property to support a major mining operation. However, timber is readily available at the sawmills at Lumby.

The property-area usually has mild winters and hot dry summers. Winter snowfall is generally light.

1.4 Property

The Deafies Creek Property comprises the following claims located in the Vernon Mining Division of British Columbia (Figure 2):

Claim Name	Record No.	No. of Units	Record Date
OK	259255(9)	20	September 20, 1985
HAZ 5	259221(7)	10	July 11, 1984

These claims are owned 100% by Zicton Gold Limited of Vancouver, British Columbia.

The property boundaries have not been surveyed.

1.5 History of Development around the Lumby Gold Mine

The history of the development of the gold mine at Lumby is most succinctly summarized by Drummond and Howard (1993). The following is paraphrased from their report by the writer.

Mineralization at the southern end of Saddle Mountain adjacent to Lumby, British Columbia was noted in the early 1900's by a local teacher whose prospect workings became known as the Teacher Showing (Figure 3). Fifty years later, the Chaput Logging Company exposed silver-lead-zinc-copper veins during road building for logging on the western slopes of Saddle Mountain. Claims were staked around what later became known as the Mine Showing (Figure 3). F.K. Explorations acquired the property in 1968. The Mine Showing was developed underground and a 50-ton flotation mill was built. From 1968 until 1970 over 1500 tons of concentrate was shipped from the mine to the smelter at Trail, B.C. Alberta Gypsum Ltd. had control of the property from 1971 to 1973 and conducted extensive surface and underground exploration. The mine was operated sporadically by Coast Interior Ventures Ltd. from 1974 until 1979. The capacity of the mill was expanded to 150 tons per day during 1980. Operations terminated in 1981.

The Quinto Mining Corporation acquired the mine and greatly expanded the property during 1983 with the plan to put it back into production based on greatly expanded ore reserves.

That year Quinto outlined co-incident soil and very low frequency electromagnetic (VLF-EM) anomalies on top of Saddle Mountain. The anomalous area was named the Plateau Zone (Figure 3). Trenching of the zone uncovered a gold-bearing shear of significant amounts. Gold was

contained sporadically throughout sheared breccia and quartz veins contained within a graphitic host.

The company was greatly encouraged by the results of initial exploration at the summit of Saddle Mountain and embarked upon a greatly expanded program of exploration in 1985. That year 10 reverse circulation drill holes and 1396 ft (425 m) comprising 13 diamond drill holes were completed to extend the area of known mineralization. In 1986, exploration comprised geological mapping of the Saddle Mountain area (Kuran, 1986), 2700 m (8856 ft) of NQ diamond drilling and extensive VLF-EM and magnetometer surveys.

Exploration was continued through 1987 with 32 additional reverse circulation and 7 more diamond drill holes totalling 3030 m (9938 ft) of drilling. Geophysical survey coverage of Quinto's property was expanded. New surveys included airborne VLF-EM and magnetometer coverage of the whole property (Figure 3). Several broad anomalies were identified over the area.

During 1987 attention was focused on development, mining and milling considerations. A metallurgical test was conducted by Lakefield Research. In 1988 Kilborn Engineering constructed a computer model of the Plateau Zone comprising 21 vertical sections and a joint venture with Golden Seville Resources Ltd. was negotiated to test a column leach process on the graphitic gold mineralization. Late that year a preliminary production feasibility study was completed by Bechtel Canada.

Underground development at the 808 m level comprising 186 m (610 ft) of 3.05 x 3.66 m (10 x 12 ft) drift and two crosscuts at portal + 140 and 190 m comprising a total of 105 m (344 ft) was driven during 1990 by Sancold Resources Contractors Ltd. for Quinto. Bradley (1990) mapped and sampled the new Plateau Zone workings.

Recently, the potential for graphite and mica production as well as gold production of the mineralization on Quinto's property was studied by Drummond and Howard (1993).

1.6 Previous Work on the Deafies Creek Property

Zicton Gold Limited's Saddle Mountain and Deafies Creek properties were staked during the mid-1980's adjacent to The Quinto Mining Corporation's Lumby Property to the east and northwest respectively.

Work by Quinto to put its Lumby gold mine back into production with new ore reserves has been paralleled by development on Zicton's properties. The two companies have had a very good working relationship. Often, information has been shared to the benefit of all.

Preliminary prospecting and geophysical work done on the north-central part of the Deafies Creek Property during the early 1980's revealed a quartz outcrop measuring 2 x 3 m (6.6 x 9.8 ft) that was associated with a local geophysical anomaly. Subsequent trenching uncovered a zone of quartz veins and sulphide pods that have been interpreted to be a shear zone (Allen, 1989).

Quinto Mining Corp.'s airborne electromagnetic survey indicated that a broad anomaly extended from Quinto Mining Corp.'s claims westward onto the Deafies Creek Property (Figure 3). It was hoped that the mineralization uncovered by the work in the main trench area was associated with a large mineralized structure that extended beneath the electromagnetic anomaly across the Deafies Creek Property (John Hilton pers. comm.).

Further trenching during 1989 exposed a length of about 25 m (82 ft) of quartz vein containing pods of pyrrhotite with minor chalcopyrite, pyrite, galena and sphalerite (Allen, 1989).

Diamond drill hole OK89-1 was collared about 15 m (50 ft) south of the quartz knob exposed near the eastern end of the main trench (Figures 7 and 12). It was drilled at an orientation of 355°/-60° for a length of 61.0 m (200 ft).

The following year, diamond drill hole Z90-2 was collared about 49 m (150 ft) west of DDH OK89-1. It was drilled at an orientation of 005°/-65° for a length of 61.6 m (202 ft) to test the rock beneath the western end of the main trench (Figures 7 and 12) (Allen, 1990).

Drilling results were reported by Allen (1989, 1990) as follow:

	Direction	Length
DDH OK89-1	355°/-60°	61.0 m
DDH Z90-2	005°/-65°	61.6 m

Summary of Drill Log for OK89-1

Bedrock was reached at a depth of 5.8 meters. The first 4.9 meters of bedrock is grey tuff with disseminated pyrite and minute pyrite-filled fractures. The underlying 23.3 meters is fine-grained black argillite with calcite veinlets and fine fractures containing pyrite, along with sulphide disseminations and irregular blebs. The bottom 27 meters is harder black and grey banded argillite with disseminated and thinly veined pyrite.

Allen, A.R.; 1989: p. 6.

Summary of Drill Log for Z90-2

Beneath 4.9 metres of overburden is 11.9 m. of black fine-grained argillite and graphitic argillite containing narrow and irregular quartz veins and pyrite in fractures and disseminations. This is underlain by 2.1 m. of grey argillite with pyrite and pyrrhotite, 12.1 m. of dark grey argillite with pyrite in irregular zones and fractures, a 6.1 m. zone of black argillite and 18.3 m. of black graphitic argillite with pyrite blebs and disseminations as well as in fractures, and chalcopyrite with blue and green oxides.

Allen, A.R., 1990: p. 4.

A series of 27 samples were taken from road outcrops of Slocan Group metasediments during the 1991 exploration program (Halliwell and Allen, 1991). Whole rock geochemical analysis conducted on those samples showed them to be typical greywackes and slates.

During the 1992 exploration program, metasedimentary road outcrops were mapped and a very low frequency electromagnetic (VLF-EM) survey was conducted over the area around the main trench. A total of 22 soil samples were taken in the grid-area (Halliwell and Allen, 1992).

Two groups of anomalies were contained within the VLF-EM results. One group was located about 200 m (656 ft) north of the main trench. The writer interpreted these anomalies to be associated with the diorite greywacke contact (Figure 7). The other group of anomalies was located around the main trench.

The current exploration program reported upon herein is an extension of the 1992 work. A soil survey was conducted over the area covered by the 1992 VLF-EM anomalies and, outcrop and float on the northern part of the property was mapped (Figures 7 and 12).

1.7 Summary of Present Work

Field work on the Deafies Creek Property was conducted from May 26 until June 10, 1993. Data compilation continued until June 28, 1993. The work was undertaken by:

John Ostler; M.Sc., P.Geo. West Vancouver, B.C.	Consulting Geologist
David R. Jones, B.Sc. Vancouver, B.C.	Geological Technician
David P. Nunuk, B.Sc. Aldergrove, B.C.	Geological Technician
John Hilton Vancouver, B.C.	Professional Prospector

The 1993 work program on the Deafies Creek Property included the following:

Geological and Geochemical Work

A. Geological Mapping 350 ha mapped at a scale of 1:5000 (Figure 7).	1.50 man-days
B. Soil Geochemical Survey Central Grid-area: 5.05 km of compass line was flagged and sampled at 50 m intervals comprising the 1993 grid covering 20 ha 99 samples were taken. All samples were analyzed for Cu, As, Pb, Zn, Ag, Au, Hg, Bi, Sb, Mo (Figures 8 to 12 and Appendices A and C).	3.50 man-days
C. Rock Geochemical Sampling: 0.1 ha of trenches and on the OK claim (Figures 7 and 12, and Appendices A and B).	0.75 man-days
D. Transportation, expediting, camp set-up, data compilation, drafting and report time	<u>9.05</u> man-days
Total time spent on the Deafies Creek Property during the May-June, 1993 work program	14.80 man-days

1.8 Claims Worked On

During 1993, work was done on the following claims:

Claim Name	Record No.	No. of Units	Current Expiry Date
OK	259255(9)	20	September 20, 1985
HAZ 5	259221(7)	10	July 11, 1984

2.0 GEOLOGY AND GEOPHYSICS

2.1 Regional Geology

The rocks in the area around Lumby were mapped by Jones (1959) as undefined units of the Monashee Group which was assigned to the Shuswap Terrane. Their age; Archean or younger, and correlation with other lithostratigraphic units was unknown. It was thought that they were bounded by two northwesterly trending faults, one of which separated them from Triassic-age Slocan Group rocks to the east.

The Shuswap Metamorphic Complex is exposed just west of the Lumby area.

It was considered by Jones and his contemporaries to be a western extension of the Canadian shield that was in fault contact with adjacent younger rocks. Okulitch (1979) found that along some of the margins of the complex, Mesozoic-age units could be traced into the complex and correlated. Consequently, Okulitch (1979) determined that the Shuswap Metamorphic Complex had a Precambrian-age core that was added to during subsequent orogenic and metamorphic events.

Okulitch (1979) found that the Lumby area was underlain by shale, massive siltstone, conglomerate and tuff (Figure 4). He could not find any evidence of a boundary fault between the Sicamous Formation of the Monashee Group and adjacent Slocan Group rocks and thus assigned everything to the Late Triassic-age Slocan Group.

The Slocan Group rocks near Lumby have been influenced profoundly by their proximity to the Shuswap Metamorphic Complex which has destroyed many sedimentary structures and lithological relationships through conversion of sedimentary and volcanic rocks to phyllites, schists and

gneisses. East of the property-area the history of these units has been discerned from extensive exploration.

The Slocan Group rocks form some of the upper part of the Kootenay Arc, which extends in southwestern British Columbia from the U.S. border to northeast of Revelstoke (Douglas et al; 1970).

Kootenay Arc sediments and volcanics were deposited at the western margin of proto-North America in the Cordilleran Geosyncline. Kootenay Arc deposition from Late Proterozoic until Middle Palaeozoic time was in a large eugeosyncline that segregated into smaller sub-basins during the Late Palaeozoic Era. Mesozoic deposition was mostly miogeosynclinal.

The older eugeosynclinal assemblage is exposed mostly in the eastern part of the Kootenay Arc; the younger miogeosynclinal assemblage is exposed mostly in the western part of the Kootenay Arc. The Lumby area is underlain by the Slocan Group which forms part of that miogeosynclinal assemblage.

Read and Wheeler (1976) mapped the Slocan Group over a broad area just northeast of the Lumby area. Their description of the Slocan Group was as follows:

The Slocan Group lies between the Kuskanax and Nelson Batholiths and extends into Vernon map-area (Jones, 1959), between Thor-Odin and Pinnacles Domes. The group consists of a thick unit of pelitic rocks overlain by approximately 4,000 feet of volcanic rocks. At the base of the group, lenses of conglomerate and sedimentary breccia ... , composed of Kaslo detritus, disconformably overlie the Kaslo Group. Near the base, limestone ... , up to 100 feet thick, forms layers intercalated with grey argillite, phyllite and fine-grained quartzite ... Near the top of (this) map unit ... , rocks become tuffaceous and pass into meta-andesite and meta-dacite tuffs and flows ... , and augite meta-basalt and meta-andesite flows and tuffs ... Between Columbia River and Slocan Lake, these volcanic rocks core the depressions of the doubly-plunging synclines ... West of Slocan Lake, an increase of metamorphic grade towards Valhalla Dome, which lies south of the map-area, has converted metasedimentary rocks of the Slocan Group to mica schist ... and marble ...

Read, P.B. and Wheeler, J.O.; 1976:
Descriptive Notes to G.S.C., O.F. 432.

Slocan Group rocks are intruded by a suite of calc-alkalic batholiths and stocks that are part of the Nelson Batholith (Read and Wheeler, 1976). Nelson Batholith intrusions are concordant intrusions elongate parallel to the westerly trend of the country rock. They are

dated by Read and Wheeler (1976) as Jurassic to Cretaceous, generally about 164 million years old.

Read and Wheeler (1976) recognized four phases of regional deformation in Kootenay Arc rocks east of the Lumby area.

The first phase of deformation produced rootless isoclinal folds with well-developed, axial plane foliation during the Middle Palaeozoic. This phase of deformation was completed before the Slocan Group was deposited and consequently, has no relevance to the geology of the Saddle Mountain or Deafies Creek properties.

The second and third phases of deformation occurred during the Middle Jurassic Period after deposition of the Slocan Group and early during emplacement of Nelson Batholith intrusions. Read and Wheeler (1976) estimated that these phases of deformation and associated regional metamorphism occurred between 178 and 164 million years ago.

Second and third-phase folds are open to tight folds with a crenulation axial plane cleavage.

Read and Wheeler (1976) recognized a late phase of deformation that produced microscopic kink folds of various orientations in phyllites.

Jurassic-age regional metamorphism in Slocan Group rocks varies from chlorite to biotite sub-facies of the greenschist facies of metamorphism. Locally, Slocan Group rocks are metamorphosed to granulite facies due to contact metamorphism during the emplacement of intrusions related to the Nelson Batholith and anatexis during extension of the Shuswap Metamorphic Complex.

Okulitch (1979) described the structures observed in Slocan assemblage rocks near Lumby as follows:

Structures in the Sicamous Formation are well-developed at all scales but are variable in their style and mutual relationships throughout the project-area. Bedding and sub-parallel foliation are ubiquitous; the latter is particularly evident although fine laminar compositional layering is also present... Attenuated isoclinal folds are common and these early structures are similar in many respects to those in adjacent older units... Megascopic early folds in the Sicamous Formation on the flanks of the Chase and Silver Star Anticlines are the same as those described in the Silver Creek and Tsalkom Formations.

Late and latest structures present in the Sicamous Formation are for the most part also the same as in adjacent units. Possibly

significant exceptions are latest brittle folds west of Adams Lake that plunge gently east, which are of anomalous orientation, and polyphase folds in Coldstream and Creighton Valleys. These features may be related to major faulting...

Okulitch; A.V.: 1979;
Descriptive Notes to G.S.C.; O.F.637, Map B

Plutonic rocks ranging in age from Ordovician to Cretaceous have been mapped in the area around Lumby and Vernon (Jones, 1959; Okulitch, 1979). Small intrusive plugs of Jurassic-age diorite and Cretaceous-age granite were mapped near some of the peaks near the Deafies Creek and Saddle Mountain properties (Okulitch, 1979). The intrusions have been related to the major orogenic events that have affected the region.

These orogenic events were described by Okulitch (1979) as follows:

The Columbian Orogeny, occurring during the Early Jurassic to Mid-Cretaceous time, was the major event affecting rocks in the project-area. Most of the polyphase ... folding, regional metamorphism and faulting took place at this time. Extensive plutonism accompanied and followed deformation...

Within the project-area, radiometric data ... suggest that closure of the K-Ar isotopic system during waning regional metamorphism and deformation took place at least 130 to 155 MA ago (Early Cretaceous to Middle Jurassic). Early Jurassic rocks were affected by most deformational phases of the orogeny; Early Cretaceous plutons ... are post-tectonic.

Uplifting and erosion followed the Columbian Orogeny. Final cooling of high-grade metamorphic rocks may not have taken place until about 50 MA ago ... or a discrete thermal event, perhaps associated with Eocene plutonic and volcanic rocks affected the Rb-Sr and K-Ar isotopic systems ... Movement along northerly trending faults and latest warping preceded or accompanied extrusion of (Eocene or Oligocene-age volcanics). Numerous feeder dykes followed fracture and fault planes. Such tensional features may be induced by post-orogenic erosion, uplift and cooling of the crust ...

Post-Eocene uplift and faulting took place predominantly in the Shuswap Metamorphic Complex and resulted in erosion of (the Tertiary-age volcanics) and further exposure of the metamorphic terrain.

Okulitch; A.V.; 1979;
Descriptive Notes to G.S.C., O.F.637, Map B

Large cliff-forming outcrops of Tertiary-age flood basalts and andesites are exposed on the eastern part of the Saddle Mountain Property about 8 km east of Lumby (Ostler, 1993). There, they unconformably overlie Slokan Group metasediments.

The Lumby area underwent significant glaciation during the Pleistocene Stage, producing broad valleys. Late Pleistocene and Recent glacio-fluvial sediments filled White Valley to its present topographic level and a thick mantle of glacial till was deposited on lower hill slopes. A thick apron of glacio-fluvial sediments covers the central part of the Saddle Mountain Property burying Slovan Group rocks located there (Ostler, 1993).

Interglacial palaeosols uncovered in the trenches at the Deafies Creek Property indicate that the profiles of upper ridge slopes in the White Valley area have been changed little by later Pleistocene glaciation. From fresh Slovan Group rocks beneath the palaeosols it appears that Tertiary-age regoliths were scraped off the hillsides by earlier Pleistocene-age glaciations.

Recent rejuvenation and valley downcutting occurred along Blue Springs Creek in the southern part of the Saddle Mountain Property resulting in the erosion and exposure of glacio-fluvial stratigraphy there.

The preceding geological history is summarized in a table of geological events and units that accompanies this report (Figure 5).

2.2 Regional Geophysics

An aeromagnetic survey was flown over the Lumby area in 1972 and published in 1973 by Geotrex Limited. The part of the survey that includes the Saddle Mountain and Deafies Creek properties is available from the Department of Energy, Mines and Resources (G.S.C.) as Map 8502G (Figure 6).

In general, the magnetic profile in the property-areas is rather flat. A significant exception is over the Oligocene to Eocene-age flood basalts exposed in the eastern part of the Saddle Mountain Property. There, an intense magnetic high corresponds with the location of the flood basalts. Mapping in that area indicates that the intensity of the magnetic high is related directly to local basalt thickness (Ostler, 1993).

The Quinto Mining Corporation conducted an extensive airborne electromagnetic survey over White Valley north of the Village of Lumby (Figure 3). That survey included the Lumby Gold Mine and parts of the Saddle Mountain and Deafies Creek properties. Several anomalies that extended from Quinto's property onto Zicton's ground were identified. At present, most of the anomalies have not been explored sufficiently.

2.3 Economic Mineralization at the Lumby Gold Mine

Saddle Mountain was named well. It has two rounded peaks which are composed of diorite and its gneissic equivalents. The peaks are separated by a saddle of fine-grained metasedimentary rocks (Ostler, 1993). The metasediments host mineralized southward dipping shears which are probably thrust faults.

The shears contain irregular segregations of quartz and graphite that are associated with disseminated pyrite containing gold. Minor amounts of galena, sphalerite, chalcopyrite, bornite, covelite and arsenopyrite occur with the pyrite.

Gold at the Lumby Gold Mine occurs as fine-grained native gold within pyrite grains and along hairline fractures within the shear zones (Drummond and Howard, 1993).

Underground sampling by Bradley (1990) along the 808 m level revealed the following:

	Width	Average Assay	Assay Range
No.140 Cross-cut	20 m	0.044 oz/ton Au	0.001 to 0.313 oz/ton Au
No.190 Cross-cut	14 m	0.060 oz/ton Au	0.005 to 0.222 oz/ton Au

Bradley (1990) observed that the mineralized rock looked "shiny black". A metallurgical test (Drummond and Howard, 1993) revealed that the mineralized rock contained 5 to 6% graphite and a significant amount of sericite mica. Graphite occurred as 0.1 to 0.3 x 2.5 micron grains interlayered with mica.

Metallurgical tests regarding the recovery of graphite and sericite as concentrate byproducts have been very encouraging. Bench tests using Dow Froth 250 to float off minerals resulted in a concentrate

weighing about 15 to 20% of the original sample that contained about 5.35% graphite and 0.136 oz/ton gold. Further flotation with a collector resulted in concentration to 0.98 oz/ton gold. The tails weighed 62.8% of the original sample weight and contained 0.038 oz/ton gold (Drummond and Howard, 1993).

Drummond and Howard (1993) calculated that the average head grade in the Lumby Gold Mine was 0.275 oz/ton and that previous estimates of 0.18 oz/ton obtained from earlier sampling were due to the loss of some of the extremely fine-grained gold during processing.

2.4 Property Geology

2.4(i) Stratigraphy

The oldest rocks exposed on the Deafies Creek Property are metasediments of the Late Triassic-age Sicamous Formation which forms part of the Slocan Group. The writer believes that Slocan Group rocks underlie the whole of the property but are obscured in most areas by a layer of ablation till (Figure 7). Sparse outcrop precludes any meaningful stratigraphic study of the area.

Exposures of Sicamous Formation rocks in the northern part of the property were mapped at a scale of 1:5000 during the 1993 exploration program (Figure 7). There, a assemblage of greywacke, siltstone and pelite containing minor conglomerate was sparsely exposed in a series of road cuts and beneath upturned roots of fallen trees. This rock assemblage has been metamorphosed to slate, phyllite, metagreywacke (Units Gw and Slt, Figure 7).

Diorite is exposed in several small outcrops at the crest of the rounded hill in the north-central part of the property (Unit D, Figure 7). Float mapping indicates that these outcrops are part of an intrusive body that covers at least 150 ha (360 a) on the claims.

Similar intrusives at Saddle Mountain; located about 7.4 km (4.6 mi) south-southeast of the Deafies Creek Property, are mapped by Okulitch (1979) (Figure 4) and Jones (1959) as Cretaceous-age granitics. The writer disagrees with that designation.

The cores of the intrusions at Saddle Mountain are granoblastic diorite and quartz diorite. Within 100 m of their margins the granoblastic texture is progressively replaced by a gneissic texture. Near the northern peak, the marginal diorite gneiss is folded within the carbonate and pelitic units around second-phase structures. The central parts of the diorite intrusions contain all of the cleavages found in the sedimentary units (Ostler, 1993). From this it is concluded that diorite intrusion predated deformation which was dated by Read and Wheeler (1976) to have occurred during the Middle Jurassic Period between 178 and 164 million years ago.

Near the margins of the diorite intrusion located in the northern part of the Deafies Creek Property the writer mapped small outcrops of volcanic breccia containing diorite and Slocan Group sediment clasts (Unit Dag, Figure 7). Slocan Group siltstone outcrops along the power line in the northwestern part of the property contained this conglomerate beds with similar diorite clasts.

The writer concluded that these diorite intrusions were a member of the Late Triassic-age Sicamous Formation. They were emplaced as shallow bodies that commonly reached surface as small volcanic mounds during deposition of Slocan Group sediments.

Late Tertiary and Pleistocene weathering malted White Valley to its present shape (Holland, 1976). A thick apron of glacio-fluvial sediments covers the central part of the valley burying Slocan Group rocks located there.

Preglacial soils on the upper slopes like those around Deafies Creek were commonly left in hollows as lithified regoliths between bedrock and Holocene-age soils. Some preglacial regolith was exposed in the trenches in the 1993 soil grid-area.

2.4 (ii) Deformation and Metamorphism of Sicamous Formation Rocks

The Sicamous Formation rocks on the property have been subjected to pervasive polyphase deformation and metamorphism. At Deafies Creek, the writer recognized three phases of deformation (Figure 7).

First-phase structures are mostly outcrop-scale tight to isoclinal folds and small crenulations which are most evident in pelitic strata. Generally, the first cleavage strikes northwest-southeast and is at a low angle to bedding. Most of the metamorphic mineral growth occurred during this phase of deformation. First-phase structures and cleavages have been reoriented by second-phase structures.

Most of the major structures in the Deafies Creek area are related to the second phase of deformation (Okulitch, 1979).

The second-phase cleavage in the Deafies Creek area strikes northwest and dips either north or south. Some of the variation in its orientation may be due to subsequent brittle deformation. This cleavage is defined by micaceous development. Metamorphic mineral development associated with the second phase of deformation is less pervasive than that developed during the first phase.

The third phase of deformation is recorded as a northeasterly striking, steeply dipping fracture cleavage.

In general, metamorphic mineral development indicates that biotite grade of the greenschist facies of regional metamorphism was achieved in Sicamous Formation rocks.

2.4 (iii) Exposed Economic Mineralization

Only one sulphide mineral showing is known to exist on the Deafies Creek Property. It is located in the main trench at the centre of the 1993 soil survey (Figures 7 and 12). Two outcrops are exposed in the trench. The easterly outcrop hosts an east-west striking, steeply dipping, 50 cm (1.6 ft) thick white quartz vein in bleached greywacke. The westerly outcrop contains a 20 cm (0.6 ft) thick white quartz vein that is oriented 121°/65° NW. There is a small amount of rust on this vein due to hematite staining where it is in contact with a indurated preglacial regolith. No economic mineralization is evident in either vein.

A small pile of massive pyrrhotite cobbles is located just south of the easterly outcrop in the trench. Such mineralization is not exposed in outcrop.

Composite chip samples were taken of the two veins and the massive pyrrhotite mineralization as follows:

DK18-1 massive pyrrhotite cobbles
DK18-2 rusty quartz vein, westerly outcrop
DK18-3 rusty gouge beneath preglacial regolith

None of the samples contained any economic mineralization (Appendix B).

It has been hoped by previous workers (Halliwell and Allen, 1991 and 1992) that these outcrops were related to a large mineralized structure responsible for the airborne geophysical anomaly located at the eastern boundary of the property (Figure 3).

In the writer's opinion, the showing exposed in the main trench is valueless. Its relation to the airborne anomaly at the eastern boundary of the property remains unknown.

3.0 SOIL GEOCHEMISTRY

3.1 1993 Soil Survey

A soil survey was conducted on the north-central part of the Deafies Creek Property to attempt to relate the mineral showing in the main trench with larger geological features.

Soil lines comprising the 1993 grid were run east-west at 50 m (164 ft) intervals from a north-south base line centred on the main trench (Figures 7 and 12). A total of 5.05 km (3.08 mi) of lines were surveyed by hip chain and compass in laying out the grid, which comprised 0.25 km of base line and 4.8 km of survey lines. The grid covered 20 ha (48 A). Soil stations were located at 50 m (164 ft) intervals along the lines.

At most sample stations, soils were sufficiently developed to enable collection of a sample from an illuviated "B" horizon. Sampling depths varied from about 0.1 to 0.3 m (0.3 to 1.0 ft).

Some of the 1993 soil survey area is disturbed by access roads and skidder trails. Soil samples from these areas tend to have spurious soil-metal contents due to contamination from road fill. To differentiate between soil-metal anomalies due to contamination from natural soil-metal anomalies from those disturbed by road building, the major roads crossing the grid-area are located on the maps of the soil survey (Figure 12).

Soils on the Deafies Creek Property are typical of those formed on glaciated lower alpine slopes where a thin layer of ablation till formed the initial regolith for soil development. Periglacial processes such as cryoturbation caused mixing with underlying rock. Subsequent post-glacial organic activity and solifluction developed and complicated soil profiles. On more moderate slopes, this resulted in well-developed soil horizons and comparatively mature soil profiles derived mostly from local parent rock.

Soil-metal concentrations commonly reflect the metal content of the underlying parent rock.

Soil samples were shipped in undyed kraft paper envelopes to Chemex Labs Limited of North Vancouver, B.C. A total of 99 samples were taken from the 1993 grid-area. Methods of analysis comprise Appendix A. Soil survey results comprise part of Appendix C. Soil-metal concentration distribution curves for copper, arsenic, lead and zinc are developed in Figures 8 to 11. Concentrations for these metals are contoured on Figure 12.

Soils were also analyzed for silver, gold, bismuth, mercury, molybdenum and antimony. No significant concentrations of these other metals were found in these soils. Consequently these metals were not plotted.

A statistical analysis using the methods of LePeltier (1969) with minor graphic variation was performed on the soil geochemical data for copper, arsenic, lead and zinc (Figures 8 to 11). Through this method, graphic representations of cumulative frequency curves resulted in the separation of data into common and anomalous populations.

Accepting the assumption that the common logs of the soil data naturally tend to form a normal distribution, these populations reflect the elimination of data below the 50th., 84th. and 97.5th. centiles and represent regional background, sub-anomalous and anomalous thresholds respectively.

The upper first and second standard deviations on the trends of lognormal distributions of populations of normal soils derived from graphic analysis as follows:

	Cu ppm	As ppm	Pb ppm	Zn ppm
84th. Centile (sub-anomalous)	82	17.5	7.3	210
97.5th. Centile (anomalous)	111	29.2	11.6	260

The LePeltier method was developed as a reliable and fast method for analyzing large amounts of data generated by regional soil surveys in which large unmineralized areas contain small, comparatively mineralized regions. The first and second standard deviations were used as soil map contours to separate mineralized and unmineralized areas. The soil-metal concentration defining the contours depended on the data used.

In an extensive regional soil survey the whole of the Deafies Creek grid-area would be considered to be unmineralized and would be plotted outside the sub-anomalous and anomalous contours. By using data from a small unmineralized area such as this, the contours representing anomalous soil-metal concentrations are set at uncommonly low levels. A possible negative result is that unmineralized areas show on the resulting soil maps as soil-metal anomalies; however, an advantage is that mildly mineralized areas and features of soil-metal transport can be more precisely defined.

The soil-metal concentrations of copper, arsenic, lead and zinc are contoured on Figure 12.

3.2 Interpretation of Soil-metal Distribution Curves

The northern part of the Deafies Creek Property is underlain by Slocan Group metasediments and intrusives which have been deformed and regionally metamorphosed. If metal concentrations in these rocks evolved through concentration from the country rocks in the area then soil-metal concentrations would plot as unimodal lognormal distributions containing many normal and a few anomalously high data. Such distributions commonly reflect data from a single population that have been subjected to a common evolution.

This does not indicate that all metals should be distributed in the same way throughout the rocks. One of the normal results of deformation and metamorphism is to separate various metals and concentrate them in different lithologies throughout the stratigraphy.

The arrangement of soil-metal data from the Deafies Creek grid into near unimodal lognormal distribution curves (Figures 8 to 11) strongly suggests that soil-metal concentrations are derived from a single population, the underlying Slocan Group rocks.

Also, the curves are unimodal with a high kurtosis which is caused by a large number of the values being clustered around the median value. Consequently the sub-anomalous and anomalous threshold values which are used to generate the contours on the soil maps are also quite close to the median value. Generally, the median value is taken to be near background concentration in a regional soil survey. This suggests that none of the rocks underlying the 1993 soil grid on the Deafies Creek Property are highly mineralized. Contours on these soil maps represent soil-metal concentration on favourable areas of a slope more than significant mineralization in underlying rocks.

As has been mentioned previously, the concentrations of gold, silver, bismuth, mercury, molybdenum and antimony were very low. These metals were not discussed in detail.

3.3 Interpretation of 1993 Soil Survey Results

Elevated soil-copper contents near the main trench-area may be the result of weathering of chalcopyrite associated with pyrrhotite in the quartz bodies such as those exposed in the main trench (Figure 12). These quartz bodies may be related to a weakly mineralized northwest-southeast trending fault that extends across the western part of the grid and through the main trench. The existence of two such structures is suggested by the distributions of arsenic and lead. One is located near the main trench and the other crosses the eastern part of the 1993 grid-area (Figure 12). The eastern structure displays a mild zinc signature as well as those of arsenic and lead.

A broad band of relatively high arsenic and lead concentrations extends across the grid along the 00 m N. line. This is probably due to moderate enrichment in these metals due to illuviation in the soil at a minor break in slope.

4.0 CONCLUSIONS AND RECOMMENDATIONS

4.1 Conclusions

4.1(i) Geology

The Deafies Creek Property is underlain by a sequence of Triassic-age turbidites and pelites carbonates and high-level diorite intrusives that form part of the Slocan Group. These rocks were complexly deformed, metamorphosed and intruded by granitic rocks in the Lumby region. Deformation and metamorphism began with the folding of the Kootenay Arc at the western margin of proto-North America during the Middle Jurassic Period and culminated during the extension of the Shuswap metamorphic terrane during the Late Cretaceous Period.

During the Pleistocene Stage, glacial erosion malted the current topography of the property-area. Minor Holocene uplift and erosion has exposed some of the older stratigraphy on the property.

Only one sulphide mineral showing is known to exist on the Deafies Creek Property. It is located in the main trench at the centre of the 1993 soil survey. Two greywacke outcrops exposed in the trench contain

white quartz veins. A small pile of massive pyrrhotite cobbles is located near the vein outcrops. Reportedly, these cobbles are cleaved from the veins.

No economic mineralization was found at this showing.

4.1(ii) Soil Survey

Soils were sampled in the north-central part of the Deafies Creek Property. The 1993 soil grid was centred on the main trench which has been the focal point of past surface exploration and drilling on the property.


Samples were analyzed for copper, arsenic, silver, gold, lead, zinc, bismuth, mercury, antimony and molybdenum.

Two northwest-southeast trending fault-like structures are indicated in the grid-area by the distributions of copper, arsenic, lead and zinc in soils. These structures seem not to be highly mineralized in the grid-area. It is very unlikely that these structures are the cause of the airborne electromagnetic anomaly discovered by Quinto Mining Corp. to be trending onto the eastern part of the property.

4.2 Recommendations

Discovery of the cause of the extensive airborne electromagnetic anomaly that trends westward from the Quinto Mining Corp.'s property onto the eastern part of the Deafies Creek Property has been the main incentive for recent exploration on the Deafies Creek Property. The area where the anomaly enters the claims is blanketed by a thin but extensive soil cover. A soil survey over all of the HAZ 5 claim would be a most useful way to investigate if this airborne anomaly is accompanied by mineralized areas.

Vancouver, British Columbia
June 28, 1993



John Ostler; M.Sc., P.Geo.
Consulting Geologist

5.0 REFERENCES

- Allen, A.R.; 1990: Diamond Drilling Report: The Lumby Property Vernon Mining Division, B.C. DD Hole Z90-2 on the OK Claim; B.C. Assessment Rept., 7 p. 5 fig.
- Allen, A.R.; 1989: Diamond Drilling Report on the Lumby Property Vernon Mining Division; B.C. Assessment Rept., 8 p. 2 fig.
- Bradley, M.; 1990: Report on Chip/Channel Sampling and Geological Mapping of the 140 East and 190 East Cross-cuts, 808 m Level Underground, Chaput 5 Claim: Report to The Quinto Mining Corporation.
- Douglas, R.J.W. et al.; 1970: Geology and Economic Minerals of Canada; Dept. Energy, Mines and Res., Economic Geology Rept. No.1, pp. 367-420.
- Drummond, A.D. and Howard, D.A.; 1993: Report on the Exploration Potential of the Muscovite-Graphite-Gold Deposit of the Quinto Mining Corporation; Report to The Quinto Mining Corporation.
- Halliwell, D.R. and Allen, A.R.; 1992: Geology, Geophysics and Geochemistry Report: The Lumby Property, Claims OK, HAZ 5; B.C. Assessment Rept. No. ?, 17 p. 11 fig.
- Halliwell, D.R. and Allen, A.R.; 1991: Geology and Prospecting Report, The Lumby Property, OK and Haz 5 Claims; B.C. Assessment Rept. No.?, 10 p. 6 fig.
- Holland, S.S.; 1976: Landforms of British Columbia, A Physiographic Outline; B.C. Min. Energy, Mines and Petr. Res., Bull. 48, pp. 73-74.
- Jones, A.G.; 1959: Vernon Map-area, British Columbia; Geol. Surv. Canada, Mem. 296.
- Kuran, D.L.; 1986: Report on the Lumby Project; Report to The Quinto Mining Corporation.
- Lepeltier, C.; 1969: A Simplified Statistical Treatment of Geochemical Data by Graphical Representation; Econ. Geology, Vol. 64, pp. 538-550.
- Okulitch, A.V.; 1979: Thompson-Shuswap-Okanagan Map-area, British Columbia; Geol Surv. Canada, Open File 637, 5 maps.
- Ostler, John; 1993: Geological and Geochemical Exploration on the B.S 3 and Hol 3 Claims of the Saddle Mountain Property; B.C. Assessment Rept., 34 p. 16 fig.

Read, P.B. and Wheeler, J.O.;1976: Geology: Lardeau West-half, British Columbia; Geol. Surv. Canada, Open File 432, 1 map + notes.

6.0 ITEMIZED COST STATEMENT FOR THE MAY-JUNE, 1993 EXPLORATION PROGRAM

	<u>Saddle Mountain</u>		<u>Deafies Creek</u>		<u>Whole Program</u>	
Wages:						
J. Ostler; M.Sc., P.Geo.						
31 days @ \$300/day\$ 7888.41		\$ 1411.59		\$ 9300.00	
D. Jones, B.Sc.; 19 days @ \$200/day.\$ 3330.66		\$ 469.34		\$ 3800.00	
D. Nunuk, B.Sc.; 16 days @ \$200/day.\$ 2804.77		\$ 395.23		\$ 3200.00	
John Hilton 3 days orientation + travel @ \$225/day\$ 591.63		\$ 83.37		\$ 675.00	
	\$14615.47	\$14615.47	\$ 2359.53	\$ 2359.53	\$16975.00	\$16975.00
Transport:						
Truck;						
1 1-ton 4X4 pick-up, 0.67 month @ \$2400/month.\$ 1577.68		\$ 222.32		\$ 1800.00	
Car Rental (John Hilton) 3 days @ \$25/day.\$ 65.74		\$ 9.26		\$ 75.00	
Gasoline and oil\$ 310.51		\$ 43.75		\$ 354.26	
Highway tolls.\$ 35.06		\$ 4.94		\$ 40.00	
	\$ 1988.99	\$ 1988.99	\$ 280.27	\$ 280.27	\$ 2269.26	\$ 2269.26
Camp and Equipment:						
Survey and sampling equipment 0.67 month @ \$600/month.\$ 350.60		\$ 49.40		\$ 400.00	
Camp + survey supplies\$ 265.49		\$ 37.41		\$ 302.90	
	\$ 616.09	\$ 616.09	\$ 86.81	\$ 86.81	\$ 702.90	\$ 702.90
Crew Costs:						
Hotel 15 nights @ \$65.23/night\$ 857.63		\$ 120.85		\$ 978.48	
L.D. telephone\$ 17.53		\$ 2.47		\$ 20.00	
Meals in Transit\$ 1062.03		\$ 149.66		\$ 1211.69	
	\$ 1937.19	\$ 1937.19	\$ 272.98	\$ 272.98	\$ 2210.17	\$ 2210.17
Balances Carried Forward		\$19157.74		\$ 2999.59		\$22157.33

	<u>Saddle Mountain</u>		<u>Deafies Creek</u>		<u>Whole Program</u>	
Balances Carried Forward		\$19157.74		\$ 2999.59		\$22157.33
Shipping and Assay:						
Shipping		\$ 95.58		\$ 18.91		\$ 114.49
Assay of 12 rocks @ \$27.65/sample.		\$ 248.85		\$ 82.95		\$ 331.80
Analysis of 611 soils @ \$14.55/sample.		<u>\$ 7449.57</u>		<u>\$ 1440.48</u>		<u>\$ 8890.05</u>
	\$ 7794.00	\$ 7794.00	\$ 1542.34	\$ 1542.34	\$ 9336.34	\$ 9336.34
Report Production:						
Drafting 113 hr @ \$25/hr		\$ 2715.00		\$ 810.00		\$ 3525.00
Published maps, enlargements, copy of base maps on mylar		\$ 458.33		\$ 126.11		\$ 584.44
Photocopy of text and small diagrams		\$ 129.00		\$ 92.00		\$ 221.00
Copy of large maps onto paper.		\$ 123.12		\$ 73.88		\$ 197.00
Report covers.		<u>\$ 38.50</u>		<u>\$ 31.50</u>		<u>\$ 70.00</u>
	\$ 3463.95	\$ 3463.95	\$ 1133.49	\$ 1133.49	\$ 4597.44	\$ 4597.44
Cost of May - June, 1993 work.		\$30415.69		\$ 5775.42		\$36091.11
G.S.T.; 7% of \$36091.11.		<u>\$ 2129.10</u>		<u>\$ 397.29</u>		<u>\$ 2526.39</u>
Total Exploration cost		\$32544.79		\$ 6172.71		\$38617.50

APPENDIX A
METHODS OF SOIL ANALYSIS AND ROCK ASSAY



Chemex Labs Ltd.

Analytical Chemists * Geochemists * Registered Assayers
212 Brooksbank Ave., North Vancouver
British Columbia, Canada V7J 2C1
PHONE: 604-984-0221

To: ZICTON GOLD LTD.

504 - 455 GRANVILLE ST.
VANCOUVER, BC
V6C 1T1

A9315497

Comments: ATTN: TOM KENNEDY CC: JOHN OSTLER

CERTIFICATE

A9315497

ZICTON GOLD LTD.

Project: DEAFIES CREEK
P.O. #:

Samples submitted to our lab in Vancouver, BC.
This report was printed on 17-JUN-93.

SAMPLE PREPARATION

CHEMEX CODE	NUMBER SAMPLES	DESCRIPTION
208	3	Assay ring to approx 150 mesh
226	3	0-5 lb crush and split

ANALYTICAL PROCEDURES

CHEMEX CODE	NUMBER SAMPLES	DESCRIPTION	METHOD	DETECTION LIMIT	UPPER LIMIT
998	3	Au oz/T: 1 assay ton	FA-AAS	0.001	20.00
1956	3	Ag ppm: ICP - assay digestion	ICP-AES	2	500
1951	3	Co %: ICP - assay digestion	ICP-AES	0.001	2.50
1952	3	Cu %: ICP, assay digestion	ICP-AES	0.001	2.50
1957	3	Fe %: ICP - assay digestion	ICP-AES	0.01	50.0
1958	3	Mn %: ICP - assay digestion	ICP-AES	0.001	2.50
1959	3	Mo %: ICP - assay digestion	ICP-AES	0.001	2.50
1953	3	Ni %: ICP, assay digestion	ICP-AES	0.001	2.50
1954	3	Pb %: ICP, assay digestion	ICP-AES	0.001	2.50
1955	3	Zn %: ICP, assay digestion	ICP-AES	0.001	2.50



Chemex Labs Ltd.

Analytical Chemists * Geochemists * Registered Assayers
212 Brooksbank Ave., North Vancouver
British Columbia, Canada V7J 2C1
PHONE: 604-984-0221

To: ZICTON GOLD LTD.

504 - 455 GRANVILLE ST.
VANCOUVER, BC
V6C 1T1

A9315659

Comments: ATTN: TOM KENNEDY CC: JOHN OSTLER

CERTIFICATE

A9315659

ZICTON GOLD LTD.

Project: DEAFIES CREEK
P.O. #:

Samples submitted to our lab in Vancouver, BC.
This report was printed on 22-JUN-93.

SAMPLE PREPARATION

CHEMEX CODE	NUMBER SAMPLES	DESCRIPTION
201	59	Dry, sieve to -80 mesh
229	59	ICP - AQ Digestion charge

ANALYTICAL PROCEDURES

CHEMEX CODE	NUMBER SAMPLES	DESCRIPTION	METHOD	DETECTION LIMIT	UPPER LIMIT
100	59	Au ppb: Fuse 10 g sample	FA-AAS	5	10000
2118	59	Ag ppm: 32 element, soil & rock	ICP-AES	0.2	200
2120	59	As ppm: 32 element, soil & rock	ICP-AES	2	10000
2123	59	Bi ppm: 32 element, soil & rock	ICP-AES	2	10000
2128	59	Cu ppm: 32 element, soil & rock	ICP-AES	1	10000
2131	59	Hg ppm: 32 element, soil & rock	ICP-AES	1	10000
2136	59	Mo ppm: 32 element, soil & rock	ICP-AES	1	10000
2140	59	Pb ppm: 32 element, soil & rock	ICP-AES	2	10000
2141	59	Sb ppm: 32 element, soil & rock	ICP-AES	2	10000
2149	59	Zn ppm: 32 element, soil & rock	ICP-AES	2	10000

Note: This soil analysis procedure is the same for certificates

A9315659

A9315660

APPENDIX B
ROCK ASSAYS



Chemex Labs Ltd.

Analytical Chemists * Geochemists * Registered Assayers
212 Brooksbank Ave., North Vancouver
British Columbia, Canada V7J 2C1
PHONE: 604-984-0221

To: ZICTON GOLD LTD.

504 - 455 GRANVILLE ST.
VANCOUVER, BC
V6C 1T1

Project: DEAFIES CREEK
Comments: ATTN: TOM KENNEDY CC: JOHN OSTLER

Page Number :1
Total Pages :1
Certificate Date: 17-JUN-93
Invoice No. :19315497
P.O. Number :
Account :HRG

CERTIFICATE OF ANALYSIS A9315497

SAMPLE	PREP CODE	Au oz/T	Ag ppm	Co %	Cu %	Fe %	Mn %	Mo %	Ni %	Pb %	Zn %
DK18-1	208 226	< 0.001	8	0.006	0.036	36.9	0.008	0.003	0.001	0.001	0.030
DK18-2	208 226	< 0.001	< 2	< 0.001	0.003	2.39	0.014	< 0.001	< 0.001	0.006	< 0.001
DK18-3	208 226	0.003	4	0.028	0.106	36.1	0.004	< 0.001	0.013	0.002	0.007

CERTIFICATION: *John Ostler*

APPENDIX C
SOIL ANALYSES



Chemex Labs Ltd.

Analytical Chemists * Geochemists * Registered Assayers
212 Brooksbank Ave., North Vancouver
British Columbia, Canada V7J 2C1
PHONE: 604-984-0221

To: ZICTON GOLD LTD.
504 - 455 GRANVILLE ST.
VANCOUVER, BC
V6C 1T1

Page Number : 1
Total Pages : 2
Certificate Date: 22-JUN-93
Invoice No. : I9315659
P.O. Number :
Account : HRG

Project : DEAFIES CREEK
Comments : ATTN: TOM KENNEDY CC: JOHN OSTLER

CERTIFICATE OF ANALYSIS A9315659

SAMPLE	PREP CODE	Au ppb FA+AA	Ag ppm	As ppm	Bi ppm	Cu ppm	Hg ppm	Mo ppm	Pb ppm	Sb ppm	Zn ppm
ON 050E	201 229	< 5	< 0.2	28	< 2	88	< 1	1	12	2	172
ON 100E	201 229	< 5	< 0.2	22	< 2	53	< 1	1	8	2	160
ON 150E	201 229	< 5	< 0.2	16	< 2	56	< 1	1	8	< 2	186
ON 200E	201 229	< 5	< 0.2	24	< 2	42	< 1	< 1	10	2	122
ON 250E	201 229	< 5	< 0.2	18	< 2	82	< 1	2	8	2	152
ON 300E	201 229	< 5	< 0.2	12	< 2	64	< 1	1	8	< 2	148
ON 350E	201 229	< 5	< 0.2	14	< 2	85	< 1	< 1	8	< 2	196
ON 400E	201 229	< 5	< 0.2	14	< 2	37	< 1	1	6	< 2	132
ON 050W	201 229	< 5	< 0.2	28	< 2	127	< 1	1	8	2	182
ON 100W	201 229	< 5	< 0.2	14	< 2	92	< 1	1	8	< 2	258
ON 150W	201 229	< 5	< 0.2	22	< 2	70	< 1	1	14	< 2	176
ON 200W	201 229	< 5	< 0.2	20	< 2	54	< 1	1	8	< 2	184
ON 250W	201 229	< 5	< 0.2	12	< 2	36	< 1	1	8	< 2	224
ON 300W	201 229	< 5	< 0.2	32	< 2	60	< 1	2	10	2	214
ON 350W	201 229	< 5	< 0.2	16	< 2	75	< 1	< 1	6	< 2	166
ON 400W	201 229	< 5	< 0.2	6	< 2	33	< 1	2	8	< 2	260
50N 050E	201 229	< 5	< 0.2	14	< 2	66	< 1	1	6	< 2	172
50N 100E	201 229	< 5	< 0.2	30	< 2	68	< 1	1	6	< 2	158
50N 150E	201 229	< 5	< 0.2	22	< 2	63	< 1	1	8	2	146
50N 200E	201 229	< 5	< 0.2	18	< 2	32	< 1	1	10	< 2	196
50N 250E	201 229	< 5	< 0.2	18	< 2	19	< 1	1	8	< 2	148
50N 300E	201 229	< 5	< 0.2	14	< 2	95	< 1	2	6	< 2	148
50N 350E	201 229	< 5	< 0.2	12	< 2	22	< 1	< 1	8	< 2	146
50N 400E	201 229	< 5	< 0.2	4	< 2	27	< 1	< 1	< 2	< 2	182
50N 000W	201 229	< 5	< 0.2	6	< 2	64	< 1	< 1	4	< 2	128
50N 050W	201 229	< 5	< 0.2	20	< 2	120	< 1	1	10	< 2	204
50N 100W	201 229	< 5	< 0.2	12	< 2	77	< 1	< 1	4	< 2	152
50N 150W	201 229	< 5	< 0.2	16	< 2	101	< 1	1	6	< 2	174
50N 200W	201 229	< 5	< 0.2	18	< 2	79	< 1	1	8	< 2	148
50N 250W	201 229	< 5	< 0.2	16	< 2	46	< 1	1	6	< 2	174
50N 300W	201 229	< 5	< 0.2	12	< 2	52	< 1	1	6	< 2	184
50N 350W	201 229	< 5	< 0.2	4	< 2	50	< 1	< 1	4	< 2	192
50N 400W	201 229	< 5	< 0.2	12	< 2	70	< 1	2	8	< 2	156
100N 000E	201 229	< 5	< 0.2	10	< 2	64	< 1	1	6	< 2	132
100N 050E	201 229	< 5	0.4	14	< 2	78	< 1	1	6	< 2	222
100N 100E	201 229	< 5	< 0.2	8	< 2	78	< 1	< 1	4	< 2	146
100N 150E	201 229	< 5	< 0.2	6	< 2	64	< 1	1	6	< 2	198
100N 200E	201 229	< 5	< 0.2	< 2	< 2	65	< 1	1	10	< 2	228
100N 250E	201 229	< 5	< 0.2	16	< 2	34	< 1	< 1	4	2	160
100N 300E	201 229	< 5	0.2	4	4	39	< 1	1	4	< 2	144

CERTIFICATION:

John A. Beckler



Chemex Labs Ltd.

Analytical Chemists * Geochemists * Registered Assayers
 212 Brooksbank Ave., North Vancouver
 British Columbia, Canada V7J 2C1
 PHONE: 604-984-0221

To: ZICTON GOLD LTD.

504 - 455 GRANVILLE ST.
 VANCOUVER, BC
 V6C 1T1

Project : DEAFIES CREEK
 Comments: ATTN: TOM KENNEDY CC: JOHN OSTLER

Page Number :2
 Total Pages :2
 Certificate Date: 22-JUN-93
 Invoice No. : 19315659
 P.O. Number :
 Account : HRG

CERTIFICATE OF ANALYSIS A9315659

SAMPLE	PREP CODE	Au ppb FA+AA	Ag ppm	As ppm	Bi ppm	Cu ppm	Hg ppm	Mo ppm	Pb ppm	Sb ppm	Zn ppm
100N 350E	201 229	< 5	< 0.2	4	< 2	31	< 1	1	4	< 2	174
100N 400E	201 229	< 5	< 0.2	14	< 2	32	< 1	1	6	< 2	160
100N 050W	201 229	< 5	< 0.2	12	< 2	63	< 1	< 1	4	< 2	172
100N 100W	201 229	< 5	< 0.2	18	< 2	60	< 1	< 1	2	< 2	154
100N 150W	201 229	< 5	< 0.2	6	< 2	78	< 1	< 1	6	< 2	132
100N 200W	201 229	< 5	< 0.2	12	< 2	43	< 1	< 1	6	< 2	202
100N 250W	201 229	< 5	< 0.2	4	< 2	43	< 1	2	12	< 2	208
100N 300W	201 229	< 5	< 0.2	< 2	< 2	61	< 1	1	6	< 2	204
100N 350W	201 229	< 5	< 0.2	6	< 2	40	< 1	1	6	< 2	194
100N 400W	201 229	< 5	< 0.2	4	< 2	50	< 1	1	4	< 2	160
150N 000W	201 229	< 5	< 0.2	26	< 2	84	< 1	1	6	< 2	234
150N 050W	201 229	< 5	< 0.2	4	< 2	53	< 1	< 1	4	< 2	144
150N 100W	201 229	< 5	< 0.2	2	< 2	71	< 1	1	4	< 2	128
150N 150W	201 229	< 5	< 0.2	6	< 2	95	< 1	1	4	< 2	126
150N 200W	201 229	< 5	< 0.2	10	< 2	103	< 1	1	6	< 2	142
150N 250W	201 229	< 5	< 0.2	6	< 2	74	< 1	< 1	4	< 2	166
150N 300W	201 229	< 5	< 0.2	4	< 2	54	< 1	1	4	< 2	206
150N 350W	201 229	< 5	< 0.2	4	< 2	26	1	1	6	< 2	228
150N 400W	201 229	< 5	< 0.2	4	< 2	35	< 1	< 1	6	< 2	166

CERTIFICATION:

John Ostler



Chemex Labs Ltd.

Analytical Chemists * Geochemists * Registered Assayers
 212 Brooksbank Ave., North Vancouver
 British Columbia, Canada V7J 2C1
 PHONE: 604-984-0221

To: ZICTON GOLD LTD.

504 - 455 GRANVILLE ST.
 VANCOUVER, BC
 V6C 1T1

Project: DEAFIES CREEK
 Comments: ATTN: TOM KENNEDY CC: JOHN OSTLER

Page Number : 1
 Total Pages : 1
 Certificate Date: 23-JUN-93
 Invoice No. : 19315660
 P.O. Number :
 Account : HRG

CERTIFICATE OF ANALYSIS A9315660

SAMPLE	PREP CODE	Au ppb FA+AA	Ag ppm	As ppm	Bi ppm	Cu ppm	Hg ppm	Mo ppm	Pb ppm	Sb ppm	Zn ppm
50S 000E	201 229	< 5	< 0.2	8	< 2	63	< 1	1	6	< 2	186
50S 050E	201 229	< 5	< 0.2	12	< 2	82	< 1	1	4	< 2	148
50S 100E	201 229	< 5	< 0.2	10	< 2	37	< 1	< 1	2	< 2	190
50S 150E	201 229	< 5	< 0.2	12	< 2	52	< 1	1	6	< 2	162
50S 200E	201 229	< 5	< 0.2	12	< 2	56	< 1	1	6	2	244
50S 250E	201 229	< 5	< 0.2	6	< 2	56	< 1	1	4	< 2	194
50S 300E	201 229	< 5	< 0.2	18	< 2	81	< 1	2	10	< 2	170
50S 350E	201 229	< 5	< 0.2	8	< 2	64	< 1	< 1	4	< 2	70
50S 400E	201 229	< 5	< 0.2	4	< 2	20	< 1	< 1	4	< 2	116
50S 050W	201 229	< 5	< 0.2	12	< 2	78	< 1	1	6	< 2	172
50S 100W	201 229	< 5	< 0.2	14	< 2	110	< 1	1	8	< 2	214
50S 150W	201 229	< 5	< 0.2	12	< 2	77	< 1	1	6	< 2	162
50S 200W	201 229	< 5	< 0.2	10	< 2	41	< 1	1	6	< 2	148
50S 250W	201 229	< 5	< 0.2	10	< 2	31	< 1	1	4	< 2	192
50S 300W	201 229	< 5	< 0.2	10	< 2	53	< 1	1	6	< 2	208
50S 350W	201 229	< 5	< 0.2	10	< 2	61	< 1	1	8	< 2	190
50S 400W	201 229	< 5	< 0.2	6	< 2	54	< 1	1	8	< 2	208
100S 000E	201 229	< 5	< 0.2	4	< 2	20	< 1	1	4	< 2	152
100S 050E	201 229	< 5	< 0.2	8	< 2	46	< 1	< 1	4	< 2	134
100S 100E	201 229	< 5	< 0.2	10	< 2	57	< 1	1	8	< 2	236
100S 250E	201 229	< 5	< 0.2	10	< 2	85	< 1	1	14	< 2	256
100S 300E	201 229	< 5	< 0.2	12	< 2	94	< 1	1	8	< 2	248
100S 350E	201 229	< 5	< 0.2	60	< 2	120	< 1	1	12	< 2	494
100S 400E	201 229	< 5	< 0.2	20	< 2	107	< 1	< 1	2	< 2	98
100S 050W	201 229	< 5	< 0.2	12	< 2	62	< 1	1	12	< 2	196
100S 100W	201 229	< 5	< 0.2	12	< 2	58	< 1	< 1	8	< 2	202
100S 150W	201 229	< 5	< 0.2	10	< 2	34	< 1	< 1	2	< 2	154
100S 200W	201 229	< 5	< 0.2	14	< 2	71	< 1	< 1	4	< 2	152
100S 250W	201 229	< 5	< 0.2	6	< 2	30	< 1	< 1	2	< 2	164
100S 300W	201 229	< 5	< 0.2	12	< 2	46	< 1	1	6	< 2	170
100S 350W	201 229	< 5	< 0.2	16	< 2	89	< 1	1	8	< 2	170
100S 400W	201 229	< 5	< 0.2	12	< 2	65	< 1	1	4	< 2	148
150N 050E	201 229	< 5	< 0.2	4	< 2	55	< 1	< 1	8	< 2	200
150N 100E	201 229	< 5	< 0.2	10	< 2	42	< 1	< 1	6	< 2	236
150N 150E	201 229	< 5	< 0.2	4	< 2	56	< 1	< 1	8	< 2	234
150N 200E	201 229	< 5	< 0.2	4	< 2	22	< 1	< 1	2	< 2	188
150N 250E	201 229	< 5	< 0.2	6	< 2	37	< 1	< 1	2	< 2	120
150N 300E	201 229	< 5	< 0.2	4	< 2	14	< 1	1	8	< 2	186
150N 350E	201 229	< 5	< 0.2	6	< 2	32	< 1	< 1	4	< 2	174
150N 400E	201 229	< 5	< 0.2	4	< 2	52	< 1	< 1	4	< 2	114

CERTIFICATION:

APPENDIX D
CERTIFICATE OF QUALIFICATION

I, John Ostler, of 2224 Jefferson Avenue in the City of West Vancouver, Province of British Columbia do hereby certify:

That I am a consulting geologist with business address at 2224 Jefferson Avenue, West Vancouver, British Columbia;

That I am a graduate of the University of Guelph in Ontario where I obtained my Bachelor of Arts degree in Geography (Geomorphology) and Geology in 1973 and that I am a graduate of Carleton University of Ottawa, Ontario where I obtained my Master of Science degree in Geology in 1977;


That I am licensed to practice as a Professional Geoscientist by the Association of Professional Engineers and Geoscientists of British Columbia and as a Professional Geologist by the Association of Professional Engineers, Geologists and Geophysicists of Alberta, and that I am a Fellow of the Geological Association of Canada;

That I have been engaged in the study and practice of the geological profession for over 20 years;


That this report is based on data in literature and a personal examination of the Deafies Creek Claim Group located in the Vernon Mining Division of British Columbia conducted on May 26 to 10, 1993;

That I have no interest in the Deafies Creek Property nor in the securities of Zicton Gold Limited nor do I expect to receive any.

Dated at West Vancouver, British Columbia this 28th day of June, 1993.



John Ostler; M.Sc., P.Geo.
Consulting Geologist



DEAFIES CREEK PROPERTY

ZICTON GOLD LIMITED

**GENERAL LOCATION
DEAFIES CREEK PROPERTY**

50°18.6'N., 118°59'W.

VERNON M.D., B.C.

N.T.S. 82L/6, L/7

JOHN OSTLER; M.Sc., P.Geo.

JUNE, 1993






Figure 1





LEGEND

- ADITS, MILL
- TARGET ZONE
- 1 MINE ZONE; Ag, Pb, Zn, Cu, Au VEINS
- 2 PLATEAU ZONE; Au VEINS, PROPOSED MINING
- 3 CONTACT ZONE; Ag-Au VEIN, Au SOIL ANOMALY
- 4 CLIFF ZONE; Au VEINS, Au SOIL, GROUND/AIRBORNE VLF/EM ANOMALY
- 5 SADDLE NORTH ZONE; AIRBORNE EM ANOMALY
- 6 COOPER'S ZONE; 10-30 m WIDE QUARTZ/SULPHIDE VEIN
- 7 DEAFIES ZONE; AIRBORNE VLF ANOMALY
- 8 VANCE ZONE; AIRBORNE EM ANOMALY
- 9 SADDLE ZONE; Au SOIL, GROUND EM ANOMALY, Au FLOAT
- 10 TEACHERS ZONE; Zn, As, Ag, Au VEINS, Au SOIL, AIRBORNE VLF ANOM

GEOLOGY

- Kg GRANDIORITE
- Jgn GRANITE, QUARTZ MONZONITE
- eJdi DIORITE
- uTsp VOLCANICS AND SEDIMENTS

NOTE: This figure is adapted from Allen, 1987 Revised, Figure 8.







Figure 3



ZICTON GOLD LIMITED
AIRBORNE ELECTROMAGNETIC SURVEY
 DEAFIES CREEK PROPERTY
 50°18.6'N., 118°59'W.
 VERNON M.D., B.C. N.T.S. 82L/6, L/7
 JOHN OSTLER; M.Sc., P.Geo. JUNE, 1993



NOTE: For legend, see Figure 4A





Figure 4



ZICTON GOLD LIMITED

REGIONAL GEOLOGY
from G.S.C. OPEN FILE 637
DEAFIES CREEK PROPERTY

50°18.6'N., 118°59'W.

VERNON M.D., B.C. N.T.S. 82L/6, L/7
 JOHN OSTLER; M.Sc., P.Geo. JUNE, 1993

**FIGURE 4A
LEGEND TO FIGURE 4
(UPPER PART)**

PHANEROZOIC

CENOZOIC

TERTIARY OR QUATERNARY
PLIOCENE OR PLEISTOCENE

TQs CONGLOMERATE (NEAR VERNON): BASALTIC ARENITE, BRECCIA, RUBBLE, CONGLOMERATE (ALONG NORTH THOMPSON AND CLEARWATER RIVERS).

TERTIARY

MIOCENE AND/OR PLEISTOCENE (MAY INCLUDE PLEISTOCENE)

mTv PLATEAU LAVA: OLIVINE BASALT, ANDESITE, RELATED ASH AND BRECCIA; BASALTIC ARENITE; MINOR BASAL SEDIMENTS; (MAY INCLUDE YOUNGER VALLEY BASALTS).

Eocene and (?) Oligocene

KAMLOOPS GROUP (PRINCETON GROUP IN SOUTHWEST CORNER; SKULL HILL FORMATION ALONG NORTH THOMPSON RIVER).

eTkV ANDESITE, BASALT, DACITE, TRACHYTE FLOWS AND DYKES, BRECCIA, TUFF, AGGLOMERATE.

KAMLOOPS GROUP (CHU CHUA FORMATION ALONG NORTH THOMPSON RIVER; TRANQUILLE BEDS NEAR WESTERNMOST SOUTH THOMPSON RIVER; INCLUDES UNIT **Tcg** ON MAP A).

eTks SANDSTONE, CONGLOMERATE, SHALE; MINOR COAL, TUFF ARKOSE.

UNCONFORMITY

PALEOCENE OR EOCENE

pTy SYENITE, GRANITE; MINOR MONZONITE, SHONKINITE.

MESOZOIC

CRETACEOUS

Kg GRANITE, GRANODIORITE; LESSER QUARTZ MONZONITE AND QUARTZ DIORITE.

BALDY BATHOLITH AND SATELLITIC STOCKS.

Kqm QUARTZ MONZONITE, GRANODIORITE; MINOR PEGMATITE.

EARLY CRETACEOUS

SALMON ARM, DEEP CREEK, NISCONLITH AND SCOTCH CREEK PLUTONS.

eKgd GRANODIORITE, GRANITE, QUARTZ MONZONITE; MINOR DIORITE, GABBRO, QUARTZ, DIORITE.

RAFT BATHOLITH

eKqm QUARTZ MONZONITE, GRANODIORITE; MINOR PEGMATITE AND DIORITE.

JURASSIC OR CRETACEOUS

 SYENITE AND FELSITE DYKES.

JURASSIC

Jgn MASSIVE AND FOLIATED, SYNTECTONIC PEGMATITE, APLITE, LEUCOCRATIC GRANITE AND QUARTZ MONZONITE BORDERING AND WITHIN SHUSWAP METAMORPHIC COMPLEX AND OKANAGAN PLUTONIC AND METAMORPHIC COMPLEX; SILVER STAR INTRUSIONS; (MAY INCLUDE ORTHOgneiss OF PALAEOZOIC AND PROTEROZOIC AGES).

LATE JURASSIC

VALHALLA PLUTONIC ROCKS

Jgd GRANODIORITE, GRANITE; MINOR GABBRO, DIORITE, QUARTZ DIORITE.

EARLY JURASSIC

LONG RIDGE PLUTON

EJg FOLIATED, LINEATED GRANITE (MAY INCLUDE PALAEOZOIC PLUTONIC ROCKS).

NELSON PLUTONIC ROCKS; THUYA BATHOLITH AND SATELLITIC STOCKS.

EJgd QUARTZ DIORITE, GRANODIORITE; MINOR DIORITE, GRANITE, AMPHIBOLITE, GABBRO AND ULTRAMAFIC ROCKS.

EJdi DIORITE; MINOR QUARTZ DIORITE AND GABBRO.

EJy SYENITE AND MONZONITE.

INTRUSIVE CONTACT

TRIASSIC AND JURASSIC

UPPER TRIASSIC AND LOWER JURASSIC

NICOLA GROUP (POSSIBLY INCLUDES SLOCAN GROUP NEAR SOUTHEAST EDGE OF AREA).

TJNV ANDESITE AND BASALT FLOW ROCKS, PORPHYRITIC AUGITE ANDESITE, BRECCIA, TUFF, AGGLOMERATE, GREENSTONE, CHLORITIC PHYLLITE; MINOR ARGILLITE, LIMESTONE, SERICITIC SCHIST.

UPPER TRIASSIC

KARNIAN AND NORIAN

NICOLA GROUP

URNs BLACK SHALE, ARGILLITE, CONGLOMERATE, LIMESTONE, SILTSTONE; MINOR TUFF AND PHYLLITE.

URNc LIMESTONE

**FIGURE 4A
LEGEND TO FIGURE 4
(MIDDLE PART)**

SLOCAN GROUP	
SICAMOUS FORMATION	
uRsc	SERICITIC, GRAPHITIC AND ARGILLACEOUS LIMESTONE; CALCAREOUS PHYLLITE, ARGILLITE.
uRsp	SHALE, ARGILLITE, MASSIVE SILTSTONE, PHYLLITE, TUFF AND CALCAREOUS PELITE; MINOR CONGLOMERATE, LIMESTONE, GREENSTONE, CHLORITIC PHYLLITE AND ANDALUCITE -, STAUROLITE - AND KYANITE - BEARING SCHIST.
uRscg	CONGLOMERATE.
PALAEOZOIC AND MESOZOIC	
OKANAGAN PLUTONIC AND METAMORPHIC COMPLEX (MAY INCLUDE METAMORPHIC EQUIVALENTS OF UNIT CP1A AND/OR OLDER ROCKS, AND TRIASSIC GNEISSIC GRANITE).	
IPMn	HORNBLende AND BIOTITE GNEISS, PARAGNEISS; MINOR SCHIST, MARBLE, QUARTZITE AND AMPHIBOLITE.
IPMnm	DIORITIC GNEISS, AMPHIBOLITE.
IPsc	MARBLE.
IPsb	QUARTZ MICA SCHIST.
PALAEOZOIC	
PERMIAN AND (?) PENNSYLVANIAN	
KASLO GROUP	
Pkvb	MASSIVE AND FOLIATED GREENSTONE, CHLORITIC PHYLLITE, AMPHIBOLITE; MINOR ULTRAMAFIC ROCKS.
Pkub	SERPENTINIZED ULTRAMAFIC ROCKS.
SLIDE MOUNTAIN GROUP	
FENNELL FORMATION	
Pf	PILLOW LAVA FLOWS, MASSIVE AND FOLIATED GREENSTONE, GREENSCHIST, ARGILLACEOUS CHERT; MINOR AMPHIBOLITE, LIMESTONE, BRECCIA.
Pft	CHERT
Pfp	ARGILLITE, SILTSTONE
Pfcg	CONGLOMERATE
Pfub	SERPENTINIZED ULTRAMAFIC ROCKS.
TSALKOM FORMATION	
Pt	GREENSTONE, CHLORITE PHYLLITE, AMPHIBOLITE; MINOR BLACK SHALE, LIMESTONE, MARBLE.
Ptub	SERPENTINIZED ULTRAMAFIC ROCKS.
Ptc	MASSIVE, WHITE LIMESTONE.
Ptcg	FOLIATED AND STRETCHED QUARTZ PEBBLE CONGLOMERATE.
Ptm	AMPHIBOLITIC GNEISS.
Ptsc	GREY, DIOPSIDIC MARBLE.
CARBONIFEROUS AND PERMIAN (MAY INCLUDE TRIASSIC)	
CHESTERIAN - MORROWAN AND WOLFCAMPIAN-GUADALUPIAN (MAY INCLUDE KARNIAN - NORIAN).	
THOMPSON ASSEMBLAGE (MAY INCLUDE UNIT uRns).	
CP1A	UNDIVIDED.
CP1As	SILICEOUS ARGILLITE, VOLCANICLASTIC SANDSTONE, QUARTZITE, SILTSTONE; MINOR LIMESTONE, SHEARED CONGLOMERATE, BRECCIA AND GREENSTONE.
CP1Av	GREENSTONE, TUFF.
CP1Ac	MASSIVE, CRYSTALLINE WHITE AND GREY LIMESTONE; MINOR CHERT PEBBLE CONGLOMERATE, ARGILLACEOUS LIMESTONE AND CHERT.
CP1Acg	CONGLOMERATE WITH LIMESTONE MATRIX.
CARBONIFEROUS	
MILFORD GROUP	
CMss	SILTSTONE, SANDSTONE, SHALE; MINOR QUARTZ GRANULE CONGLOMERATE.
CMsp	BLACK SHALE, ARGILLITE; MINOR SANDSTONE.
CMvd	GREENSTONE, CHLORITIC PHYLLITE.
MISSISSIPPIAN	
OSAGEAN - MERAMECIAN	
MILFORD GROUP	
Mmc	FINE GRAINED GREY LIMESTONE; MINOR DOLOMITE AND SHALE.
Mmcg	GRANULE TO BOULDER CONGLOMERATE, SOME WITH LIMESTONE AND GREENSTONE CLASTS.
MISSISSIPPIAN (?) OR OLDER	
OLD DAVE INTRUSIONS (INCLUDES ULTRAMAFIC ROCKS ASSOCIATED WITH UNITS COEbv AND TRJnv).	
Pub	SERPENTINITE AND SERPENTINIZED ULTRAMAFIC ROCKS; MINOR PYROXENITE AND PERIDOTITE.
CHAPPERON GROUP	
PCv	CHLORITIC PHYLLITE, GREENSTONE, MICACEOUS SCHIST; MINOR LIMESTONE AND ULTRAMAFIC ROCKS.
DEVONIAN	
LATE DEVONIAN	
MOUNT FOWLER BATHOLITH, SOUTH FOSTHALL PLUTON.	
lDgn	FOLIATED AND LINEATED LEUCOCRATIC GRANITE, GRANITIC FELDSPAR PORPHYRY, QUARTZ MONZONITE, GRANODIORITE, MINOR FEGMATITE AND QUARTZ DIORITE.

FIGURE 4A LEGEND TO FIGURE 4 (LOWER PART)

ORDOVICIAN
LATE ORDOVICIAN
LITTLE SHUSWAP GNEISS

L Ogn LEUCOCRATIC GRANITE GNEISS, QUARTZ MONZONITE GNEISS, GRANODIORITE GNEISS; MINOR DIORITE GNEISS.

CAMBRIAN AND ORDOVICIAN
EAGLE BAY FORMATION

E OEBvg FOLIATED ACID VOLCANIC ROCKS; CHERT, SILICEOUS PHYLLITE; SHEARED AND ALTERED QUARTZ FELDSPAR PORPHYRY AND/OR QUARTZ GRANULE CONGLOMERATE; GNEISSIC ACID IGNEOUS ROCKS NEAR SHUSWAP LAKE.

E OEBv GREENSTONE, CHLORITIC PHYLLITE; MINOR AGGLOMERATE, SERICITIC PHYLLITE, QUARTZITE, LIMESTONE AND TUFF.

E OEBq SERICITIC, SILICEOUS PHYLLITE, SERICITIC QUARTZITE, QUARTZ BIOTITE SCHIST, QUARTZ BIOTITE GARNET SCHIST; MINOR TUFF AND LAYERS OF UNITS **E OEBv**, **E OEBc**.

E OEBp BLACK ARGILLITE, ARGILLACEOUS PHYLLITE, SHALE; MINOR LIMESTONE.

E OEBc MASSIVE WHITE CRYSTALLINE LIMESTONE, DARK GREY FOLIATED LIMESTONE; MINOR LIMESTONE WITH CHERT NODULES.

E OEBqg CONGLOMERATE, SOME WITH BLACK QUARTZ CLASTS; MINOR BRECCIA AND AGGLOMERATE.

TSHINAKIN LIMESTONE MEMBER

E OEB_T MASSIVE WHITE CRYSTALLINE LIMESTONE; MINOR GREENSTONE AND GREENSCHIST.

SILVER CREEK FORMATION

E OScq QUARTZ BIOTITE, SERICITE AND GARNET SCHIST, MINOR QUARTZO-FELDSPATHIC BIOTITE GNEISS, PEGMATITE, AMPHIBOLITE, MARBLE.

CHASE QUARTZITE MEMBER

E OScC QUARTZITE, SILICEOUS MARBLE, CRYSTALLINE LIMESTONE; MINOR PELITIC SCHIST.

PROTEROZOIC AND PALAEOZOIC (MAY INCLUDE ARCHAEOAN)

SHUSWAP METAMORPHIC COMPLEX

E IPns UNDIVIDED: GRANITOID GNEISS, PARAGNEISS, SCHIST; MINOR QUARTZITE, MARBLE, AMPHIBOLITE.

E IPsb QUARTZ MICA SCHIST, COMMONLY GARNET-AND SILLIMANITE-BEARING.

E IPsq QUARTZITE; MINOR PELITIC SCHIST.

E IPsc MARBLE, DIOPSIDIC MARBLE; MINOR CALCIUM SILICATE GNEISS AND AMPHIBOLITE.

E IPm AMPHIBOLITE, AMPHIBOLITIC GNEISS, MINOR HORNBLENDE BIOTITE SCHIST.

E IPsqc SILICEOUS MARBLE, CALCAREOUS QUARTZITE, CALCIUM SILICATE GNEISS; MINOR PELITIC SCHIST.







E IPgdn GRANODIORITE, DIORITE AND TONALITE GNEISS; AUSEN GNEISS.

--- GEOLOGICAL BOUNDARIES (APPROXIMATE, ASSUMED).




Faults

- ▲▲▲▲ MYLONITE ZONES (TEETH ON HANGING WALL).
- ▲▲▲▲ THRUST FAULTS (APPROXIMATE, ASSUMED; TEETH ON HANGING WALL).
- ~ ~ ~ ~ HIGH ANGLE FAULTS (APPROXIMATE, ASSUMED).




PLANAR STRUCTURES

-  BEDDING (TOPS KNOWN: INCLINED, OVERTURNED).
-  BEDDING (TOPS UNKNOWN: HORIZONTAL, INCLINED, VERTICAL).
-  FOLIATION, SCHISTOSITY; GNEISSIC LAYERING OR CLEAVAGE (HORIZONTAL, INCLINED, VERTICAL); EARLIEST OR ONLY OBSERVED.
-  AXIAL PLANES (INCLINED, VERTICAL) OF MESOSCOPIC FOLDS OBSERVED TO HAVE DEFORMED BEDDING; EARLIEST OR ONLY OBSERVED.
-  AXIAL PLANES (INCLINED, VERTICAL) OF LATER MESOSCOPIC FOLDS OBSERVED TO HAVE DEFORMED BEDDING, FOLIATION OR PRE-EXISTING STRUCTURES.
-  AXIAL PLANES (INCLINED, VERTICAL) OF LATEST MESOSCOPIC FOLDS OBSERVED TO HAVE DEFORMED BEDDING AND TWO PHASES OF PRE-EXISTING STRUCTURES.

LINEAR STRUCTURES

-  LINEATIONS (PLUNGING, HORIZONTAL) FORMED BY FOLD AXES (F), BEDDING/FOLIATION INTERSECTION (X), MINERAL ALIGNMENT OR RODDING (R) AND BOUNDINASE AXES (A); (UNDETERMINED LINEATIONS NOT LABELLED); EARLIEST OR ONLY OBSERVED.
-  LINEATIONS (PLUNGING, HORIZONTAL) OBSERVED TO BE ASSOCIATED WITH LATE FOLDS OR SUPERIMPOSED UPON PRE-EXISTING STRUCTURES.
-  LINEATIONS (PLUNGING, HORIZONTAL) OBSERVED TO BE ASSOCIATED WITH LATEST FOLDS OR SUPERIMPOSED UPON TWO PHASES OF PRE-EXISTING STRUCTURES.

FOLDS

-  EARLY AXIAL TRACE (ANTIFORM: UPRIGHT, OVERTURNED OR RECUMBENT).
-  EARLY AXIAL TRACE (SYNFORM: UPRIGHT, OVERTURNED OR RECUMBENT).
-  LATE AXIAL TRACE (ANTIFORM, SYNFORM).

STRATIGRAPHIC SAMPLE SITE



-  PALAEOONTOLOGIC SAMPLE
-  RADIO-METRIC SAMPLE

FIGURE 5

TABLE OF GEOLOGICAL EVENTS AND LITHOLOGICAL UNITS
IN THE LUMBY AREA

Time	Formation or Event
Pleistocene to Recent	-valley rejuvenation
Eocene to Pleistocene	-erosion of the Shuswap Highland and broadening of valleys, -deposition of glacial sediments
Eocene	-brittle deformation and development of north-east striking fracture cleavage
Oligocene to Eocene	-extrusion of flood basalts and andesites
Palaeocene to Oligocene	-erosion of stratigraphy in the property-area unroofing during the Oligocene Stage
Jurassic to Palaeocene	-enlargement of Shuswap Metamorphic Complex (173 to 50 m.y*) -anatexis and metasomatism of Slocan Group rocks -intrusion of granitic to dioritic plutons -thrust faulting and development of gold-bearing shear zones
Triassic to Jurassic	-folding and metamorphism of Slocan Group rocks (173 to 164 m.y.*) resulting in: 1. development of structures and cleavages of the first and second phases of deformation; equivalent to Read and Wheeler's (1976) second and third phases 2. middle greenschist regional metamorphism
Triassic	-deposition of the Slocan Group a coarsening-upward, basin-filling sequence of variably carbonaceous pelite, variably calcareous siltstone and greywacke intruded by shallow dioritic bodies some of which reached surface


* million years ago





LEGEND

- ISOMAGNETIC LINES (absolute total field)
- 500 gammas
- 100 gammas
- 20 gammas
- 10 gammas
- Magnetic depression
- Flight lines 15 687
- Flight altitude 1000 feet above ground level




ZICTON GOLD LIMITED
AEROMAGNETISM:
E.M.R. MAP 8502G
DEAFIES CREEK PROPERTY

50°18.6'N., 118°59'W.

VERNON M.D., B.C. N.T.S. 82L/6, L/7
 JOHN OSTLER; M.Sc., P.Geo. JUNE, 1993

Figure 6



LEGEND

- N = 99 for the 1993 soil survey on the central part of the property
- b = median value which approximates the mean value
- b+s = first positive standard deviation
- b+2s = second positive standard deviation

NOTE: For map of distribution of copper in soils, see Figure 12.





Figure 8

CASSIAR EAST YUKON EXP. LTD.

ZICTON GOLD LIMITED
COPPER DISTRIBUTION CURVE:
CENTRAL AREA
DEAFIES CREEK PROPERTY
50°18.6' N., 118°59' W.
 VERNON M.D., B.C. N.T.S. 82L/6, L/7
 JOHN OSTLER; M.Sc., P.Geo. JUNE, 1993



LEGEND

- N = 99 for the 1993 soil survey on the central part of the property
- b = median value which approximates the mean value
- b+s = first positive standard deviation
- b+2s = second positive standard deviation

NOTE: For map of distribution of arsenic in soils, see Figure 12.





Figure 9

CASSIAR EAST YUKON EXP. LTD.

ZICTON GOLD LIMITED
ARSENIC DISTRIBUTION CURVE:
CENTRAL AREA
DEAFIES CREEK PROPERTY
50°18.6'N., 118°59'W.
 VERNON M.D., B.C. N.T.S. 82L/6, L/7
 JOHN OSTLER; M.Sc., P.Geo. JUNE, 1993



LEGEND

- N = 99 for the 1993 soil survey on the central part of the property
- b = median value which approximates the mean value
- b+s = first positive standard deviation
- b+2s = second positive standard deviation

NOTE: For map of distribution of lead in soils, see Figure 12.





Figure 10

CASSIAR EAST YUKON EXP. LTD.

ZICTON GOLD LIMITED
LEAD DISTRIBUTION CURVE:
CENTRAL AREA
DEAFIES CREEK PROPERTY
50°18.6'N., 118°59'W.

VERNON M.D., B.C. N.T.S. 82L/6, L/7
 JOHN OSTLER; M.Sc., P.Geo. JUNE, 1993



LEGEND

- N = 99 for the 1993 soil survey on the central part of the property
- b = median value which approximates the mean value
- b+s = first positive standard deviation
- b+2s = second positive standard deviation

NOTE: For map of distribution of zinc in soils, see Figure 12.




Figure 11

CASSIAR EAST YUKON EXP. LTD.

ZICTON GOLD LIMITED
ZINC DISTRIBUTION CURVE:
CENTRAL AREA
DEAFIES CREEK PROPERTY
50°18.6'N., 118°59'W.

VERNON M.D., B.C. N.T.S. 82L/6, L/7
 JOHN OSTLER; M.Sc., P.Geo. JUNE, 1993



LEGEND

TABLE OF LITHOLOGIC UNITS

AGE	FORMATION AND LITHOLOGY
Triassic	Slocan Group; Sicamous Formation
	diorite intrusions
	agglomerate and breccia with diorite and sedimentary clasts in diorite matrix
	slate and phyllite; variably carbonaceous, black weathering to dark grey
	greywacke with minor slate, conglomerate and tuff.

STRUCTURE

Geological contact	defined	approximate	assumed
	—	- - -	· · ·

Bedding

upright	overturned	tops unknown	parallel with cleavage
↕	↘	↗	↔

Cleavage

first	second	third	Vein attitude
↖	↗	↘	↔

TOPOGRAPHY

Topographic contour	Road or skidder trail	Property boundary
~	—	—




Limit of outcrop

Trench	Creek	Diamond drill hole
⊖	↘	○

Rock assay location

DK 18
○

SCALE



Magnetic declination for the western margin of N.T.S. Map 82 L/7 as of July 1, 1993. Declination decreases 7.9' annually.

Figure 7

ZICTON GOLD LIMITED





GEOLOGY
NORTH of DEAFIES CREEK

DEAFIES CREEK PROPERTY
50°18.6'N., 118°59'W.

VERNON M.D., BRITISH COLUMBIA N.T.S. L/6, L/7
JOHN OSTLER, M.Sc., P.Geo. JUNE, 1993

GEOLOGICAL BRANCH
ASSESSMENT REPORT

22,954



LEGEND

TOPOGRAPHY:
 Road
 Trench
 cut
 bulldozer tracks
 fill
 limit of outcrop
 assay location

SOIL SURVEY:
 Diamond drill hole
 1993 soil grid
 soil station
 soil analysis in parts per million
 82 ppm contour

NOTE: For location on property, see Figure 7.

SCALE

0 20 40 60 80 100 120 140 160 180 200 220 240 260 280 300 metres

0 100 200 300 400 500 600 700 800 900

N.
 20°15' N. mag.

Magnetic declination for the western margin of N.T.S. Map 82 L/7 as of July 1, 1993. Declination decreases 7.9' annually.

PROFESSIONAL
 JOHN D. OSTLER
 BRITISH COLUMBIA
 GEOSCIENTIST

ZICTON GOLD LIMITED

**COPPER, ARSENIC, LEAD and ZINC
 in SOILS: CENTRAL AREA**

DEAFIES CREEK PROPERTY
 50°18.6'N., 118°59'W.

VERNON M.D., BRITISH COLUMBIA N.T.S. L/6, L/7

JOHN OSTLER, M.Sc., P. Geo. 1993
**GEOLOGICAL BRANCH
 ASSESSMENT REPORT**

Figure 12

22,954