

GEOLOGICAL REPORT

on the

TAM MINERAL CLAIMS

**Clinton Mining Division
British Columbia**

**N.T.S. 092P/14
Latitude 51° 57' 00" N
Longitude 121° 15' 00" W**

for

**Paul Reynolds
4035 West 35th Avenue
Vancouver, B.C.
V6S 1Y7**

by

**P. REYNOLDS, B.Sc., P.Geo.
07 October 1999**

**GEOLOGICAL SURVEY BRANCH
ASSESSMENT REPORT**

26,038

TABLE OF CONTENTS

1.	SUMMARY	2
2.	INTRODUCTION	2
3.	LOCATION, ACCESS AND PHYSIOGRAPHY	2
4.	CLAIM STATUS	3
5.	HISTORY	3
6.	REGIONAL GEOLOGY	4
7.	PROPERTY GEOLOGY	5
8.	CONCLUSION AND RECOMMENDATIONS	5
9.	REFERENCES	6
10.	CERTIFICATE	7

LIST OF FIGURES

FIGURE 1	LOCATION MAP	APPENDIX II
FIGURE 2	CLAIM MAP	APPENDIX II
FIGURE 3	REGIONAL GEOLOGY	APPENDIX II
FIGURE 4	PROPERTY GEOLOGY	APPENDIX II

APPENDICES

APPENDIX I	STATEMENT OF COSTS
APPENDIX II	FIGURES 1-4

1. SUMMARY

- 1.1 The Tam property consists of five contiguous mineral claims totaling 35 units. The claims are located approximately 21 kilometres northeast of Lac La Hache, B.C. The claims are accessible by good gravel roads from Lac La Hache.
- 1.2 The property is underlain by Nicola Group volcanoclastic and sedimentary rocks of Upper Triassic age. These rocks are intruded by a number of Upper Triassic-Lower Jurassic plutons, stocks and dykes which are part of an alkaline intrusive complex underlying the Spout Lake area.
- 1.3 The claims are located at the southeast end of a large arcuate aeromagnetic anomaly which reflects magnetite rich phases in the Spout Lake alkaline intrusive complex and/or alteration zones along its margin. Past exploration work along the southern limb of this anomaly located a number of occurrences mineralized with copper plus values in gold, silver +/- molybdenum and tungsten over a strike length of 15 kilometres.
- 1.4 Mineralization on the property is clearly associated with syenitic intrusives and structures. More exploration work consisting of detailed geological mapping is required in order to understand the nature of this relationship. In addition to geological mapping, soil sampling should also be conducted.

2. INTRODUCTION

- 2.1 This report has been prepared to satisfy assessment requirements.
- 2.2 The information for the following report was obtained from sources cited under references and from fieldwork conducted by the Author and Michael H. Sanguinetti, P. Eng. from 12 October to 15 October, 1999. The fieldwork conducted consisted of preliminary geological mapping at a scale of 1:2,500.
- 2.3 The registered owner of the Tam claims is Mr. Paul Reynolds. The claims lie approximately 21 kilometres northeast of Lac La Hache, B.C. Access to the claims is by paved highway and a network of logging roads.

3. LOCATION, ACCESS AND PHYSIOGRAPHY

- 3.1 The Tam claims are located approximately 21 kilometres northeast of Lac La Hache, B.C. The claims are centered at 51° 57' 00" north latitude and 121° 15' 00" west longitude on NTS map sheet 092P/14. The claims are in the Clinton Mining Division.
- 3.2 Access to the property is provided by the Timothy Lake Road from Lac La Hache and thence along a network of logging roads. The southern part of the property is accessed from the Timothy Mountain turn-off via Fly Lake while the northern part is accessible from Rail Lake via the "1700 Road". Total road distance from Lac La Hache is approximately 30 kilometres.

3.3 The claims are located on the western edge of the Quesnel Highlands in an area characterized by relatively low, rounded hills and ridges separated by depressions often occupied by small lakes, swamps or streams. The area is well forested with Fir, Spruce and Balsam. Approximately 40% of the property has been clear cut logged.

4. CLAIM STATUS

4.1 The Tam property comprises five mineral claims totaling 35 units. Complete claim information is as follows:

<u>NAME</u>	<u>UNITS</u>	<u>TENURE NO.</u>	<u>EXPIRY DATE *</u>
Tam 1	12	348485	14 July 2000
Tam 3	20	348486	14 July 2000
Mat 1	1	348482	14 July 2000
Mat 2	1	348483	14 July 2000
Mat 3	1	348484	14 July 2000

* Includes assessment currently being applied.

4.3 All claims are recorded in the name of Mr. Paul Reynolds. Any legal aspect of claim ownership is beyond the scope of this report.

5. HISTORY

5.1 The area of the Tam claims has been explored intermittently since 1966 when the area was first worked by the Coranex syndicate. Their claims, located west and northwest of the present Tam claims, and the surrounding area were explored by the syndicate from 1966 to 1968. Exploration work by Asarco in 1969 and Amax Exploration in 1972 identified three main mineral occurrences: Peach, Miracle and Tim. The Peach and Miracle showings are located west and northwest of the Tam claims respectively. The Tim showing is located on the Tam claims. The showings consisted of chalcopyrite-pyrite and lesser magnetite in syenodiorite and/or altered andesite, proximal to syenodiorite/andesite contacts.

5.2 During the same period, Amax Exploration was exploring their own claims located to the west of the Coranex ground. They located several significant skarn altered zones well mineralized with magnetite and chalcopyrite (WC showing). One zone, the North Zone, was estimated to contain 1.5 million tons grading 0.54% copper, 0.01 ounces/ton gold and 0.10 ounces/ton silver. A second zone, the South Zone, was not as well tested.

5.3 Craigmont Mines Ltd. drilled the North Zone in 1974, extending the known mineralized strike length to 660 metres. This drilling also indicated the possibility of a mineralized zone parallel to the north zone. Both the North and South zones are open along strike and to depth.

- 5.4 The Tam claim area was re-staked in 1979 by Stallion Resources Ltd. They conducted soil sampling, trenching and limited diamond drilling, mostly in areas of induced polarization (IP) anomalies generated during the Coranex work in 1969.
- 5.5 In 1983, Stallion diamond drilled six short holes totaling 312 metres on the Tim No. 1 showing. Diamond drill hole DDH-1 was well mineralized from surface to 42.7 metres depth. This section graded 2.76% copper, 0.74 ounces/ton silver and 0.018 ounces/ton gold (Assessment Report 12192). Hole DDH-2 intersected copper mineralization from 19.8 to 33.5 metres depth. This 13.7 metre intersection returned 1.03% copper with strongly anomalous gold and silver values. DDH-1 may have been drilled partially down the volcanic-intrusive contact.
- 5.6 In 1988, Liberty Gold Corp. optioned the claims and conducted a VLF-EM and magnetometer survey over the entire property. In 1989, Liberty conducted a soil geochemical survey over the property utilizing the previous years grid. They also conducted an IP survey over the central portion of the property. Significant copper-gold geochemical anomalies were indicated. In addition, prominent IP chargeability anomalies coincident with the geochemical anomalies were noted. In 1990, they conducted 17.8 kilometres of detailed IP survey, 736 metres of percussion drilling in seven holes and 1,245 metres of NQ diamond drilling in 12 holes. Not all of this work was publicly documented.

6. REGIONAL GEOLOGY

- 6.1 The Tam claims lie within the Quesnel Trough, a 30 to 60 kilometre wide belt of Lower Mesozoic volcanic and related sedimentary rocks bounded by older sedimentary rocks of the Cache Creek Group to the east and younger Coast Intrusions to the west. In the area of the Tam claims the Quesnel Trough is dominated by Upper Triassic Nicola Group andesites, basalts, tuffs and argillites. The Nicola Group is intruded by the Upper Triassic-Lower Jurassic Spout Lake Intrusions. These include plutons that vary in composition from granodiorite to quartz diorite and small alkali stocks which vary in composition from syenite through diorite to pyroxenite.
- 6.2 The Late Jurassic-Early Cretaceous Takomkane Batholith intrudes the Nicola Group to the east of the Tim claims. The Takomkane Batholith is composed of granodiorite.
- 6.3 Portions of this area are obscured by Tertiary Plateau lavas. Bedrock exposure in the area of the Tim claims amounts to about 10%, the rest being covered by glacial drift.

7. PROPERTY GEOLOGY

7.1 Preliminary geological mapping at a scale of 1:10,000 was previously completed by the writer's over portions of the Tam property. During the fall 1998 field program the writer's conducted geological mapping at a scale of 1:2,500 over a one square kilometre area on the west-central portion of the Tam 3 claim. This mapping has been confined, for a large part, to road-cuts and old trenches because thick deposits of glacial drift cover a large portion of the property. Additional copper mineralization was documented as a result of this new work. A brief description of the geological units follows:

UNIT 1

7.2 The oldest and most abundant rocks on the property are the upper Triassic volcanic and related pyroclastic rocks of the Nicola Group. These comprise tuffs, flows and ignimbrites of andesitic composition. Bedded rocks strike west to northwest and dip steeply north. Near syenodiorite contacts, the volcanics are strongly fractured and sheared along northeast and northwest trends. The volcanics exhibit weak to moderate magnetic response. Locally, the volcanics exhibit weak to moderate epidote and weak chlorite alteration. The andesite becomes porphyritic locally.

UNIT 2

7.3 This unit consists of grey and white very fine grain groundmass with 5-15 millimetre diameter rounded fragments of white, feldspar phenocrysts.

UNIT 3

7.4 Dark greenish gray to gray, medium grained diorite. Unit 3a is similar to Unit 3 but with about 10% quartz and is labeled quartz diorite.

UNIT 4

7.5 Pink to pinkish gray medium grained syenodiorite. Along the contacts with the andesite, both the andesite and syenodiorite are altered to epidote. An andesite-syenodiorite hybrid consisting of andesite with minor (<15%) syenodiorite appears to form along the andesite-syenodiorite contact. Much of the copper mineralization is concentrated within this hybrid phase (which is often mapped as andesite). This unit is highly magnetic.

7.6 Mineralization consists of malachite and chalcopryrite in intensely epidote and quartz altered andesite, syenodiorite and syenite breccia. Much of the mineralization is concentrated in an andesite-syenodiorite hybrid that appears to form within shears and fractures. Chip and grab samples of mineralized rock, taken by the author during previous exploration programs, returned copper values ranging from trace to one percent.

8. CONCLUSION AND RECOMMENDATIONS


8.1 The Tam claims are underlain by andesites of the Nicola Group which are intruded by syenitic rocks of the Spout Lake Intrusions. Copper mineralization consisting of malachite,

10. CERTIFICATE

I, Paul Reynolds, of the city of Vancouver in the province of British Columbia do hereby certify that:

- 1) I am a Professional Geoscientist registered with the Association of Professional Engineers and Geoscientists of British Columbia.
- 2) I am a graduate of the University of British Columbia with a B.Sc. degree in geology.
- 3) I have practiced my profession as exploration geologist since graduation in 1987.
- 4) This report is based on a review of previous reports and on fieldwork conducted by the Author from 12 October to 15 October, 1999.
- 5) I am the registered owner of the Tam claims.


Dated this 7th day of October, 1999.



APPENDIX I
STATEMENT OF COSTS

STATEMENT OF COSTS
October 12 to October 15, 1998

Paul Reynolds, P. Geo.	4 days @ \$500/day	2,000
Michael H. Sanguinetti, P. Eng.	4 days @ \$500/day	2,000
Truck Rental	4 days @ \$55/day	220
Mileage	1,200 km @ \$0.20/km	240
Assays and Analyses		143
Food, Gas & Lodging		316
Field Supplies		50
Report, Drafting & Printing		1,000
TOTAL		\$5,969





GEOCHEMICAL ANALYSIS CERTIFICATE

Sanguinetti Engineering Ltd. PROJECT TAM File # 9804976
422 - 470 Granville St., Vancouver BC V6C 1V5


SAMPLE#	Mo ppm	Cu ppm	Pb ppm	Zn ppm	Ag ppm	Ni ppm	Co ppm	Mn ppm	Fe %	As ppm	U ppm	Au ppm	Th ppm	Sr ppm	Cd ppm	Sb ppm	Bi ppm	V ppm	Ca %	P %	La ppm	Cr ppm	Mg %	Ba ppm	Ti %	B ppm	Al %	Na %	K %	W ppm	Au* ppb
R 3139	3	3054	13	97	3.1	7	15	1624	4.34	9	<8	<2	<2	46	1.9	<3	<3	118	3.74	.162	4	16	.70	50	.07	3	1.23	.07	.50	<2	53
R 3140	1	1839	11	89	2.0	6	13	1848	3.77	7	<8	<2	<2	36	1.9	<3	<3	99	2.64	.153	4	16	.57	68	.06	4	1.31	.07	.60	<2	19
R 3141	4	52	<3	92	.3	12	19	1012	5.46	17	<8	<2	<2	125	.4	<3	<3	178	1.61	.244	2	14	1.69	87	.18	<3	2.25	.18	1.23	<2	6
R 3142	2	113	7	112	<.3	9	31	1044	6.09	18	<8	<2	<2	123	.5	<3	<3	197	2.31	.242	2	18	2.05	64	.20	<3	2.45	.14	1.07	<2	13
R 3143	7	217	4	42	<.3	7	31	635	4.90	14	<8	<2	<2	106	.2	<3	<3	139	1.20	.214	3	17	1.52	75	.17	<3	1.99	.15	.72	<2	15
R 3144	6	23	7	28	<.3	5	7	455	2.61	5	<8	<2	2	140	.2	<3	<3	53	1.07	.145	7	5	.52	46	.07	9	1.15	.15	.35	<2	11
RE R 3144	5	23	7	28	<.3	4	7	458	2.63	7	<8	<2	2	143	<.2	<3	<3	53	1.08	.145	8	4	.52	49	.07	9	1.19	.17	.36	<2	4
R 3145	<1	48	8	92	<.3	9	27	911	4.35	11	<8	<2	<2	139	<.2	<3	<3	127	1.51	.246	3	17	1.74	76	.17	<3	2.11	.08	1.07	<2	4
R 3146	2	171	5	107	<.3	11	27	1229	4.17	17	<8	<2	<2	161	.2	<3	<3	143	1.48	.250	3	18	2.00	113	.20	5	2.53	.08	1.47	<2	17
STANDARD C3/AU-R	26	65	37	153	5.6	33	12	739	3.21	54	17	3	21	29	22.5	17	17	77	.54	.085	17	160	.55	139	.07	20	1.79	.05	.17	15	496
STANDARD G-2	2	2	5	40	<.3	7	5	530	2.02	<2	<8	<2	4	84	<.2	<3	<3	39	.66	.094	7	77	.57	292	.10	<3	1.08	.14	.53	2	<1

ICP - .500 GRAM SAMPLE IS DIGESTED WITH 3ML 2-2-2 HCL-HNO3-H2O AT 95 DEG. C FOR ONE HOUR AND IS DILUTED TO 10 ML WITH WATER.
THIS LEACH IS PARTIAL FOR MN FE SR CA P LA CR MG BA TI B W AND MASSIVE SULFIDE AND LIMITED FOR NA K AND AL.
ASSAY RECOMMENDED FOR ROCK AND CORE SAMPLES IF CU PB ZN AS > 1%, AG > 30 PPM & AU > 1000 PPB
- SAMPLE TYPE: ROCK AU* - IGNITED, AQUA-REGIA/MIBK EXTRACT, GF/AA FINISHED.(10 GM)
Samples beginning 'RE' are Reruns and 'RRE' are Reject Reruns.

DATE RECEIVED: NOV 9 1998 DATE REPORT MAILED: *Nov 19/98* SIGNED BY: *CL* D. TOYE, C. LEONG, J. WANG; CERTIFIED B.C. ASSAYERS

APPENDIX II


FIGURES 1 – 4



MOUNT TIMOTHY PROJECT
LAC LA HACHE AREA, BRITISH COLUMBIA
 CLINTON MINING DIVISION
 NTS 092 P/14 E, W


TAM CLAIM GROUP
LOCATION MAP

JULY 1996, FIGURE 1



REYNOLDS GEOLOGICAL LTD.

**TAM PROPERTY
CLAIM MAP**



LEGEND

- QUESHEL TROUGH**
- Upper Triassic Lower Jurassic
- RJ** BASALTIC BRECCIAS, MINOR FLOWS, TUFF, SANDSTONE, CONGLOMERATE & LIMESTONE; INCLUDES COMAGMATIC ALKALIC STOCKS, SILLS & DYKES
- Upper Triassic
- UR** ARGILLITE, AUGITE-PORPHYRY BRECCIA, BASALTIC TO ANDESITIC TUFF POSSIBLE DYKES & SILLS

GOLD OCCURRENCES

- ▲ Cu-Au Porphyry
- Au Stratabound
- × Au Bearing Veins
- P Placer Au (major occurrence)




After Salekin & Simpson (1981)



REYNOLDS GEOLOGICAL LTD.	
TAM PROPERTY REGIONAL GEOLOGY	
Clinton M. D.	NTS 092P/14
DATE October 93	SCALE 1:50,000
FIGURE 3	

SAMPLE NO.	TYPE	WIDTH (m)	Cu (ppm)	Au (ppb)
3139	GRAB	-	3,054	53
3140	GRAB	-	1,839	19
3141	CHIP	1.7	52	6
3142	CHIP	1.0	113	13
3143	CHIP	1.4	217	15
3144	CHIP	0.5	23	11
3145	CHIP	2.6	48	4
3146	CHIP	2.5	171	17



LEGEND

- Road
- ~ Stream
- Lake
- Clearcut
- Outcrop
- Mal Malachite
- Cpy Chalcopyrite

- Tertiary
 - 5 Basalt
- Triassic-Jurassic
 - 4 Syenodiorite
 - 3 Diorite 3a Quartz Diorite
 - 2 Syenite Breccia
- Upper Triassic
 - 1 Andesite.
 - 1a Plagioclase feldspar crystal tuff.
 - 1b Porphyritic andesite.

GEOLOGICAL SURVEY BRANCH
 (SEPARATE REPORT)

26,038



REYNOLDS GEOLOGICAL LTD.

**TAM CLAIMS
 GEOLOGY MAP**

NTS: 092P/14	SCALE: 1:2,500	CLINTON M.D.
DATE: SEPTEMBER 99		FIGURE: 4

