



ASSESSMENT REPORT

describing

PROSPECTING, MAPPING AND GEOCHEMICAL SURVEYS

on the

OLD IRONSIDES 2 CLAIM

Tenure No. 369542

Latitude 50° 13'N; Longitude 124° 07'W

NTS 92K/1

in the

VANCOUVER MINING DIVISION

BRITISH COLUMBIA

ARND BURGERT
OCTOBER 30, 1999

**GEOLOGICAL SURVEY BRANCH
ASSESSMENT REPORT**

26,243

TABLE OF CONTENTS

	PAGE
INTRODUCTION	1
PROPERTY, LOCATION AND ACCESS	2
GEOMORPHOLOGY	5
REGIONAL GEOLOGY	6
REGIONAL MINERALIZATION	8
REGIONAL GEOCHEMISTRY	33
PROPERTY GEOLOGY	34
PROPERTY MINERALIZATION	38
PROPERTY GEOCHEMISTRY	40
CONCLUSIONS AND RECOMMENDATIONS	51
REFERENCES	52

FIGURES

NO. DESCRIPTION	PAGE
1 PROPERTY LOCATION	3
2 CLAIM LOCATION	4
3 REGIONAL GEOLOGY	7
4 REGIONAL MINERALIZATION	9
5 PROPERTY GEOLOGY	35
6 GEOCHEMISTRY HISTOGRAMS	41
7 SAMPLE LOCATIONS	43
8 COPPER SOIL GEOCHEMISTRY	45
9 BARIUM SOIL GEOCHEMISTRY	46
10 ZINC SOIL GEOCHEMISTRY	47
11 SILVER SOIL GEOCHEMISTRY	48
12 ARSENIC SOIL GEOCHEMISTRY	49

APPENDICES

	APPENDIX
AUTHOR'S STATEMENT OF QUALIFICATIONS	I
CERTIFICATES OF ASSAY	II
STATEMENT OF EXPENDITURES	III

INTRODUCTION

The Old Ironsides 2 mineral claim was staked during June, 1999 to protect a previously unstaked volcanogenic massive sulphide (VMS) base and precious metals target identified during prospecting during the summer of 1998. One claim was staked over rocks of the lower Cretaceous Gambier group.

Gambier group rocks host the Britannia deposit on Howe Sound as well as the Northair deposit near Whistler. In the Powell River region, uneconomic base metals occurrences lying within the Gambier group include the Mt. Diadem workings overlooking Jervis Inlet and the Hummingbird past producer on Goat Island in Powell Lake.

All work was conducted personally by the author, whose Statement of Qualifications appears in Appendix I.


PROPERTY, LOCATION AND ACCESS

The Old Ironsides 2 claim is located in southwest British Columbia at 50° 13'N latitude and 124° 07'W longitude on NTS mapsheet 92K/1 (Figure 1). It consists of a sixteen-unit four-post claim (Figure 2) registered with the Vancouver Mining Recorder's Office. The Old Ironsides 2 claim was staked during June 1999 to protect a geochemical soil anomaly indicated by reconnaissance soil sampling during a 1998 exploration program. The Old Ironsides 2 claim is contiguous with the Old Ironsides and Old Ironsides 3 claims, but none of the claims are grouped. Claim registration data is summarized below.

<u>Claim Name</u>	<u>Units</u>	<u>Tenure Number</u>	<u>Expiry Date*</u>
Old Ironsides 2	16	369542	June 17, 2002

*if assessment credit for work described in this report is granted

Exploration was conducted from June 21 to August 29, 1999 from a helicopter-supported fly camp near the claim's southern boundary. The work, which was delayed due to an extreme snow pack, consisted of prospecting, soil sampling and geological mapping. A Statement of Expenditures appears in Appendix III.



-  HIGHWAY
-  MAJOR ROAD
-  LAKE
-  RIVER
-  SELECTED CITY





FIGURE 1

Old Ironsides 2 Claim

PROPERTY LOCATION

DRAWN BY: AB	PRODUCED AT: BAR
DATE: OCT 12, 1999	FILE: GENERAL\BCMAP.DWG



- ROAD, DRIVEABLE
- ROAD, DEACTIVATED
- LAKE
- RIVER, CREEK
- LEGAL CORNER POST




FIGURE 2

OLD IRONSIDES 2 CLAIM

CLAIM LOCATION

DRAWN BY: AB	PRODUCED AT: 1:50000
DATE: JUL 22, 1999	FILE: IRONSIDES/SOCLAIM.OWG

GEOMORPHOLOGY

The Old Ironsides 2 Claim is situated in mountainous terrain of the Coast Ranges. Topography is steep, typically 20° to 30°, with the exception of a broad U-shaped valley at the claim's southern end. Elevations range from 760m in the valley to 1700m along the ridge near the northern claim boundary.

Creeks draining the northern end of the claim are part of the Barkshack Creek watershed, which in turn flows into the Skwakwa River and eventually Jervis Inlet. Creeks draining the southern slope drain into a tributary of the Eldred River, which flows into Goat Lake and eventually Powell Lake.


Vegetation consists of sparse stands of juvenile second growth fir, douglas fir, hemlock and western red cedar on the valley floor. The lower valley slope is covered with old growth fir and hemlock, giving way to old growth yellow cedar scrub above 1150m. In the old growth, density of underbrush varies, and with increasing elevation, vegetation gradually becomes sparser. Above 1370m, scattered buckbrush, dwarf balsam and moss dominate, while steep talus slopes and cliffs are vegetated only by lichen. Vegetation on the ridge top is limited to grasses, moss and scattered clumps of dwarf conifers.

REGIONAL GEOLOGY

The Old Ironsides 2 Claim covers steeply dipping blocks or pendants of metasedimentary and metavolcanic rocks which lie engulfed in the main mass of the Coast Plutonic Complex (Figure 3). Pendants of Gambier Group, named for their type locality on Gambier Island in Howe Sound, extend discontinuously from North Vancouver in the southeast to north of Bella Bella in the northwest.

These pendants are thought to represent fault slices along which plutonic rock was thrust upwards (Roddick, 1976). The bounding shear zones in places still exist, and in many places are flanked by diorite. The dioritic rocks may represent remnants of a primitive granitoid basement upon which sedimentary and volcanic rocks were deposited.

The metamorphic rocks have undergone deep burial and subsequent deformation, probably in response to compressive forces transmitted through the North America Plate against oceanic crust. With the eventual onset of subduction, plutonic masses, formed during the compressive stage, began their movement upwards bounded by synplutonic faults.







- Major City or Town
-  Quaternary
-  Coast Plutonic Complex
diorite, quartz diorite, granodiorite, quartz monzonite, gabbro
-  Gombier
greenstone, volcanic breccia, argillite, minor conglomerate, limestone and schist
-  Other layered rocks
Upper Karmutsen; Bowen Island; Gorbaldi; other stratified rocks, Triassic to Tertiary


FIGURE 3
OLD IRONSIDES 2 CLAIM
 REGIONAL GEOLOGY





Geology by Roddick et al, 1976; Roddick et al, 1979; Woodsworth, 1977
 DRAWN BY: AB PRODUCED AT: 1:1000000
 DATE: SEPT 9, 1999 FILE: GENERAL\REGGEO.DWG

REGIONAL MINERALIZATION

A number of significant base metals deposits occur within pendants of Gambier group (Figure 4). Some of those that are known or suspected to be volcanocenic in origin are described in this section. The descriptions are taken from the British Columbia Ministry of Energy and Mines Minfile database, where references may be found.

The most valuable deposit discovered to date in rocks of the Gambier group is the **Britannia Deposit** at Britannia Beach on Howe Sound. The Britannia district is underlain by a roof pendant of mid- Mesozoic volcanic and sedimentary rocks, within the Cenozoic- Mesozoic Coast Plutonic Complex. A broad, steeply south dipping zone of complex shear deformation and metamorphism, the Britannia shear zone, crosses the pendant in a northwest direction; all orebodies are in the shear zone. A narrow zone of foliated rocks, the Indian River shear zone, is subparallel to the Britannia shear zone and transects the northeast part of the Britannia pendant. The deformed rocks are cut by dacite dykes and several major sets of faults. The Britannia roof pendant is one of many northwest trending bodies within, and in part metamorphosed by, the Coast Plutonic Complex. The pendant is comprised of fresh to weakly metamorphosed rocks with sharp contacts against plutonic rocks,



- Major City or Town
-  Quaternary
-  Coast Plutonic Complex
diorite, quartz diorite, granodiorite, quartz monzonite, gabbro
-  Gambier
greenstone, volcanic breccia, argillite, minor conglomerate, limestone and schist
-  Other layered rocks
Upper Karmutsen; Bowen Island; Garibaldi; other stratified rocks, Triassic to Tertiary




FIGURE 4
OLD IRONSIDES 2 CLAIM
 REGIONAL MINERALIZATION

DRAWN BY: AB	PRODUCED AT: 1:1000000
DATE: SEPT 9, 1999	FILE: GENERAL\REGGEOI.DWG

and belongs to the Lower Cretaceous Gambier Group. The Coast plutonic rocks consist of older, commonly foliated bodies ranging from diorite to granodiorite and younger quartz diorite to quartz monzonite intrusions (Squamish pluton). The plutonic rocks have produced contact metamorphic aureoles up to a hundred metres wide in the Britannia pendant.

The Britannia mine area within the Britannia shear zone is dominated by strongly foliated pyroclastic rocks of dacitic to andesitic volcanism intercalated near the top and overlain by dark marine shales and siltstones. Extensive units of fine-grained andesitic rocks were formed in the mine area during hiatuses in dacitic volcanism; one hiatus occurred during the period of formation of massive sulphides and related deposits after extrusion of a dacite tuff breccia. The lower pyroclastic sequence and the upper shale-siltstone sequence are cut by many dacitic and andesitic dykes. The lower sequence is composed of pyroclastic dacite tuff breccia (locally called the Bluff tuff breccia) that commonly grades to dacitic crystal and lithic tuffs. This unit contains prominent dark, wispy fragments and grades at the top into distinctive beds which consist of intercalated black argillite and plagioclase crystal tuffs. These may be regularly interbedded, convoluted or disaggregated by soft rock deformation. Within the pyroclastic sequence there are also minor

intercalations of black or green argillite or volcanic sandstone; fragments of argillite also form a normal component of the pyroclastic flow rocks. Overlying the dacite tuff breccias are a sequence of andesitic tuffaceous sediments, andesitic tuffs and cherty andesitic sedimentary rocks. The overlying black argillite and siltstone are relatively featureless, poorly bedded, but commonly displays cleavage. Intercalations of greywacke may show graded bedding, shale sharpstones and minor slump structures. Although gross stratigraphic units can be defined over much of the area, numerous lateral lithologic variations, the scarcity of marker units in the mine area, and complex deformation hampers detailed stratigraphic and structural interpretation.

Intruding this package are two major dyke sequences and a group of small mafic dykes. The early dyke intrusions are composed of dark grey-green andesites that commonly have a slightly mottled texture that reflects a fragmental nature; they may also contain abundant quartz and chlorite amygdules. They are clearly almost contemporaneous with the pyroclastic flow rocks and may be highly deformed and mineralized. The second group are massive grey-green porphyritic dacites, which show no deformation or slight deformation on their margins. Their emplacement postdates major mineralization but they have a close spatial and structural relationship to orebodies. Late dykes are common but

volumetrically insignificant and include lamprophyre, basalt and andesite.

Sulphide and genetically related deposits of anhydrite, quartz, silicified rock, cherty andesitic sedimentary rocks, bedded chert, and minor barite formed from volcanogenic hydrothermal solutions after formation of the dacite tuff breccia and during deposition of the overlying andesitic sedimentary and tuffaceous rocks. Sulphides occur as massive and stringer deposits and as disseminations and bedding plane concentrations. Massive deposits are mainly along and slightly above the upper contact of the dacite tuff breccia and commonly in or near cherty andesitic rocks. Stringer deposits are mainly in silicified dacite tuff breccia below the massive sulphide deposits. The ratio of stringer (80 per cent of ore) to massive deposits is much greater at Britannia than in most volcanogenic sulphide deposits. Original deposits and alteration halos are modified by shear deformation and segmented by faults. The massive sulphide-type orebodies mined were: Jane, Fairview Zinc (1.5 per cent of total ore mined); No. 8 (top), Beta, 040, Bluff (4.5 per cent of total ore mined); and No. 8 (bottom), No. 10, Empress, Victoria, West Victoria (15 per cent of total ore mined). Stringer-type orebodies mined were the Bluff, East Bluff, Jane, No. 4 (Bluff), No. 5, No. 10 and Fairview Veins (79 per cent of total ore mined). Other zones within and near the

mine area include the Daisy, Homestake, Robinson, Furry Creek, Fairwest and 074.

The sulphide orebodies of Britannia are highly heterogeneous mixtures of sulphides, remnant altered host rocks, and discrete veins. The main mineralogy of orebodies is simple and fairly constant. Pyrite is by far the most abundant mineral, with less chalcopyrite and sphalerite and minor erratically distributed galena, tennantite, tetrahedrite and pyrrhotite. The main nonmetallic minerals include quartz and muscovite (chlorite), anhydrite and siderite. The main massive orebodies, the Bluff, East Bluff, No. 5, No. 8 and 040 all show a marked zonal structure in which they have one or more high-grade chalcopyrite cores enveloped successively by a lower-grade zone and overlapping pyrite and siliceous zones. Zinc-rich ore tends to occur in the upper central parts of massive bodies and as almost sheet-like masses, like the Fairview Zinc vein. In section, the main orebodies have a crude lens-like shape oriented within the schistosity and are commonly connected to a steeply plunging root which may or may not be of ore grade. The other orebodies such as the Fairview Veins are stringer lodes and veins composed of thin sheet-like masses of chalcopyrite and pyrite with some quartz that appear generally parallel to the schistosity but actually cut across schistosity in plan at a small angle. Trace realgar,

orpiment, scheelite, fluorite and pyrolusite occur in post-dacite, northeast trending gash quartz-carbonate veins in the No. 10 orebody.

The ore contains thin layers of sphalerite, pyrite and barite parallel to the bedding planes (So). Galena forms irregular intergrowths in sphalerite and is abundant in a few thin layers in zinc and zinc-copper ore. Gold is abundant in scattered narrow veins in the Homestake showing, in high-grade quartz veinlets in the No. 8 orebody and throughout the No. 5 and East Bluff orebodies. Massive ore in the No. 10 mine contains pyrrhotite and argentite inclusions within the chalcopyrite-rich massive orebody. Many of the orebodies contain several types of sulphide concentrations; the No. 8 massive orebodies grade from zinc-copper to copper. The No. 8 and No. 8A ore zones contain more zinc than the No. 8B. In the Bluff deposit, sphalerite is abundant only above the 1800 level; locally in this region siliceous copper-zinc stringer ore grades into massive zinc-copper ore toward the structural footwall (stratigraphic top).

A broad zone of pervasively silicified rock surrounds all stringer orebodies in the dacite tuff breccia except the Fairview veins. Quartz and quartz-pyrite veins occur throughout the silicified halos and increase in abundance and sulphide content

toward an orebody. Pyrite is abundant as beds and nodules in andesitic sedimentary rocks above the Fairview Zinc orebody and locally pyritic layers show slumping features characteristic of soft sediment deformation. Anhydrite is abundant in pyritic andesitic sedimentary rocks and less abundant in the dacite tuff breccia in a broad elongate tabular halo around ore centres. Locally anhydrite forms massive deposits in tuffaceous sedimentary rocks, flanking and above orebodies, and is also found as distinct crosscutting veins in tensional zones. Locally the anhydrite has been converted to gypsum, especially near permeable zones where the gypsum occurs as narrow replacement veinlets. Within 60 to 90 metres of surface the conversion of anhydrite to gypsum is complete. James (1929) reports the presence of native sulphur in the mine. While the native sulphur may have gypsum or anhydrite associated with it none is present in the large gypsum masses (Open File 1991-15, page 35). Barite is disseminated and/or well bedded in zinc ore and nearby zinc-rich sedimentary rocks. Cherty andesitic sedimentary rocks and tuffs, locally with abundant pyrite, occur in and near massive sulphide bodies and host most of the No. 8 ore lenses.

Structure at the Britannia mine is complex; the earliest deformation (Do) produced widespread, open, concentric, flexural-slip folds (Fo) with subhorizontal to gently plunging, west-

northwest trending axes. A major anticline was formed in the dacitic pyroclastic rocks and a major syncline was formed in argillite to the north. Further flexural-slip deformation was localized along the Britannia anticline, which became overturned to the north. Under continued stress, deformation consisting of several episodes of inhomogeneous strain produced the Britannia and other shear zones. Rocks were crystallized to S-tectonites with phase assemblages the same as those of lower greenschist facies regional metamorphism. East of the Jane basin, the axis of the Britannia shear zone follows the axis of the Britannia anticline; from the Jane basin to the west, the shear zone cuts across the south limb of the Britannia anticline. On the surface, the shear zone narrows to a single fault west of the Jane basin, whereas at depth and to the east it widens.

The first episode of shear deformation (D1) was the most intense. Parallel orientation of recrystallized chlorite and sericite plates and flattened lithic fragments define a foliation (S1). Numerous isoclinal folds (F1) were formed with S1 as an axial plane cleavage. In the second episode of shear deformation (D2), some sericite which had formed parallel to S1 during D1 was recrystallized to define S2 into steeply dipping west plunging mesoscopic and microscopic folds (F2). A critical factor regarding the origin of the Britannia sulphide deposits is whether they are

pre- or post- D1 (and D2). Recent observations support the hypothesis that sulphide and related deposits at Britannia were deformed during D1 (see Economic Geology, Payne, et. al. 1980, for extensive discussion). The existence of stratabound ore lenses within a felsic volcanic sequence, including pyroclastic breccias, suggests that the Britannia area was a structural locus for all initial and subsequent geological processes. Volcanism, hydrothermal activity, shear deformation, faulting, and metamorphism were all dynamic forces centred along the axis presently known as the Britannia shear zone.

Rocks were altered by volcanogenic hydrothermal solutions during sulphide deposition and by metasomatic hydrothermal solutions during shear deformation. Near orebodies, alteration during deformation was superimposed on ore-stage alteration such that the two are indistinguishable. Alteration is more pronounced in andesitic than in dacitic rocks. Andesitic rocks were altered to an assemblage of quartz-chlorite-sericite (epidote-albite-potassium feldspar-calcite). Some strongly altered andesitic rocks are distinguished from strongly altered dacitic rocks by the andesite's much higher TiO_2 content. Studies of rocks near several of the orebodies show that much of the variation in chemical composition in all rock types is produced by ore-stage introduction of quartz, sulphides and sulphates.

A major compressional event (ending with D2) was followed by a period of relaxation of stress during which dacitic magma was intruded into dilated zones within the shear zone and surrounding rocks. In the shear zone, dacite formed dykes subparallel to S1 mainly in or near the dacite tuff breccia. Near the axis of the Britannia anticline, dykes coalesce upward and to the west and appear to cap some of the orebodies. Thin continuous andesite dykes are subparallel to S1 and cut the dacite dykes. Outside the shear zones, sills, dykes and irregular bodies of several varieties of dacite cut the Gambier Group rocks. The evidence suggests that most of the dykes at Britannia were intruded in the late stages of D2 deformation.

A third metamorphic foliation (S3) was formed locally, possibly following the dacite intrusion. It is parallel to northeast trending gash fractures in and near the dacites and to a set of northeast trending faults. The faults cut the dacite dykes and late andesite dykes and commonly contain vuggy quartz-carbonate veins. They have siderite-kaolinite alteration halos that are most intensely developed in rocks with abundant chlorite. A fourth metamorphic foliation (S4) is a widespread strain-slip cleavage and may have formed from a release of compression perpendicular to the shear zone.

A major set of post-dacite dyke faults cuts the Britannia shear zone subparallel to its margins and to S1. The faults converge upward and to the west to form one major fault. To the east, successive faults branch off a major footwall zone and cut diagonally across the shear zone subparallel to S1. These faults are characterized by a few centimetres to metres of gouge and/or strongly sheared rock. Many are braided and coalesce. In the major fault blocks, minor faults of a similar nature are abundant. Some show more than one age of movement. All the orebodies are cut by the minor faults and many are bounded by, or are near, one or more major faults.

Because many orebodies have contacts at or near major east striking faults and because most appear to be parts of a typical volcanogenic sulphide deposit, the present orebodies may represent faulted segments of a few original major sulphide deposits. A predeformation reconstruction suggests that the orebodies are segments of two original massive sulphide deposits; this requires a near vertical displacement along one fault zone followed by sub-horizontal offset with a cumulative right-lateral displacement of a couple of thousand of metres (Economic Geology, Payne et. al., 1980).

In summary, the Britannia ore deposits were formed from hydrothermal solutions genetically related to dacitic volcanism. Massive zinc, zinc-copper and copper deposits were formed near the contact of dacite tuff breccia and overlying fine andesitic tuff and sedimentary rocks. Siliceous stringer zones were formed in the dacitic tuff breccia and grade upward into massive deposits. Massive to disseminated bodies of anhydrite, pyrite, and minor barite were formed near the orebodies from exhalite solutions. Cherty andesitic sedimentary rocks are common near the orebodies. A northeast trending compressive stress couple produced the following events: a) Broad concentric folds, under continued stress, became tighter and slightly overturned at Britannia. The early part of deformation overlapped the late stages of dacitic volcanism and hydrothermal activity, and produced a series of subparallel fractures which acted as channelways for hydrothermal solutions. b) With continuing stress, several episodes of inhomogeneous strain produced the schistose rocks which define the Britannia shear zone. Rocks were recrystallized into S-tectonites and sulphide deposits were deformed in part by fracture and in part by plastic flow, and were segmented into a series of en echelon stringers parallel to S1. Sulphides and quartz in the orebodies show typical deformation textures similar to those of the enclosing rock. c) Ore-stage hydrothermal solutions and deformation stage solutions caused chemical alteration. Andesitic

rocks were effected more than dacitic rocks and show increases in Al_2O_3 , K_2O , SiO_2 and H_2O and decreases in CaO , FeO and MnO . TiO_2 remains relatively constant and its content can be used to distinguish some strongly altered andesitic rocks from similarly altered dacitic rocks. d) Orebodies were deformed during several periods of faulting. Following an early period of right-lateral movement, dacite dyke swarms were intruded into the shear zone generally parallel to S1 and concentrated in the dacitic tuff breccia. Dykes were cut by northeast trending quartz-carbonate gash fractures, which near orebodies contain sulphides, mainly chalcopyrite and pyrrhotite, remobilized from the orebodies. e) A major set of late east faults displaces the rock and orebodies with a cumulative right-lateral horizontal component of motion to a maximum of 2438 metres (Economic Geology, Payne, J.G. et. al., 1980).

Measured and drill indicated reserves in the No. 10 mine at the time of closure were 1,424,147 tonnes grading 1.9 per cent copper (Property File -- Northcote, K.). Past work consisted of extensive underground and surface development. Between 1905 and 1977, the Britannia orebodies yielded approximately 52.7 million tonnes of ore grading 1.1 per cent copper, 0.65 per cent zinc, 6.8 grams per tonne silver and 0.6 grams per tonne gold. The mine site

became the B.C. Museum of Mining, a National Historic Site in 1975.

The **Northair Deposit** is located in a Lower Cretaceous roof pendant of Gambier Group volcanic and sedimentary rocks within the southern Coast Plutonic Complex. This particular pendant, known as the Callaghan Creek pendant, is comprised of variably metamorphosed northwest trending volcanic and volcanically-derived sedimentary rocks, commonly characterized by a strong northwest foliation. The pendant rocks exhibit regional lower greenschist facies metamorphism, except near their contact with intrusive bodies, where they have locally undergone contact metamorphism.

The plutonic rocks in the area have a compositional range which varies from quartz monzonite to diorite. The plutonic rocks vary in age from Early Tertiary to Late Jurassic. Pendant contacts with adjacent plutonic rocks are often sharp and commonly marked by narrow shear zones which are parallel to the foliation within the pendant rocks.

Previous mapping in the Northair mine area has divided the geology of the 5000-metre thick Gambier Group into two major units. Unit 1 is a lower, volcanic-derived, sediment-rich unit characterized by well-sorted wacke with low fragment (clast)

variation and minor volcanic tuffs, indicating a relatively long depositional history. Sedimentary features such as graded bedding and crossbedding are present with indicated tops to the northeast. Thin magnetite beds are locally present in wacke sediments. The stratigraphy appears to have a north to northwest strike and a steep dip to the northeast.

Unit 2 is comprised of a volcanic tuff of predominantly andesitic composition which stratigraphically overlies unit 1. Most of the southern contact between these two units is a fault which locally is occupied by a Tertiary felsic dyke. The upper 2500 metres of unit 2 is characterized by a high variability of clast size (ash tuff to block breccia) representing a rapid depositional environment. Depositional cycles are evident by the northeastward and southward fining of these fragmentals. Locally emergent conditions are indicated by features such as hematitic clasts which are well-rounded and similar in size. This is found particularly in the upper portion of the stratigraphy (northwest part of the property).

A proximal environment is indicated for the lower 1000 metres of unit 2, which is characterized by the absence of sediments, almost chaotic and locally clast-supported angular block and ash tuffs, volcanic breccias and lapilli tuffs which represent a

brief, rapid depositional history. The significance of the lower unit lies in the fact that it hosts more of the ore.

Recent workers have interpreted the Gambier Group rocks on the property as a homoclinal succession (Assessment Report 18402). No minor fold structures have been observed. The bedding varies in strike from 160 to 200 degrees and dips from 45 to 89 degrees east. A pervasive cleavage is moderately well-developed and is common in the volcanic rocks; it has a strike of 160 to 180 degrees and is steeply inclined. Rock analyses show that the volcanics are calc-alkaline basalt to dacite in composition, with the majority of the samples falling into the andesite to dacite fields (Assessment Report 18402). Host rocks to the ore deposits at the Northair mine are andesitic pyroclastic breccia and lapilli tuffs. The ore deposits are comprised of 3 or 4 steeply dipping, fault-dismembered tabular zones, 1 to 7 metres wide and approximately 1200 metres long. They dip steeply southwest and are known to extend downdip at least 300 metres. The four mineralized segments are separated by north trending faults and are named from south to north as: Manifold, Warman, C and Discovery.

The mineralized segments are generally small bodies. The sulphides comprise pyrite, galena, sphalerite and minor chalcopyrite disseminations, veins and locally discontinuous,

banded segregations in quartz-calcite gangue. Anastomosing veins of pyrite, galena and sphalerite are common; often they are irregular sulphide pods and lenses, separated by barren, brecciated country rock (horses). Locally, spectacular ribbon-banded, quartz-chlorite-pyrite veins (with minor lead-zinc sulphides) are present in the ore zone. The vein zone which comprises most of the ore, as a whole has a steep southwest dip which is broadly discordant to the perceived northeast dip of the volcanic stratigraphy. A general pattern of sulphide mineralogy indicates silver-rich, base metal-poor mineralization in the Manifold zone, progressing to more base metals and less silver toward the northwest (through Warman, C and Discovery zones). The width of the mineralization increases from the south to the northwest. Local banded, massive sphalerite and galena were reported at the Discovery zone. Other minerals reported at the mine are tetrahedrite, argentite, bornite, pyrargyrite and electrum with trace amounts of gold and stromeyerite (Geology in British Columbia 1977-1981, page 100).

At the northwest end of the "Northair horizon" (C and Discovery zones), where highest base metal values are indicated, the tested extent of mineralization is essentially less than 150 metres below surface. This locality was considered to have the best chance for massive sulphides discovery because of reported

local occurrences of banded sulphides and shallow testing by previous exploration (Assessment Report 18402).

A consistent black, biotite/chlorite hydrothermal alteration zone is closely associated with the mineralization. This alteration forms an envelope to the sulphide vein zone, and is in some cases asymmetrical; more often it appears to be broadest in the structural hanging wall. The biotite content increases toward the sulphide vein system; it is a pervasive, fine-grained overprint of dark green chlorite. A gradation exists from a dark green, pervasive chlorite-altered tuff to a black, biotite-dominant tuff, most strongly altered nearest the mineralization. The biotite forms 6 to 7-millimetre clumps or aggregates in the altered host rock very close to, and within the mineralized vein system. Pervasive sericite alteration is also evident, but it appears to be an earlier event, and much more extensive; it is not directly related to the mineralization. Near the sulphide vein system within the alteration is a quartz-calcite stockwork which contains weak metal sulphides.

A long standing controversy has existed regarding the origin of the Northair mineralization. Two views are that the sulphides represent (1) volcanogenic massive sulphide mineralization or (2) that it is vein-type mineralization, related either to a

synvolcanic hydrothermal system, or to nearby intrusions of the Coast Plutonic Complex; the latter genesis is proposed (Assessment Report 18402).

Production at the Northair mine began in 1974 and was suspended in mid-July, 1982 due mainly to low grades and low gold prices. Indicated reserves are 59,071 tonne grading 26.73 grams per tonne silver, 9.08 per tonne gold and 2 per cent combined lead-zinc (Canadian Mines 1986-87, page 285).

The **Hummingbird**-Romana Copper showing is located on the north side of Goat Island on Powell Lake.

The showing was extensively worked in the late 1920s including numerous opencuts, a gloryhole and 2 tunnels exceeding a total of 183 metres. Romana Copper Mines Ltd. acquired Hummingbird and nine other claims in 1928. The Hummingbird claim was Crown granted in 1929. A tramway was constructed in 1928. Tunnels were driven in 1929 and 1930. The property lay dormant until 1983 when explored by Corinth Resources. In 1988, Ashworth Explorations Ltd. conducted a geochemical exploration program on the Humming Bird (Lot 4815a) Reverted Crown grant and Clover claims covering the property. The property was owned by J. Fleishman.

The area of interest consists of a roof pendant which forms a 100-metre wide belt of highly altered volcanic and sedimentary rocks unconformably overlying diorite, quartz diorite and granodiorite of the Cretaceous Coast Plutonic Complex. The apparent strike of the belt, thought to be part of the Lower Cretaceous Gambier Group, is about 220 degrees.

Within this roof pendant is a contact metamorphosed zone containing garnetite, epidote and mineralization. The mineralization, manifested by rusty zones and malachite stain, consists of pods, streaks, veins and lenses of massive sulphides composed of varying proportions of pyrite and chalcopyrite. Most samples were moderately magnetic, and magnetite was identified in some specimens.

The best silver values occur in the opencut from which previous ore shipments were made. In 1983, a chip sample over unknown length assayed 17.40 per cent copper and 320.17 grams per tonne silver (Assessment Report 11884). Eight rock chip samples were taken during property exploration in 1988. Sample CL88-R2 yielded 3.08 per cent copper, 52.80 grams per tonne silver and 0.27 gram per tonne gold (Assessment Report 18531). The sample was a 100-centimetre chip sample across malachite stained, heavily altered metavolcanics striking 160 degrees and dipping vertical.

One hundred and forty tonnes of ore are quoted as being mined and shipped several years before 1928 assaying 8 to 11 per cent copper, 240 to 685 grams per tonne silver and minor gold (Minister of Mines Annual Report 1928).

Mineralization in the **Mount Diadem** area became known in 1928, when several massive sulphide showings containing pyrite, pyrrhotite, chalcopyrite and sphalerite were discovered near the headwaters of No Man's Creek. Both Britain River Mining Co. Ltd. and Mount Diadem Mines Ltd. staked claims west and north of Mount Diadem. Numerous trenches were excavated where sulphide showings occurred in altered limestone and other sedimentary rocks. Some adits were driven and work continued sporadically over the years. The original claims lapsed and were restaked in 1947 by Nickel Mining Company of Canada Ltd. The new claims were optioned to Bralorne Mines Ltd. in 1949. Considerable work has been carried out since 1949 by various operators. Geological mapping, limited diamond drilling and sampling of old adits and trenches were performed by Sphere Development Corp. in 1967. In 1970, Tiger Silver Mines Ltd. performed geophysical magnetic and geochemical soil surveys. Britain River Syndicate performed geological, geophysical and geochemical surveys in 1971. Some new anomalies were discovered. Minor rock sampling was conducted by Fury

Explorations in 1980. The claims were transferred to Fury Explorations Ltd. in the early 1980s. In 1983, Anaconda Ltd. optioned these claims and conducted a drilling program, consisting of nine holes and 899 metres. In the late 1980s, Covenant Resources staked the Diadem claims, surrounding the claim owned by Fury Exploration and the Fox claim owned by R. Schmidt.

Immediately above the head of No Man's Creek on the northern slopes of Mount Diadem an old adit is located at an elevation of 900 metres. The adit lies within the Cretaceous Coast Plutonic Complex near its western boundary with the Insular Belt. The complex consists mainly of diorites, granodiorites, gneisses and migmatites enclosing a northwest trending belt (pendant) of Lower Cretaceous Gambier volcanic and sedimentary rocks. Only in the eastern and possibly basal part of the belt are mafic flows and interbedded tuff evident. These rocks have been metamorphosed to greenschist and less commonly amphibolite grade. Structural deformation has been intense with the early development of tight, moderate to steep, north plunging folds characterized by an axial planar cleavage. This has been overprinted with later, open style folds. Two shear orientations predominate, both of which appear to locally control massive sulphide mineralization. One is subparallel to regional banding and parallel to the penetrative

foliation. The other set strikes 060 to 100 degrees and is steeply dipping.

Seven rock units have been defined locally. These are: (1) tuffaceous sandstone, siltstone and argillite; andesitic flows, lapilli tuff and chloritic schist and massive diorite, (2) green-grey, chlorite-rich tuff, tuffaceous sandstone; felsic lapilli and vesicular flows and breccias and massive diorite, (3) rusty to black weathering, thinly bedded argillite, (4) well banded, grey-green interbedded argillite, siltstone, sandstone, black chert and lapilli tuffs, (5) siliceous argillite, tuffaceous siltstone, chert and lapilli tuff, (6) andesitic breccia and (7) feldspar-rich diorite, quartz diorite and granite. The adit is collared at the contact of the volcanic rocks with the intrusive rocks. The adit penetrates the silicified, recrystallized volcanics for 12 metres, at which distance a 0.61-metre shear is intersected. Pods consisting of galena, sphalerite, pyrite and small amounts of chalcopyrite are exposed in the shear.

A 0.25-metre wide sample of the shear southeast of the adit assayed 0.017 per cent copper, greater than 1 per cent lead, greater than 1 per cent zinc, greater than 200 grams per tonne silver and 0.18 gram per tonne gold (Assessment Report 11641). A grab sample from the adit assayed 4.9 grams per tonne gold, 264

grams per tonne silver, 8.89 per cent lead, 8.62 per cent zinc and 0.02 per cent copper (Assessment Report 11641).

Diamond drilling completed under option to Anaconda has tested up to 175 metres along strike, the contact between sheared argillite - chloritized volcanics. Three zones were believed intersected; the North, Central and South. The best drilling results were obtained from the Central zone. Diamond-drill hole 84-3 intersected 0.79 per cent copper, 2.74 per cent lead, 1.61 per cent zinc and 148.80 grams per tonne silver over 12.0 metres (Assessment Report 18207). The Central zone was also intersected by drillholes 84-1, 84-5, 84-6, and 84-8. The South zone was intersected in drillhole 84-9, approximately 60 metres below the surface. A 7.7-metre section yielded 0.1 per cent copper, 1.48 per cent lead, 1.53 per cent zinc and 44.91 grams per tonne silver (Assessment Report 18207). Mineralization in all intersections is hosted in intensely deformed argillite.

REGIONAL GEOCHEMISTRY

A regional stream sediment survey published by the Geological Survey of Canada in 1988 indicates geochemical anomalies in streams that drain the Old Ironsides 2 claim. Streams are moderately anomalous for molybdenum and weakly anomalous for copper, zinc and lead.


PROPERTY GEOLOGY

Much of the Old Ironsides 2 Claim was mapped in detail during 1999 (Figure 5). A soil geochemistry grid (Figure 7) was used for control over most of the ground mapped, while elevation was used for control in outlying areas. With the exception of a small portion in its southwestern corner, the property covers Gambier group metamorphic rocks that strike northwest and dip steeply to vertically.

The southernmost rock type mapped on the property is medium to coarse grained, unfoliated granodiorite. It is a part of the **Coast Mountain intrusive** suite, and has not been subjected to regional metamorphism. Its occurrence on the property is limited to a small area south of the creek in the property's southwestern corner. The remainder of the rocks mapped belong to Gambier group.


The next rock type to the north is fine grained black **amphibolite**. It is massive to weakly foliated and sometimes contains a small amount of fine grained and evenly disseminated pyrite. This rock weathers grey to light grey.

To the north of the amphibolite lies a thick package of interlayered **schist** and **felsite**. The contact between this package



- 7 Coast Mountain intrusives
- 6 Mafic agglomerate
- 5 Mixed unit
- 4 Granodiorite gneiss
- 3 Felsite
- 2 Schist
- 1 Amphibolite

- P197001 □ Rock float sample with sample number
- P197002 ■ Rock outcrop sample with sample number
- ↖ 090 Foliation attitude
- ↕ 090 Vertical foliation attitude



- Lake, stream
- ⋯ Outcrop boundary
- - - Geologic contact
- ▲ Fly camp location

FIGURE 5

OLD IRONSIDES 2 CLAIM

Property Geology

DRAWN BY: AB	PRODUCED AT: 1:10000
DATE: OCT 12, 1999	FILE: 012\10GEOI.DWG

and the amphibolite appears to be conformable. The schist is very dark grey and moderately to strongly foliated. It is composed of 50% anhedral black grains, 1mm to 2mm in size, which have not been positively identified, in a very fine grained black matrix. This rock often contains up to 2% very fine grained pyrite in disseminations and thin bands, and usually weathers rusty. Near L20700N, 19800E, interbeds to 50mm in thickness of medium grained greywacke were noted.

The felsite is fine to medium grained and white. It is mostly unfoliated to weakly foliated, except in rare localities where it is strongly foliated and schistose. The felsite commonly contains up to 5% very fine to fine grained pyrite as irregular disseminations, thin bands and fracture coatings. Near L20200N, 20100E, a 3m thick section contains mica flakes and has an intrusive texture that is distinct from most occurrences of this rock. Near L20800N, 20600E, the felsite hosts a 0.4m thick quartz vein which contains sporadically disseminated medium grained pyrite. A specimen of sulphide-bearing vein was analysed and returned an anomalous gold value of 1700ppb.

A medium grained, **granodiorite gneiss** lies within the schist-felsite assemblage near the northern end of the control grid. It is weakly to moderately foliated and inequigranular with brownish

quartz grains and biotite flakes, both up to 3mm in size, in a light-coloured fine grained groundmass. This rock may be a partially recrystallized grit or immature sandstone.

The felsite grades into a rapidly changing **mixed unit** composed of felsic and mafic volcanics, some of which appear intrusive, and minor interbeds of schist and grit. Near the ridgetop north of rock sample P197016 is an exposure of boulder conglomerate.

At the top of the ridge is a grey weathering **mafic agglomerate** which is dark grey to black, rarely with a very dark greenish tinge. It consists of 35% very fine grained black groundmass and 65% fine grained dark grey to black angular to subangular clasts. The clasts range in size from 2mm to 70mm. To the north of the claim boundary, on ground covered by the Old Ironsides 3 claim, the agglomerate is host to an occurrence of pyrite and chalcopyrite.

On the ridgetop, the agglomerate is intruded by granodiorite dykes which have not been affected by regional metamorphism and are probably of the same age as the Coast Plutonics diorite to the north of the property.

PROPERTY MINERALIZATION

Fine bands of pyrite are common within the felsics at numerous localities. This style of mineralization resembles that observed at many known VMS occurrences. Samples of the pyrite-bearing felsics were submitted to Chemex Labs in North Vancouver, BC, where they were crushed, split and pulverized to -150 mesh, digested in nitric aqua regia and analysed for 32 elements using an induced coupled plasma (ICP) technique. Certificates of Analysis appear in Appendix II. Sample P197015, a sulphide-bearing specimen of felsite, returned values of 1305ppm Cu and 4ppm silver.

Near L20800N, 20600E, the felsite hosts a 0.4m thick quartz vein which contains sporadically disseminated medium grained pyrite. A specimen (P197019) of sulphide-bearing vein was analysed and returned a value of 69.4 ppm silver. A 30 gram sample of the rock was further analysed for gold by fire assay with atomic absorption finish and returned a value of 1700ppb gold. VMS deposits account for a considerable proportion of gold and silver production in Canada (Franklin, 1996), and the gold produced at the Britannia deposit was recovered from quartz veins (Payne et al, 1980).

At the top of the ridge, on ground now protected by the Old Ironsides 3 claim, an occurrence of pyrite and chalcopyrite was found during 1998. The sulphide occurs as coarse grained blobs in the matrix of a mafic agglomerate and comprises about 20% of the rock. Two mineralized specimens were analyzed and returned values of 1055ppm and 1140ppm copper.


PROPERTY GEOCHEMISTRY

During 1999, 123 soil samples were collected from a grid with a sample density of 100 by 100m. The grid was established using flagging tape and aluminum tags inscribed with grid coordinates. The tags were fastened either to foliage with wire or to lath pickets with staples. To mark the grid location permanently, a 50cm long, 12mm diameter steel pin was inserted into the ground to a depth of 40cm at each end of baseline 20000E. All soil samples were submitted to Chemex Labs in North Vancouver, BC, where they were dried, screened to -150 mesh, split, digested in nitric aqua regia and analysed for 32 elements using an induced coupled plasma (ICP) technique. Certificates of analysis appear in Appendix II.

An elementary statistical analysis was performed on a population of soil samples collected during a regional exploration program conducted in the Powell River region during 1998 and 1999. A total of 522 soil samples is included in the analysis, all of which were collected over roof pendants of metamorphic Gambier group rocks. The resulting data were used to establish thresholds for geochemical anomalies. A summary of the data and histogram plots for elements of primary interest is shown in Figure 6, while the threshold values are summarized in the following table.


Figure 6: Regional Soil Geochemistry Histograms
 Horizontal Axes: geochemical value range; Vertical Axes: number of values in range
 Number of samples n=522
 NB: outliers trimmed from histograms but included in calculations

Copper




mean	52	peak	17.5
median	24	95th percentile	187
standard deviation	99	90th percentile	149

Arsenic




mean	19	peak	750
median	8	95th percentile	66
standard deviation	46	90th percentile	46

Zinc




mean	57	peak	9650
median	42	95th percentile	118
standard deviation	162	90th percentile	102

Silver




mean	0.32	peak	48.8
median	0.00	95th percentile	1.00
standard deviation	2.15	90th percentile	0.60

Barium (partial digestion)



mean	153	peak	1510
median	70	95th percentile	650
standard deviation	248	90th percentile	429

Cobalt



mean	9.2	peak	63
median	7.0	95th percentile	25
standard deviation	9.4	90th percentile	19



ANOMALOUS THRESHOLDS FOR 522 SOIL SAMPLES

Element	Anomalous threshold (ppm)				
	Background	Weak	Moderate	Strong	Peak
Cu	25	45	90	180	1775
Zn	15	30	60	120	3650
Ag	<0.2	0.3	0.6	1.0	48.8
Co	4	6	12	25	83
Ba*	75	150	300	600	1510
As	7	15	30	60	750
Pb	2	5	10	20	584

*partial digestion

Overburden is considered largely residual or colluvial, and usually of thickness of less than two metres. It was observed that soil development north of L20600N is extremely poor, with no differentiable horizons. Material commonly sampled was a brown-grey C horizon. The poor soil development suggests that the soil geochemistry may not reflect an accurate signature of bedrock mineralization. South of L20600N, red-brown or orange-brown B horizons were most commonly sampled.

The grid covers the area around two 1998 reconnaissance soil lines (Figure 7) which had returned anomalous values for copper and barium. It covers a zone in which the volcanics change in composition from predominantly mafic to predominantly felsic. A considerable portion of the area of geochemical interest could not be sampled because of cliffs.



○ Soil sample location with sample number

○ Lake, stream

FIGURE 7

OLD IRONSIDES 2 CLAIM

SAMPLE LOCATIONS



DRAWN BY: AB	PRODUCED AT: 1:10000
DATE: OCT 12, 1999	FILE: 012\10GEOL.DWG

The grid soil samples confirm and partially define the 1998 copper and barium anomalies on the lower portion of the hillside (Figures 8,9). Peak values are 181ppm copper and 400ppm barium. Anomalous values were also returned over an extensive area for zinc (Figure 10). Barium values are likely understated because of incomplete solution of barium sulphate by the aqua regia. Barite is an accessory mineral at the Britannia deposit (Payne et al, 1980) and the Red Dog deposit in Alaska (Koehler et al, 1991), as well as numerous other VMS deposits (Hoffman, 1986).

North of L20600N, where the slope of the hillside decreases, there is a sharp drop in geochemical background for most metals, and soil horizons are more poorly developed. There are several subtle anomalies in this portion of the grid.

The twelve samples in the northeast corner of the grid returned subtly but distinctly anomalous values for any of copper, zinc, barium, silver (Figure 11), or a combination of these elements. This is the area where pyritic felsic volcanics host a quartz vein which contains anomalous amounts of silver and gold.

Another significant anomaly occurs at L21000N, 20000E, where silver, copper and zinc values are anomalous. Arsenic values (Figure 12) are extremely anomalous 200m to the south and also to





- 20 Soil sample location with copper value in ppm
- < Soil sample location with copper value below detection limit
- Lake, stream

FIGURE B

OLD IRONSIDES 2 CLAIM

COPPER SOIL GEOCHEMISTRY

DRAWN BY: AB	PRODUCED AT: 1:10000
DATE: OCT 12, 1999	FILE: 012\10GEOI.DWG



- 120 Soil sample location with barium value in ppm
NE: partial digestion only
- < Soil sample location with barium value below detection limit


 Lake, stream



FIGURE 9

OLD IRONSIDES 2 CLAIM

BARIUM
SOIL GEOCHEMISTRY

NB: Partial digestion only

DRAWN BY: AB	PRODUCED AT: 1:10000
DATE: OCT 12, 1999	FILE: 012\10GEOI.DWG



- 100 Soil sample location with zinc value in ppm
- < Soil sample location with zinc value below detection limit


Lake, stream

FIGURE 10

OLD IRONSIDES 2 CLAIM

ZINC
SOIL GEOCHEMISTRY

DRAWN BY: AB	PRODUCED AT: 1:10000
DATE: OCT 12, 1999	FILE: 01210GEO1.DWG



- 0.4 Soil sample location with silver value in ppm
- < Soil sample location with silver value below detection limit

Lake, stream


FIGURE 11

OLD IRONSIDES 2 CLAIM

SILVER
SOIL GEOCHEMISTRY

DRAWN BY: AB
DATE: OCT 12, 1999

PRODUCED AT: 1:10000
FILE: 012\10GEO1.DWG




○20 Soil sample location with arsenic value in ppm
 < Soil sample location with arsenic value below detection limit
 Lake, stream

FIGURE 12

OLD IRONSIDES 2 CLAIM

ARSENIC
SOIL GEOCHEMISTRY

DRAWN BY: AB	PRODUCED AT: 1:10000
DATE: OCT 12, 1999	FILE: 012\10GEO1.DWG

the west of this site, with a peak value of 274ppm.

CONCLUSIONS AND RECOMMENDATIONS

The Old Ironsides 2 mineral claim was staked in June, 1999 to protect a VMS base and precious metals target lying within a metamorphic roof pendant of Gambier Group rocks. During 1999, prospecting, geological mapping and soil sampling were carried out. 123 soil samples and 21 rock samples were collected. 1999 exploration resulted in the discovery of a gold-silver bearing quartz vein, partial definition of a copper-barium soil geochemical anomaly and the discovery of two new soil geochemical anomalies. Due to poor access, helicopter support is recommended for future work.

It is recommended that the soil grid be extended to the north to define two anomalies at the northern end of the current grid. The soil samples in the northeastern quadrant of the current grid should be reanalyzed for gold, as a gold-silver bearing vein was found in that area. Geologic mapping and prospecting should also be extended northward.

Respectfully submitted,

Arnd Burgert

REFERENCES

- Franklin, J. M., 1996: Volcanogenic-associated massive sulphide base metals; in *Geology of Canadian Mineral Deposit Types*, (ed.) O. R. Eckstrand, W. D. Sinclair, and R. I. Thorpe; Geological Survey of Canada, *Geology of Canada*, no. 8, p. 158-183
- Geological Survey of British Columbia: Minfile Database
- Geological Survey of Canada, 1988: Stream sediment and water geochemical survey, northern Vancouver Island and adjacent mainland, Geological Survey Open File 2039
- Hoffman, S. J., 1986: The volcanogenic massive sulphide target, in *Exploration geochemistry: design and interpretation of soil surveys*, *Reviews in economic geology*, (ed.) James M. Robertson, Society of economic geologists, pp. 139-146.
- Koehler, George F., and George D. Tikkanen, 1991: Red Dog, Alaska: Discovery and definition of a major zinc-lead-silver deposit, *Economic Geology*, Monograph 8, pp. 268-274.
- Payne, J. G., J. A. Bratt and B. G. Stone, 1980: Deformed mesozoic volcanogenic Cu-Zn sulphide deposits in the Britannia district, British Columbia, *Economic Geology*, Volume 75, pp. 700-721.
- Roddick, J. A., 1976: Notes on the stratified rocks of Bute Inlet map-area, Geological Survey of Canada Open File 480.
- Roddick, J. A., W. W. Hutchison and G. J. Woodsworth, 1976: *Geology of Bute Inlet map area*, Geological Survey of Canada Open File 480.
- Roddick, J. A., and G. J. Woodsworth, 1979: *Geology of Vancouver, west half, and mainland part of Alberni*, Geological Survey of Canada Open File 611.
- Woodsworth, G. J., 1977: *Geology of Pemberton map-area*, Geological Survey of Canada Open File 482.

APPENDIX I

AUTHOR'S STATEMENT OF QUALIFICATIONS

AUTHOR'S STATEMENT OF QUALIFICATIONS

I, Arnd Burgert, geologist, with business and residential address in New Westminster, British Columbia, do hereby certify that:

1. I graduated from the University of British Columbia in 1995 with a B.Sc. in Geology.
2. From 1989 to present, I have been actively engaged in mineral exploration in British Columbia, the Northwest Territories and the Yukon Territory.
3. I have personally performed the work reported herein.

A. Burgert, B.Sc.

Dated this 30th day of October, 1999

APPENDIX II
CERTIFICATES OF ASSAY



Chemex Labs Ltd.

Analytical Chemists * Geochemists * Registered Assayers
 212 Brooksbank Ave., North Vancouver
 British Columbia, Canada V7J 2C1
 PHONE: 604-984-0221 FAX: 604-984-0218

To: BURGERT, ARND

P.O. BOX 1208
 PORT MCNEILL, BC
 V0N 2R0

Project: OLD IRONSIDES
 Comments: ATTN: ARND BURGERT CC: ARND BURGERT

Page Number : 1-A
 Total Pages : 4
 Certificate Date: 09-SEP-1999
 Invoice No. : 19927561
 P.O. Number :
 Account : QHB

CERTIFICATE OF ANALYSIS A9927561

SAMPLE	PREP CODE	Ag ppm	Al %	As ppm	B ppm	Ba ppm	Be ppm	Bi ppm	Ca %	Cd ppm	Co ppm	Cr ppm	Cu ppm	Fe %	Ga ppm	Hg ppm	K %	La ppm	Mg %	Mn ppm
L20000N 19400E	201 229	0.8	4.71	8	< 10	160	< 0.5	< 2	0.24	< 0.5	12	14	32	4.92	10	< 1	0.53	< 10	1.28	525
L20000N 19500E	201 229	0.2	3.22	14	< 10	180	< 0.5	< 2	0.20	< 0.5	14	27	33	4.32	10	< 1	0.58	< 10	1.14	600
L20000N 19600E	201 229	0.2	2.49	4	< 10	80	< 0.5	< 2	0.11	< 0.5	10	4	16	2.97	< 10	< 1	0.23	< 10	0.81	730
L20000N 19700E	201 229	0.2	3.37	12	< 10	200	< 0.5	< 2	0.22	< 0.5	13	6	28	4.22	10	< 1	0.43	< 10	1.27	720
L20000N 19800E	201 229	0.2	3.06	28	< 10	100	< 0.5	< 2	0.06	< 0.5	9	52	43	5.75	10	< 1	0.27	< 10	0.67	285
L20000N 19900E	201 229	0.2	2.89	22	< 10	120	< 0.5	< 2	0.06	< 0.5	10	53	37	4.08	< 10	< 1	0.20	< 10	0.53	410
L20000N 20000E	201 229	0.2	2.67	18	< 10	110	< 0.5	< 2	0.17	< 0.5	11	59	32	4.24	10	< 1	0.29	10	0.73	475
L20000N 20100E	201 229	0.4	3.24	14	< 10	180	< 0.5	< 2	0.24	< 0.5	12	57	43	4.38	10	< 1	0.61	< 10	0.93	445
L20000N 20200E	201 229	0.2	4.09	4	< 10	220	< 0.5	< 2	0.67	< 0.5	20	9	30	4.66	10	< 1	0.37	< 10	2.15	740
L20000N 20350E	201 229	0.2	3.72	8	< 10	140	< 0.5	< 2	0.06	< 0.5	7	57	34	5.44	10	< 1	0.28	< 10	1.14	350
L20100N 19400E	201 229	< 0.2	1.64	2	< 10	230	< 0.5	< 2	0.20	< 0.5	6	1	9	2.09	10	< 1	0.49	< 10	0.80	165
L20100N 19500E	201 229	0.4	3.22	14	< 10	150	< 0.5	< 2	0.25	< 0.5	20	21	31	4.37	10	< 1	0.40	< 10	1.14	1150
L20100N 19600E	201 229	0.2	6.41	8	< 10	310	< 0.5	< 2	1.09	< 0.5	33	9	12	4.95	10	< 1	0.52	< 10	2.77	735
L20100N 19700E	201 229	0.2	5.14	2	< 10	260	< 0.5	< 2	0.50	< 0.5	18	8	28	4.86	10	< 1	0.62	< 10	2.21	835
L20100N 19800E	201 229	0.2	2.87	12	< 10	200	< 0.5	< 2	0.19	< 0.5	15	45	26	4.50	10	< 1	0.36	< 10	0.93	705
L20100N 19900E	201 229	0.2	3.50	24	< 10	170	< 0.5	< 2	0.22	< 0.5	16	57	48	4.51	< 10	< 1	0.39	< 10	0.91	835
L20100N 20000E	201 229	0.2	3.24	20	< 10	90	< 0.5	< 2	0.03	< 0.5	8	65	40	4.71	10	< 1	0.26	< 10	0.74	245
L20100N 20100E	201 229	0.2	2.77	14	< 10	170	< 0.5	< 2	0.18	< 0.5	14	38	37	3.75	< 10	< 1	0.54	< 10	0.76	635
L20100N 20200E	201 229	0.2	1.51	120	< 10	30	< 0.5	< 2	0.01	< 0.5	3	16	17	3.51	< 10	< 1	0.05	10	0.12	75
L20100N 20300E	201 229	0.6	2.40	20	< 10	60	< 0.5	< 2	0.03	< 0.5	5	56	25	6.20	10	< 1	0.13	< 10	0.24	115
L20100N 20350E	201 229	0.2	3.80	20	< 10	40	< 0.5	< 2	0.07	< 0.5	7	62	54	6.25	10	1	0.13	< 10	0.54	175
L20200N 19600E	201 229	0.6	5.31	14	< 10	140	< 0.5	< 2	0.42	< 0.5	28	< 1	38	5.54	< 10	< 1	0.27	< 10	0.51	1275
L20200N 19700E	201 229	0.2	2.79	4	< 10	60	< 0.5	< 2	0.10	< 0.5	10	6	31	3.19	10	< 1	0.09	< 10	1.03	420
L20200N 19800E	201 229	0.4	5.02	10	< 10	270	< 0.5	< 2	0.19	< 0.5	18	7	48	4.29	10	< 1	0.62	< 10	1.36	585
L20200N 19900E	201 229	0.6	4.80	14	< 10	150	< 0.5	< 2	0.11	< 0.5	28	46	52	5.54	10	< 1	0.20	< 10	0.67	1000
L20200N 20000E	201 229	0.2	4.64	14	< 10	120	< 0.5	< 2	0.13	< 0.5	15	21	44	4.11	10	< 1	0.33	< 10	1.24	375
L20200N 20100E	201 229	0.2	2.72	20	< 10	170	< 0.5	< 2	0.16	< 0.5	16	48	42	3.86	< 10	< 1	0.59	< 10	0.74	805
L20200N 20200E	201 229	0.2	4.61	14	< 10	150	< 0.5	< 2	0.07	< 0.5	16	95	87	4.80	10	< 1	0.26	< 10	0.88	385
L20200N 20300E	201 229	0.4	5.50	16	< 10	130	< 0.5	< 2	0.11	< 0.5	15	84	98	5.26	10	< 1	0.39	< 10	1.04	390
L20200N 20400E	201 229	0.2	2.24	4	< 10	30	< 0.5	< 2	0.03	< 0.5	3	27	8	3.44	10	< 1	0.06	< 10	0.15	55
L20200N 20500E	201 229	< 0.2	1.02	< 2	< 10	30	< 0.5	< 2	0.03	< 0.5	2	7	3	3.48	< 10	< 1	0.03	< 10	0.12	140
L20200N 20600E	201 229	< 0.2	1.52	2	< 10	20	< 0.5	< 2	0.04	< 0.5	3	1	< 1	4.08	10	< 1	0.08	< 10	0.30	425
L20200N 20650E	201 229	0.4	5.54	< 2	< 10	60	0.5	< 2	0.04	< 0.5	9	54	46	4.26	10	< 1	0.14	< 10	0.30	160
L20300N 19700E	201 229	0.2	5.60	6	< 10	170	< 0.5	< 2	0.26	< 0.5	19	7	39	4.22	10	< 1	0.29	< 10	1.54	660
L20300N 19800E	201 229	< 0.2	4.43	12	< 10	280	< 0.5	< 2	0.29	< 0.5	23	5	47	4.28	10	< 1	0.74	< 10	1.55	660
L20300N 19900E	201 229	0.4	2.20	8	< 10	40	< 0.5	< 2	0.04	< 0.5	4	8	31	2.36	< 10	< 1	0.05	< 10	0.26	80
L20300N 20000E	201 229	0.2	2.71	22	< 10	70	< 0.5	< 2	0.04	< 0.5	5	41	24	4.51	10	< 1	0.08	< 10	0.35	190
L20300N 20100E	201 229	0.2	3.78	12	< 10	80	< 0.5	< 2	0.04	< 0.5	9	79	53	4.99	10	< 1	0.20	< 10	0.65	210
L20300N 20200E	201 229	0.2	3.07	16	< 10	140	< 0.5	< 2	0.16	0.5	20	41	49	3.62	< 10	< 1	0.48	< 10	0.70	760
L20300N 20300E	201 229	0.2	2.10	14	< 10	70	< 0.5	< 2	0.04	< 0.5	6	15	13	3.61	10	< 1	0.20	< 10	0.40	450

CERTIFICATION: _____



Chemex Labs Ltd.

Analytical Chemists * Geochemists * Registered Assayers
 212 Brooksbank Ave., North Vancouver
 British Columbia, Canada V7J 2C1
 PHONE: 604-984-0221 FAX: 604-984-0218

To: BURGERT, ARND

P.O. BOX 1208
 PORT MCNEILL, BC
 V0N 2R0

Page Number : 1-B
 Total Pages : 4
 Certificate Date: 09-SEP-1999
 Invoice No. : 19927561
 P.O. Number :
 Account : QHB

Project : OLD IRONSIDES
 Comments : ATTN: ARND BURGERT CC: ARND BURGERT

CERTIFICATE OF ANALYSIS A9927561

SAMPLE	PREP CODE	Mo ppm	Na %	Ni ppm	P ppm	Pb ppm	S %	Sb ppm	Sc ppm	Sr ppm	Ti %	Tl ppm	U ppm	V ppm	W ppm	Zn ppm
L20000N 19400E	201 229	< 1	0.07	7	670	< 2	0.06	< 2	9	14	0.22	< 10	< 10	120	< 10	74
L20000N 19500E	201 229	< 1	0.04	13	760	< 2	0.06	< 2	9	13	0.19	< 10	< 10	107	< 10	80
L20000N 19600E	201 229	< 1	0.03	3	430	< 2	0.05	< 2	5	9	0.17	< 10	< 10	93	< 10	42
L20000N 19700E	201 229	1	0.04	5	850	< 2	0.08	< 2	8	15	0.16	< 10	< 10	114	< 10	96
L20000N 19800E	201 229	< 1	0.02	23	510	6	0.05	< 2	6	14	0.20	< 10	< 10	115	< 10	74
L20000N 19900E	201 229	< 1	0.01	25	520	4	0.06	< 2	5	10	0.14	< 10	< 10	73	< 10	70
L20000N 20000E	201 229	1	0.03	30	590	10	0.05	< 2	6	12	0.13	< 10	< 10	65	< 10	88
L20000N 20100E	201 229	< 1	0.03	35	730	4	0.04	< 2	8	21	0.17	< 10	< 10	75	< 10	104
L20000N 20200E	201 229	1	0.08	8	620	< 2	0.05	< 2	14	28	0.27	< 10	< 10	165	< 10	102
L20000N 20350E	201 229	2	0.05	10	210	2	0.05	< 2	10	12	0.29	< 10	< 10	142	< 10	86
L20100N 19400E	201 229	< 1	0.05	< 1	580	2	0.06	< 2	9	13	0.15	< 10	< 10	69	< 10	50
L20100N 19500E	201 229	1	0.04	12	730	2	0.07	< 2	8	15	0.19	< 10	< 10	112	< 10	80
L20100N 19600E	201 229	1	0.38	7	580	< 2	0.03	< 2	12	107	0.30	< 10	< 10	185	< 10	128
L20100N 19700E	201 229	1	0.13	4	520	6	0.05	< 2	12	32	0.26	< 10	< 10	155	< 10	94
L20100N 19800E	201 229	1	0.03	19	710	2	0.06	< 2	7	23	0.16	< 10	< 10	106	< 10	84
L20100N 19900E	201 229	< 1	0.02	36	830	4	0.08	< 2	7	18	0.15	< 10	< 10	93	< 10	102
L20100N 20000E	201 229	< 1	0.01	29	480	4	0.06	< 2	7	9	0.18	< 10	< 10	100	< 10	78
L20100N 20100E	201 229	< 1	0.03	22	920	4	0.05	< 2	6	16	0.15	< 10	< 10	53	< 10	82
L20100N 20200E	201 229	1	0.01	8	240	8	0.01	< 2	1	3	0.09	< 10	< 10	64	< 10	30
L20100N 20300E	201 229	3	0.01	15	410	8	0.03	< 2	4	5	0.26	< 10	< 10	129	< 10	38
L20100N 20350E	201 229	< 1	0.02	24	570	8	0.05	< 2	6	10	0.21	< 10	< 10	128	< 10	78
L20200N 19600E	201 229	5	0.07	< 1	990	4	0.27	< 2	7	31	0.17	< 10	< 10	77	< 10	48
L20200N 19700E	201 229	< 1	0.04	3	410	8	0.05	< 2	6	10	0.28	< 10	< 10	132	< 10	50
L20200N 19800E	201 229	< 1	0.04	7	600	< 2	0.05	< 2	10	16	0.21	< 10	< 10	128	< 10	90
L20200N 19900E	201 229	< 1	0.02	34	1000	6	0.08	< 2	5	15	0.13	< 10	< 10	87	< 10	134
L20200N 20000E	201 229	1	0.03	12	490	< 2	0.03	< 2	8	7	0.23	< 10	< 10	118	< 10	74
L20200N 20100E	201 229	< 1	0.03	28	850	6	0.05	< 2	6	13	0.15	< 10	< 10	55	< 10	116
L20200N 20200E	201 229	< 1	0.01	55	670	6	0.04	< 2	8	12	0.22	< 10	< 10	114	< 10	126
L20200N 20300E	201 229	< 1	0.02	61	1040	6	0.06	< 2	8	7	0.19	< 10	< 10	116	< 10	134
L20200N 20400E	201 229	< 1	0.01	5	170	2	0.02	< 2	3	3	0.22	< 10	< 10	103	< 10	18
L20200N 20500E	201 229	< 1	0.01	< 1	210	2	0.02	< 2	1	3	0.24	< 10	< 10	39	< 10	16
L20200N 20600E	201 229	< 1	0.01	< 1	150	2	0.01	< 2	3	1	0.28	< 10	< 10	16	< 10	30
L20200N 20650E	201 229	< 1	0.01	17	340	8	0.04	< 2	4	5	0.15	< 10	< 10	73	< 10	52
L20300N 19700E	201 229	< 1	0.06	5	470	< 2	0.04	< 2	10	16	0.28	< 10	< 10	149	< 10	80
L20300N 19800E	201 229	< 1	0.05	6	510	< 2	0.04	< 2	11	18	0.23	< 10	< 10	134	< 10	110
L20300N 19900E	201 229	1	0.01	10	210	< 2	0.03	< 2	2	6	0.14	< 10	< 10	60	< 10	22
L20300N 20000E	201 229	1	0.01	14	430	2	0.04	< 2	4	10	0.17	< 10	< 10	89	< 10	52
L20300N 20100E	201 229	1	0.02	35	420	2	0.03	< 2	8	5	0.25	< 10	< 10	130	< 10	92
L20300N 20200E	201 229	1	0.01	36	860	4	0.05	< 2	6	12	0.12	< 10	< 10	53	< 10	176
L20300N 20300E	201 229	< 1	0.01	6	220	2	0.02	< 2	4	7	0.25	< 10	< 10	39	< 10	48

CERTIFICATION:



Chemex Labs Ltd.

Analytical Chemists * Geochemists * Registered Assayers

212 Brooksbank Ave., North Vancouver
British Columbia, Canada V7J 2C1
PHONE: 604-984-0221 FAX: 604-984-0218

To: BURGERT, ARND

P.O. BOX 1208
PORT MCNEILL, BC
V0N 2R0

Project : OLD IRONSIDES
Comments: ATTN: ARND BURGERT CC: ARND BURGERT

Page Number :2-A
Total Pages :4
Certificate Date: 09-SEP-1999
Invoice No. :19927561
P.O. Number :
Account :QHB

CERTIFICATE OF ANALYSIS A9927561

SAMPLE	PREP CODE	Ag ppm	Al %	As ppm	B ppm	Ba ppm	Be ppm	Bi ppm	Ca %	Cd ppm	Co ppm	Cr ppm	Cu ppm	Fe %	Ga ppm	Hg ppm	K %	La ppm	Mg %	Mn ppm
L20300N 20350E	201 229	0.2	3.82	10	< 10	90	< 0.5	< 2	0.05	< 0.5	9	75	60	6.85	10	< 1	0.24	< 10	0.75	290
L20400N 20000E	201 229	0.4	4.39	30	< 10	140	< 0.5	< 2	0.06	< 0.5	13	64	54	4.83	10	< 1	0.25	< 10	0.74	345
L20400N 20100E	201 229	0.2	4.71	32	< 10	240	< 0.5	< 2	0.08	< 0.5	22	50	88	4.97	10	< 1	0.54	< 10	1.01	500
L20400N 20175E	201 229	0.2	3.82	18	< 10	100	< 0.5	< 2	0.06	< 0.5	9	61	77	5.41	10	< 1	0.24	< 10	0.79	235
L20400N 20300E	201 229	0.6	3.90	10	< 10	200	< 0.5	< 2	0.15	< 0.5	19	30	90	4.14	< 10	< 1	0.51	< 10	1.06	595
L20400N 20350E	201 229	< 0.2	1.46	< 2	< 10	80	< 0.5	< 2	0.06	< 0.5	4	4	5	2.35	< 10	< 1	0.25	< 10	0.30	450
L20500N 19850E	201 229	0.2	3.60	12	< 10	70	< 0.5	< 2	0.04	< 0.5	5	32	44	6.09	10	< 1	0.16	< 10	0.57	185
L20500N 19900E	201 229	0.2	1.97	14	< 10	90	< 0.5	< 2	0.10	< 0.5	8	39	24	2.91	< 10	1	0.14	10	0.36	325
L20500N 20000E	201 229	0.2	3.73	4	< 10	100	< 0.5	< 2	0.04	< 0.5	9	64	41	4.71	10	< 1	0.20	< 10	0.66	270
L20500N 20100E	201 229	< 0.2	3.67	18	< 10	50	< 0.5	< 2	0.03	< 0.5	6	89	27	5.63	10	< 1	0.17	< 10	0.67	155
L20600N 19950E	201 229	0.2	5.21	14	< 10	140	< 0.5	< 2	0.05	< 0.5	17	62	66	4.77	10	< 1	0.34	< 10	1.04	410
L20600N 20000E	201 229	0.2	4.40	10	< 10	60	< 0.5	< 2	0.04	< 0.5	9	41	21	5.07	10	< 1	0.22	< 10	0.58	240
L20600N 20100E	201 229	0.2	4.00	12	< 10	70	< 0.5	< 2	0.05	< 0.5	9	51	28	5.30	10	< 1	0.18	< 10	0.58	150
L20600N 20200E	201 229	< 0.2	0.90	2	< 10	10	< 0.5	< 2	0.01	< 0.5	< 1	4	4	3.28	10	< 1	0.03	< 10	0.04	45
L20600N 20300E	201 229	0.4	3.66	2	< 10	10	< 0.5	< 2	0.06	< 0.5	5	40	10	7.59	20	< 1	0.03	< 10	0.28	115
L20600N 20400E	201 229	< 0.2	1.01	< 2	< 10	< 10	< 0.5	< 2	0.04	< 0.5	3	13	7	4.41	10	< 1	0.02	< 10	0.12	90
L20600N 20500E	201 229	< 0.2	0.43	< 2	< 10	< 10	< 0.5	< 2	0.01	< 0.5	< 1	3	2	0.51	< 10	< 1	0.01	20	0.01	5
L20600N 20600E	201 229	< 0.2	0.36	10	< 10	< 10	< 0.5	< 2	0.03	< 0.5	< 1	35	1	1.55	< 10	< 1	0.01	10	0.03	45
L20600N 20700E	201 229	0.2	1.95	< 2	< 10	20	< 0.5	< 2	0.05	< 0.5	4	13	11	6.04	20	< 1	0.05	< 10	0.29	135
L20600N 20800E	201 229	< 0.2	2.00	< 2	< 10	40	< 0.5	< 2	0.09	< 0.5	6	9	3	1.58	10	< 1	0.07	< 10	0.57	170
L20600N 20900E	201 229	< 0.2	1.96	2	< 10	30	< 0.5	< 2	0.10	< 0.5	6	16	15	4.79	10	< 1	0.09	< 10	0.35	215
L20700N 19750E	201 229	0.2	1.01	4	< 10	10	< 0.5	< 2	0.02	< 0.5	2	12	6	3.41	10	< 1	0.01	10	0.04	40
L20700N 19800E	201 229	0.4	2.41	20	< 10	20	< 0.5	< 2	0.06	< 0.5	4	14	17	5.10	10	< 1	0.05	< 10	0.25	125
L20700N 19900E	201 229	0.2	1.80	26	< 10	10	< 0.5	< 2	0.02	< 0.5	3	11	11	4.80	10	< 1	0.02	< 10	0.09	65
L20700N 20000E	201 229	0.2	1.10	14	< 10	10	< 0.5	< 2	0.03	< 0.5	2	11	10	3.59	10	< 1	0.03	< 10	0.09	65
L20700N 20050E	201 229	0.4	3.14	40	< 10	20	< 0.5	< 2	0.03	< 0.5	5	14	9	8.50	30	< 1	0.04	< 10	0.32	155
L20700N 20075E	201 229	0.2	1.76	8	< 10	30	< 0.5	< 2	0.04	< 0.5	4	4	9	5.66	10	< 1	0.10	< 10	0.23	305
L20700N 20100E	201 229	0.4	2.35	12	< 10	20	< 0.5	< 2	0.03	< 0.5	3	9	13	4.50	10	< 1	0.03	< 10	0.24	115
L20700N 20200E	201 229	0.2	1.44	6	< 10	50	< 0.5	< 2	0.03	< 0.5	4	10	5	2.08	< 10	< 1	0.13	< 10	0.25	200
L20700N 20300E	201 229	< 0.2	0.98	< 2	< 10	30	< 0.5	< 2	0.08	< 0.5	4	7	16	1.47	< 10	< 1	0.15	< 10	0.37	265
L20700N 20400E	201 229	< 0.2	0.54	2	< 10	< 10	< 0.5	< 2	0.02	< 0.5	1	5	5	1.21	< 10	< 1	0.01	< 10	0.01	35
L20700N 20500E	201 229	< 0.2	0.52	< 2	< 10	< 10	< 0.5	< 2	0.08	< 0.5	2	1	4	1.16	< 10	< 1	0.09	< 10	0.16	210
L20700N 20600E	201 229	0.2	1.91	4	< 10	30	< 0.5	< 2	0.06	< 0.5	5	10	17	4.27	20	< 1	0.13	< 10	0.55	215
L20700N 20700E	201 229	< 0.2	0.89	2	< 10	< 10	< 0.5	< 2	0.02	< 0.5	2	6	8	2.81	10	< 1	0.01	< 10	0.04	35
L20700N 20800E	201 229	0.2	1.72	< 2	< 10	60	< 0.5	< 2	0.12	< 0.5	9	5	2	2.09	< 10	< 1	0.42	< 10	0.81	330
L20700N 20900E	201 229	0.8	0.46	< 2	< 10	10	< 0.5	< 2	0.83	< 0.5	< 1	1	2	0.08	< 10	< 1	0.01	< 10	0.04	20
L20800N 19625E	201 229	< 0.2	1.06	18	< 10	10	< 0.5	< 2	0.02	< 0.5	3	16	9	4.71	20	< 1	0.03	< 10	0.09	55
L20800N 19700E	201 229	0.2	2.21	< 2	< 10	30	< 0.5	< 2	0.03	< 0.5	4	24	12	4.01	10	< 1	0.03	< 10	0.25	85
L20800N 19800E	201 229	0.2	3.09	42	< 10	30	< 0.5	< 2	0.03	< 0.5	5	16	17	5.75	10	1	0.07	< 10	0.41	210
L20800N 19900E	201 229	0.4	1.86	70	< 10	10	< 0.5	< 2	0.04	< 0.5	4	6	8	3.45	10	< 1	0.06	< 10	0.18	170

CERTIFICATION: *[Signature]*



Chemex Labs Ltd.

Analytical Chemists * Geochemists * Registered Assayers

212 Brooksbank Ave., North Vancouver
British Columbia, Canada V7J 2C1
PHONE: 604-984-0221 FAX: 604-984-0218

To: BURGERT, ARND

P.O. BOX 1208
PORT MCNEILL, BC
V0N 2R0

Page Number :2-B
Total Pages :4
Certificate Date:09-SEP-1999
Invoice No. :19927561
P.O. Number :
Account :QHB

Project : OLD IRONSIDES
Comments: ATTN: ARND BURGERT CC: ARND BURGERT

CERTIFICATE OF ANALYSIS

A9927561

SAMPLE	PREP CODE	Mo ppm	Na %	Ni ppm	P ppm	Pb ppm	S %	Sb ppm	Sc ppm	Sr ppm	Ti %	Ti ppm	U ppm	V ppm	W ppm	Zn ppm
L20300N 20350E	201 229	1	0.03	38	530	8	0.04	< 2	7	6	0.26	< 10	< 10	107	< 10	82
L20400N 20000E	201 229	< 1	0.02	36	560	4	0.05	< 2	7	13	0.20	< 10	< 10	103	< 10	88
L20400N 20100E	201 229	1	0.02	65	1090	2	0.05	< 2	7	10	0.18	< 10	< 10	91	< 10	118
L20400N 20175E	201 229	< 1	0.02	42	830	6	0.08	< 2	7	7	0.15	< 10	< 10	96	< 10	82
L20400N 20300E	201 229	1	0.03	26	670	2	0.03	< 2	5	25	0.19	< 10	< 10	73	< 10	106
L20400N 20350E	201 229	5	0.02	2	510	6	0.06	< 2	3	10	0.16	< 10	< 10	19	< 10	30
L20500N 19850E	201 229	< 1	0.01	16	480	4	0.04	< 2	5	13	0.13	< 10	< 10	70	< 10	56
L20500N 19900E	201 229	1	0.01	20	620	2	0.05	< 2	4	13	0.13	< 10	< 10	60	< 10	44
L20500N 20000E	201 229	< 1	0.02	34	460	6	0.03	< 2	8	5	0.23	< 10	< 10	100	< 10	82
L20500N 20100E	201 229	< 1	0.03	22	390	6	0.04	< 2	8	4	0.24	< 10	< 10	142	< 10	66
L20600N 19950E	201 229	< 1	0.02	40	450	6	0.02	< 2	10	6	0.22	< 10	< 10	123	< 10	112
L20600N 20000E	201 229	2	0.01	24	560	2	0.03	< 2	7	5	0.19	< 10	< 10	87	< 10	70
L20600N 20100E	201 229	< 1	0.02	22	390	2	0.04	< 2	8	3	0.19	< 10	< 10	107	< 10	54
L20600N 20200E	201 229	1	0.01	< 1	190	< 2	0.01	< 2	< 1	1	0.21	< 10	< 10	41	< 10	4
L20600N 20300E	201 229	< 1	0.02	3	340	8	0.04	< 2	2	1	0.32	< 10	< 10	128	< 10	18
L20600N 20400E	201 229	< 1	0.02	< 1	150	6	0.01	< 2	1	1	0.31	< 10	< 10	125	< 10	8
L20600N 20500E	201 229	< 1	< 0.01	< 1	190	< 2	0.01	< 2	< 1	1	0.04	< 10	< 10	17	< 10	< 2
L20600N 20600E	201 229	< 1	0.01	1	130	2	0.01	< 2	< 1	2	0.07	< 10	< 10	59	< 10	< 2
L20600N 20700E	201 229	< 1	0.03	1	270	2	0.03	< 2	2	3	0.37	< 10	< 10	144	< 10	16
L20600N 20800E	201 229	< 1	0.02	3	130	4	0.05	< 2	3	5	0.25	< 10	< 10	60	< 10	26
L20600N 20900E	201 229	3	0.02	2	220	2	0.04	< 2	2	5	0.31	< 10	< 10	130	< 10	18
L20700N 19750E	201 229	2	0.01	2	200	6	0.02	< 2	< 1	3	0.31	< 10	< 10	171	< 10	8
L20700N 19800E	201 229	2	0.02	3	360	2	0.04	< 2	3	4	0.27	< 10	< 10	134	< 10	20
L20700N 19900E	201 229	1	0.01	3	330	4	0.04	< 2	1	4	0.19	< 10	< 10	80	< 10	12
L20700N 20000E	201 229	1	0.01	1	360	8	0.03	< 2	1	2	0.20	< 10	< 10	69	< 10	12
L20700N 20050E	201 229	1	0.03	< 1	390	6	0.05	< 2	3	3	0.38	< 10	< 10	112	< 10	24
L20700N 20075E	201 229	1	0.02	< 1	400	8	0.03	< 2	3	3	0.38	< 10	< 10	89	< 10	22
L20700N 20100E	201 229	1	0.01	1	320	2	0.04	< 2	2	3	0.19	< 10	< 10	73	< 10	20
L20700N 20200E	201 229	1	0.01	7	180	6	0.01	< 2	3	3	0.18	< 10	< 10	40	< 10	30
L20700N 20300E	201 229	4	0.01	2	100	2	0.01	< 2	1	3	0.16	< 10	< 10	42	< 10	30
L20700N 20400E	201 229	1	0.01	< 1	80	6	< 0.01	< 2	< 1	3	0.21	< 10	< 10	74	< 10	2
L20700N 20500E	201 229	< 1	0.01	< 1	80	< 2	0.01	< 2	< 1	2	0.11	< 10	< 10	33	< 10	20
L20700N 20600E	201 229	1	0.02	2	270	4	0.03	< 2	4	3	0.28	< 10	< 10	152	< 10	28
L20700N 20700E	201 229	1	0.01	< 1	170	6	0.02	< 2	< 1	2	0.27	< 10	< 10	129	< 10	6
L20700N 20800E	201 229	< 1	0.02	2	130	< 2	0.04	< 2	2	6	0.23	< 10	< 10	66	< 10	36
L20700N 20900E	201 229	3	0.01	1	570	< 2	0.46	< 2	< 1	38	0.01	< 10	< 10	3	< 10	8
L20800N 19625E	201 229	1	0.02	1	240	8	0.01	< 2	1	2	0.32	< 10	< 10	163	< 10	12
L20800N 19700E	201 229	< 1	0.01	6	190	2	0.02	< 2	4	4	0.21	< 10	< 10	133	< 10	22
L20800N 19800E	201 229	1	0.01	1	200	12	0.04	< 2	3	3	0.26	< 10	< 10	116	< 10	34
L20800N 19900E	201 229	1	0.01	1	260	8	0.03	< 2	1	2	0.18	< 10	< 10	71	< 10	18

CERTIFICATION:



Chemex Labs Ltd.

Analytical Chemists * Geochemists * Registered Assayers

212 Brooksbank Ave., North Vancouver
British Columbia, Canada V7J 2C1
PHONE: 604-984-0221 FAX: 604-984-0218

To: BURGERT, ARND

P.O. BOX 1208
PORT MCNEILL, BC
V0N 2R0

Project: OLD IRONSIDES
Comments: ATTN: ARND BURGERT CC: ARND BURGERT

Page Number : 3-A
Total Pages : 4
Certificate Date: 09-SEP-1999
Invoice No. : 19927561
P.O. Number :
Account : OHB

CERTIFICATE OF ANALYSIS A9927561

SAMPLE	PREP CODE	Ag ppm	Al %	As ppm	B ppm	Ba ppm	Be ppm	Bi ppm	Ca %	Cd ppm	Co ppm	Cr ppm	Cu ppm	Fe %	Ga ppm	Hg ppm	K %	La ppm	Mg %	Mn ppm
L20800N 20000E	201 229	< 0.2	2.35	234	< 10	30	< 0.5	< 2	0.04	< 0.5	4	11	9	5.68	10	< 1	0.07	< 10	0.43	190
L20800N 20100E	201 229	0.2	3.01	< 2	< 10	40	< 0.5	< 2	0.04	< 0.5	4	11	16	5.26	10	< 1	0.09	< 10	0.47	210
L20800N 20200E	201 229	< 0.2	0.45	< 2	< 10	< 10	< 0.5	< 2	0.01	< 0.5	< 1	2	3	0.12	< 10	< 1	0.01	10	< 0.01	30
L20800N 20300E	201 229	< 0.2	0.74	< 2	< 10	10	< 0.5	< 2	0.06	< 0.5	4	11	6	4.28	10	< 1	0.04	< 10	0.10	75
L20800N 20400E	201 229	< 0.2	1.29	2	< 10	10	< 0.5	< 2	0.05	< 0.5	3	10	7	1.30	10	< 1	0.05	< 10	0.36	175
L20800N 20500E	201 229	0.2	1.82	< 2	< 10	10	< 0.5	< 2	0.07	< 0.5	4	8	13	3.40	10	< 1	0.05	< 10	0.26	185
L20800N 20600E	201 229	0.8	2.28	< 2	< 10	120	< 0.5	< 2	0.16	< 0.5	7	15	18	1.72	< 10	< 1	0.52	< 10	0.93	390
L20800N 20700E	201 229	0.2	7.48	< 2	< 10	10	< 0.5	< 2	0.01	< 0.5	8	4	9	6.14	10	< 1	0.05	< 10	0.77	375
L20800N 20800E	201 229	0.2	4.83	< 2	< 10	20	< 0.5	< 2	0.06	< 0.5	6	26	17	5.09	10	< 1	0.06	< 10	0.57	235
L20800N 20900E	201 229	0.2	3.86	< 2	< 10	100	< 0.5	< 2	0.01	< 0.5	8	4	19	4.36	10	< 1	0.32	< 10	1.36	315
L20900N 19700E	201 229	0.6	2.89	2	< 10	10	< 0.5	< 2	0.03	< 0.5	4	15	12	6.80	30	1	0.03	< 10	0.10	60
L20900N 19800E	201 229	< 0.2	2.32	6	< 10	40	< 0.5	< 2	0.03	< 0.5	3	5	11	5.63	10	< 1	0.10	< 10	0.33	230
L20900N 19875E	201 229	0.2	3.90	6	< 10	40	< 0.5	< 2	0.03	< 0.5	4	9	13	5.47	10	< 1	0.08	< 10	0.41	240
L20900N 20000E	201 229	0.6	3.07	14	< 10	40	< 0.5	< 2	0.03	< 0.5	5	13	17	5.91	10	< 1	0.10	< 10	0.57	230
L20900N 20100E	201 229	1.0	3.59	< 2	< 10	30	< 0.5	< 2	0.04	< 0.5	3	8	20	5.53	10	< 1	0.06	< 10	0.21	105
L20900N 20200E	201 229	0.4	2.16	2	< 10	20	< 0.5	< 2	0.02	< 0.5	3	9	14	7.59	20	< 1	0.05	< 10	0.21	130
L20900N 20300E	201 229	0.8	2.11	8	< 10	40	< 0.5	< 2	0.02	< 0.5	3	9	9	4.98	10	< 1	0.09	< 10	0.25	190
L20900N 20400E	201 229	0.2	1.40	12	< 10	10	< 0.5	< 2	0.01	< 0.5	1	5	5	2.57	< 10	< 1	0.11	10	0.21	135
L20900N 20500E	201 229	0.2	1.15	< 2	< 10	10	< 0.5	< 2	0.06	< 0.5	5	9	6	3.72	10	< 1	0.08	< 10	0.32	160
L20900N 20600EA	201 229	0.6	2.11	6	< 10	20	< 0.5	< 2	0.04	< 0.5	3	25	15	7.58	20	< 1	0.03	< 10	0.16	90
L20900N 20600EB	201 229	1.2	3.48	< 2	< 10	100	< 0.5	< 2	0.12	< 0.5	10	22	42	3.58	< 10	< 1	0.31	< 10	0.96	425
L20900N 20700E	201 229	0.8	4.01	< 2	< 10	10	< 0.5	< 2	0.06	< 0.5	5	21	16	6.28	10	< 1	0.03	< 10	0.34	155
L20900N 20800E	201 229	0.2	3.11	< 2	< 10	20	< 0.5	< 2	0.10	< 0.5	9	32	33	4.94	10	< 1	0.06	< 10	1.01	290
L20900N 20900E	201 229	< 0.2	2.86	< 2	< 10	30	< 0.5	< 2	0.07	< 0.5	6	24	15	4.69	10	< 1	0.11	< 10	0.64	240
L2100N 19600E	201 229	0.4	2.67	6	< 10	30	< 0.5	< 2	0.03	< 0.5	3	13	13	4.92	10	< 1	0.05	< 10	0.26	145
L2100N 19700E	201 229	0.6	2.41	4	< 10	20	< 0.5	< 2	0.04	< 0.5	4	15	15	6.04	10	< 1	0.05	< 10	0.45	140
L2100N 19800E	201 229	0.2	2.24	274	< 10	40	< 0.5	< 2	0.01	< 0.5	3	35	13	5.97	10	< 1	0.08	< 10	0.18	80
L2100N 19900E	201 229	0.2	1.98	6	< 10	10	< 0.5	< 2	0.04	< 0.5	3	8	12	5.58	10	< 1	0.03	< 10	0.20	120
L2100N 20000E	201 229	1.4	3.41	< 2	< 10	70	< 0.5	< 2	0.02	< 0.5	6	4	96	8.90	10	< 1	0.24	< 10	0.55	555
L2100N 20100E	201 229	0.8	1.40	< 2	< 10	20	< 0.5	< 2	< 0.01	< 0.5	1	6	39	8.50	10	< 1	0.07	< 10	0.10	110
L2100N 20200E	201 229	0.4	2.49	< 2	< 10	50	< 0.5	< 2	0.01	< 0.5	4	7	16	6.96	20	< 1	0.15	< 10	0.39	185
L2100N 20300E	201 229	0.2	1.32	< 2	< 10	20	< 0.5	< 2	0.01	< 0.5	3	7	9	4.19	10	< 1	0.05	< 10	0.21	110
L2100N 20400E	201 229	0.2	3.48	< 2	< 10	60	< 0.5	< 2	0.03	< 0.5	8	25	16	5.14	10	< 1	0.13	< 10	1.11	440
L2100N 20500E	201 229	0.2	3.55	< 2	< 10	60	< 0.5	< 2	0.05	< 0.5	7	17	12	4.37	10	< 1	0.11	< 10	0.61	295
L2100N 20600E	201 229	0.2	2.80	< 2	< 10	50	< 0.5	< 2	0.07	< 0.5	5	18	33	2.48	< 10	< 1	0.16	< 10	0.49	310
L2100N 20700E	201 229	< 0.2	2.94	< 2	< 10	20	< 0.5	< 2	0.08	< 0.5	8	15	12	4.26	10	< 1	0.10	< 10	0.69	300
L2100N 20800E	201 229	0.2	4.74	< 2	< 10	30	< 0.5	< 2	0.06	< 0.5	6	20	19	4.64	10	< 1	0.09	< 10	0.50	230
L2100N 20900E	201 229	< 0.2	2.60	2	< 10	120	< 0.5	< 2	0.08	< 0.5	10	12	11	3.46	10	< 1	0.41	< 10	1.01	440
S137	201 229	0.2	3.83	20	< 10	180	< 0.5	< 2	0.05	< 0.5	3	20	17	3.76	< 10	< 1	0.28	< 10	0.99	355
S138	201 229	< 0.2	0.85	28	< 10	20	< 0.5	< 2	0.01	< 0.5	1	4	7	2.65	< 10	< 1	0.04	< 10	0.08	65

CERTIFICATION:



Chemex Labs Ltd.

Analytical Chemists * Geochemists * Registered Assayers

212 Brooksbank Ave., North Vancouver
British Columbia, Canada V7J 2C1
PHONE: 604-984-0221 FAX: 604-984-0218

To: BURGERT, ARND

P.O. BOX 1208
PORT MCNEILL, BC
V0N 2R0

Project: OLD IRONSIDES
Comments: ATTN: ARND BURGERT CC: ARND BURGERT

Page Number :3-B
Total Pages :4
Certificate Date:09-SEP-1999
Invoice No. :19927561
P.O. Number :
Account :GHB

CERTIFICATE OF ANALYSIS A9927561

SAMPLE	PREP CODE	Mo ppm	Na %	Ni ppm	P ppm	Pb ppm	S %	Sb ppm	Sc ppm	Sr ppm	Ti %	Tl ppm	U ppm	V ppm	W ppm	Zn ppm
L20800N 20000E	201 229	1	0.02	2	280	2	0.03	< 2	3	3	0.29	< 10	< 10	86	< 10	30
L20800N 20100E	201 229	< 1	0.02	1	270	2	0.03	< 2	4	3	0.23	< 10	< 10	103	< 10	34
L20800N 20200E	201 229	< 1	< 0.01	< 1	110	< 2	0.01	< 2	< 1	1	0.02	< 10	< 10	5	< 10	< 2
L20800N 20300E	201 229	1	0.03	< 1	240	4	0.02	< 2	1	3	0.39	< 10	< 10	190	< 10	6
L20800N 20400E	201 229	2	0.01	2	140	8	0.01	< 2	2	4	0.20	< 10	< 10	57	< 10	20
L20800N 20500E	201 229	2	0.02	< 1	150	< 2	0.02	< 2	1	4	0.31	< 10	< 10	98	< 10	18
L20800N 20600E	201 229	< 1	0.03	7	220	< 2	0.10	< 2	4	12	0.17	< 10	< 10	49	< 10	58
L20800N 20700E	201 229	< 1	0.02	< 1	260	6	0.06	< 2	6	< 1	0.28	< 10	< 10	78	< 10	26
L20800N 20800E	201 229	< 1	0.02	4	240	2	0.04	< 2	4	3	0.23	< 10	< 10	94	< 10	30
L20800N 20900E	201 229	< 1	0.02	1	280	2	0.03	< 2	9	1	0.18	< 10	< 10	108	< 10	44
L20900N 19700E	201 229	< 1	0.03	< 1	380	8	0.04	< 2	1	1	0.42	< 10	< 10	182	< 10	10
L20900N 19800E	201 229	< 1	0.02	< 1	440	2	0.03	< 2	4	3	0.26	< 10	< 10	62	< 10	28
L20900N 19875E	201 229	< 1	0.02	1	400	8	0.05	< 2	3	3	0.20	< 10	< 10	65	< 10	34
L20900N 20000E	201 229	< 1	0.01	1	320	2	0.04	< 2	4	2	0.21	< 10	< 10	93	< 10	38
L20900N 20100E	201 229	< 1	0.01	< 1	430	8	0.06	< 2	3	4	0.22	< 10	< 10	101	< 10	20
L20900N 20200E	201 229	< 1	0.02	1	470	2	0.05	< 2	3	3	0.25	< 10	< 10	149	< 10	26
L20900N 20300E	201 229	< 1	0.02	< 1	230	10	0.05	< 2	4	3	0.28	< 10	< 10	113	< 10	22
L20900N 20400E	201 229	2	0.01	< 1	200	2	0.02	< 2	1	1	0.16	< 10	< 10	45	< 10	10
L20900N 20500E	201 229	1	0.03	1	280	< 2	0.02	< 2	1	3	0.35	< 10	< 10	155	< 10	14
L20900N 20600EA	201 229	< 1	0.02	3	420	8	0.03	< 2	2	2	0.26	< 10	< 10	163	< 10	18
L20900N 20600EB	201 229	4	0.03	9	280	2	0.04	< 2	4	9	0.19	< 10	< 10	75	< 10	56
L20900N 20700E	201 229	< 1	0.02	2	270	4	0.13	< 2	4	3	0.28	< 10	< 10	112	< 10	20
L20900N 20800E	201 229	< 1	0.03	9	180	2	0.01	< 2	5	5	0.31	< 10	< 10	116	< 10	50
L20900N 20900E	201 229	< 1	0.02	5	250	2	0.03	< 2	3	4	0.25	< 10	< 10	84	< 10	30
L2100N 19600E	201 229	< 1	0.01	3	340	8	0.03	< 2	3	3	0.20	< 10	< 10	93	< 10	24
L2100N 19700E	201 229	1	0.02	1	300	4	0.03	< 2	3	3	0.28	< 10	< 10	149	< 10	34
L2100N 19800E	201 229	< 1	0.01	16	500	10	0.04	< 2	2	4	0.19	< 10	< 10	72	< 10	26
L2100N 19900E	201 229	< 1	0.02	< 1	280	8	0.02	< 2	1	3	0.27	< 10	< 10	133	< 10	14
L2100N 20000E	201 229	< 1	0.02	< 1	680	10	0.07	< 2	7	4	0.22	< 10	< 10	127	< 10	82
L2100N 20100E	201 229	< 1	0.01	< 1	610	10	0.05	< 2	1	1	0.22	< 10	< 10	102	< 10	16
L2100N 20200E	201 229	< 1	0.02	< 1	450	6	0.04	< 2	4	3	0.30	< 10	< 10	160	< 10	38
L2100N 20300E	201 229	1	0.01	< 1	390	4	0.03	< 2	1	3	0.22	< 10	< 10	109	< 10	12
L2100N 20400E	201 229	< 1	0.02	4	270	4	0.04	< 2	9	2	0.21	< 10	< 10	122	< 10	50
L2100N 20500E	201 229	< 1	0.01	4	260	4	0.03	< 2	3	1	0.26	< 10	< 10	108	< 10	32
L2100N 20600E	201 229	1	0.01	4	280	< 2	0.04	< 2	2	7	0.14	< 10	< 10	48	< 10	42
L2100N 20700E	201 229	< 1	0.03	4	270	< 2	0.03	< 2	2	3	0.30	< 10	< 10	92	< 10	32
L2100N 20800E	201 229	< 1	0.02	3	320	2	0.06	< 2	4	6	0.19	< 10	< 10	78	< 10	28
L2100N 20900E	201 229	< 1	0.03	5	200	4	0.02	< 2	3	3	0.30	< 10	< 10	88	< 10	40
S137	201 229	< 1	0.02	3	490	2	0.10	< 2	9	14	0.14	< 10	< 10	75	< 10	58
S138	201 229	< 1	0.01	1	230	6	0.02	< 2	1	3	0.15	< 10	< 10	48	< 10	8

CERTIFICATION:



Chemex Labs Ltd.

Analytical Chemists * Geochemists * Registered Assayers

212 Brooksbank Ave., North Vancouver
British Columbia, Canada V7J 2C1
PHONE: 604-984-0221 FAX: 604-984-0218

To: BURGERT, ARND

P.O. BOX 1208
PORT MCNEILL, BC
V0N 2R0

Project: OLD IRONSIDES
Comments: ATTN: ARND BURGERT CC: ARND BURGERT

Page Number : 1-A
Total Pages : 1
Certificate Date: 10-SEP-1999
Invoice No. : I9927530
P.O. Number :
Account : QHB

CERTIFICATE OF ANALYSIS A9927530

SAMPLE	PREP CODE	Ag ppm	Al %	As ppm	B ppm	Ba ppm	Be ppm	Bi ppm	Ca %	Cd ppm	Co ppm	Cr ppm	Cu ppm	Fe %	Ga ppm	Hg ppm	K %	La ppm	Mg %	Mn ppm
P197001	205 226	1.0	5.75	8	< 10	10	< 0.5	< 2	3.08	< 0.5	14	24	65	0.28	10	1	0.31	< 10	0.43	280
P197002	205 226	0.2	2.70	< 2	< 10	120	< 0.5	< 2	0.06	< 0.5	5	81	26	3.33	< 10	< 1	0.41	10	0.59	125
P197003	205 226	< 0.2	1.57	< 2	< 10	70	< 0.5	< 2	0.04	< 0.5	25	56	36	3.17	< 10	< 1	0.44	< 10	0.68	240
P197004	205 226	< 0.2	1.33	4	< 10	30	< 0.5	< 2	0.18	< 0.5	12	46	28	4.27	< 10	1	0.41	< 10	0.85	250
P197005	205 226	0.8	2.73	176	< 10	120	< 0.5	< 2	0.15	< 0.5	27	35	45	3.32	< 10	< 1	1.20	< 10	1.32	430
P197006	205 226	0.2	4.86	6	< 10	140	< 0.5	< 2	1.96	< 0.5	11	85	21	3.14	< 10	1	0.74	< 10	1.18	345
P197007	205 226	0.2	1.41	< 2	< 10	80	< 0.5	< 2	0.03	< 0.5	< 1	68	1	2.06	< 10	< 1	0.83	< 10	1.13	320
P197008	205 226	< 0.2	3.10	30	< 10	90	< 0.5	< 2	0.78	< 0.5	8	28	16	2.82	< 10	< 1	0.62	< 10	1.08	310
P197009	205 226	< 0.2	1.23	94	< 10	120	< 0.5	< 2	0.01	< 0.5	1	48	8	2.75	< 10	< 1	0.34	< 10	0.51	105
P197010	205 226	0.2	1.57	34	< 10	120	< 0.5	< 2	0.02	< 0.5	< 1	62	7	1.96	< 10	< 1	0.36	< 10	0.60	90
P197011	205 226	< 0.2	1.65	2	< 10	170	< 0.5	< 2	0.02	< 0.5	4	87	5	1.48	< 10	< 1	0.48	20	0.72	230
P197012	205 226	< 0.2	4.00	6	< 10	220	< 0.5	< 2	1.76	< 0.5	7	96	8	3.57	< 10	< 1	0.67	< 10	0.80	370
P197013	205 226	0.2	0.58	2	< 10	60	< 0.5	< 2	0.54	< 0.5	13	38	80	1.25	< 10	< 1	0.07	< 10	0.15	105
P197014	205 226	< 0.2	2.50	6	< 10	260	< 0.5	< 2	0.32	< 0.5	13	70	20	3.26	< 10	< 1	0.81	< 10	1.17	410
P197015	205 226	4.0	0.49	8	< 10	50	< 0.5	2	0.07	0.5	2	202	1305	7.27	< 10	< 1	0.06	< 10	0.13	195
P197016	205 226	0.6	1.15	6	< 10	70	< 0.5	< 2	0.03	< 0.5	7	83	35	5.55	< 10	< 1	0.23	< 10	0.53	150
P197017	205 226	< 0.2	1.59	2	< 10	20	< 0.5	< 2	1.72	< 0.5	8	43	22	5.13	< 10	< 1	0.08	< 10	0.26	270
P197018	205 226	0.8	3.50	< 2	< 10	180	< 0.5	< 2	0.40	< 0.5	16	54	179	6.28	< 10	< 1	0.55	< 10	1.05	505
P197019	205 226	69.4	0.06	32	< 10	10	< 0.5	< 2	< 0.01	< 0.5	2	101	22	3.94	< 10	< 1	0.04	< 10	< 0.01	5
P197020	205 226	1.4	1.30	4	< 10	70	< 0.5	< 2	0.04	< 0.5	8	77	24	2.15	< 10	< 1	0.10	10	0.86	205
P197021	205 226	1.0	1.90	10	< 10	80	< 0.5	< 2	0.06	< 0.5	9	57	29	4.53	< 10	< 1	0.10	< 10	1.05	895

CERTIFICATION: _____



Chemex Labs Ltd.

Analytical Chemists * Geochemists * Registered Assayers

212 Brooksbank Ave., North Vancouver
British Columbia, Canada V7J 2C1
PHONE: 604-984-0221 FAX: 604-984-0218

To: BURGERT, ARND

P.O. BOX 1208
PORT MCNEILL, BC
V0N 2R0

Project: OLD IRONSIDES
Comments: ATTN: ARND BURGERT CC: ARND BURGERT

Page Number :1-B
Total Pages :1
Certificate Date: 10-SEP-1999
Invoice No. :19927530
P.O. Number :
Account :QHB

CERTIFICATE OF ANALYSIS A9927530

SAMPLE	PREP CODE	Mo ppm	Na %	Ni ppm	P ppm	Pb ppm	S %	Sb ppm	Sc ppm	Sr ppm	Ti %	Tl ppm	U ppm	V ppm	W ppm	Zn ppm
P197001	205 226	6	0.27	2	790	6	2.01	8	3	139	0.12	< 10	10	34	< 10	38
P197002	205 226	< 1	0.02	14	390	2	0.55	< 2	3	11	0.05	< 10	10	27	< 10	58
P197003	205 226	< 1	0.02	7	200	2	2.68	< 2	1	6	< 0.01	< 10	< 10	8	< 10	46
P197004	205 226	< 1	0.07	3	530	< 2	2.60	< 2	19	7	0.06	< 10	< 10	136	< 10	64
P197005	205 226	< 1	0.03	17	650	< 2	0.65	2	8	3	0.15	< 10	< 10	102	< 10	66
P197006	205 226	< 1	0.39	9	230	2	1.18	< 2	7	168	0.06	< 10	< 10	78	< 10	64
P197007	205 226	< 1	0.04	< 1	60	2	0.31	< 2	4	6	0.07	< 10	< 10	106	< 10	28
P197008	205 226	< 1	0.12	5	320	4	0.82	< 2	5	68	0.06	< 10	< 10	42	< 10	46
P197009	205 226	< 1	0.02	2	280	4	0.34	< 2	< 1	6	< 0.01	< 10	< 10	6	< 10	12
P197010	205 226	< 1	0.03	< 1	130	2	0.27	< 2	1	8	0.01	< 10	< 10	9	< 10	12
P197011	205 226	< 1	0.03	3	60	2	0.40	2	< 1	13	< 0.01	< 10	< 10	3	< 10	40
P197012	205 226	3	0.15	11	290	< 2	0.77	< 2	10	159	0.17	< 10	< 10	73	< 10	66
P197013	205 226	< 1	0.09	1	1620	< 2	0.51	< 2	3	21	0.05	< 10	< 10	15	< 10	20
P197014	205 226	< 1	0.07	6	1010	< 2	0.72	2	11	15	0.11	< 10	< 10	105	< 10	62
P197015	205 226	11	0.04	4	160	10	2.59	8	1	10	0.04	< 10	10	5	< 10	50
P197016	205 226	8	0.04	14	570	< 2	0.67	< 2	2	23	0.03	< 10	10	27	< 10	36
P197017	205 226	< 1	0.07	8	1130	< 2	0.01	4	3	158	0.14	< 10	10	85	< 10	32
P197018	205 226	< 1	0.13	3	1040	< 2	0.87	2	10	38	0.10	< 10	10	88	< 10	96
P197019	205 226	12	< 0.01	4	10	< 2	3.36	8	< 1	3	< 0.01	< 10	< 10	< 1	< 10	2
P197020	205 226	2	0.01	3	70	< 2	0.84	2	1	15	< 0.01	< 10	< 10	8	< 10	36
P197021	205 226	8	0.05	6	240	2	1.01	4	2	11	< 0.01	< 10	< 10	16	< 10	34

CERTIFICATION:



Chemex Labs Ltd.

Analytical Chemists * Geochemists * Registered Assayers
212 Brooksbank Ave., North Vancouver
British Columbia, Canada V7J 2C1
PHONE: 604-984-0221 FAX: 604-984-0218

To: BURGERT, ARND

P.O. BOX 1208
PORT MCNEILL, BC
V0N 2R0

Project : OLD IRONSIDES
Comments: ATTN: ARND BURGERT CC: ARND BURGERT

Page Number :1
Total Pages :1
Certificate Date: 22-SEP-1999
Invoice No. :19929087
P.O. Number :
Account :QHB

CERTIFICATE OF ANALYSIS A9929087

SAMPLE	PREP CODE		Au ppb FA+AA									
P197019	244	--	1700									

CERTIFICATION: *Markus...*

APPENDIX III
STATEMENT OF EXPENDITURES

Old Ironsides 2 Mineral Claim
1999 Statement of Expenditures

Prospecting			
Prospector			
11 days	@ \$225 /day		\$ 2,475
Camp costs			
11 days	@ 45 /day		495
Laboratory			
21 rock ICP	@ 13 /sample		273
1 rock Au	@ 10 /sample		10
Mapping			
Geologist			
9 days	@ 275 /day		2,475
Camp costs			
9 days	@ 45 /day		405
Soil Sampling			
Soil Sampler			
8 days	@ 180 /day		1,440
Camp costs			
8 days	@ 45 /day		360
Laboratory			
123 soil samples	@ 9 /sample		1,107
Freight			
shipping, Powell River - Vancouver			86
Supplies			
flagging tape, soil bags, tags, etc.			109
General			
4 wheel drive truck			
28 days	@ 70 /day		1,960
fuel and oil			238
drafting			752
report preparation; photocopying			625
helicopter charter			1,752
Total			<u>\$ 14,562</u>