

**ASSESSMENT REPORT  
EAGLET PROPERTY**

New Assay Data from  
1980-1984 Drill Core Samples  
DDH 08-80, 59-82, & DDH 82 to 104, 110 to 114-84  
Prepared by Eaglet Mines

Including Structural Data from  
DDH Logs 1 to 2, 6 to 13, 17 to 50, 59 to 67, 82, 85 to 89-84,  
& 111-85

**Q CLAIMS** (Eaglet Property)  
Tenure Number 401809  
**Cariboo Mining Division**

**N.T.S Map Area 93A-10W**  
Latitude 52°33' north; Longitude 121°00' west

Minfile Number **093A 046**


**For:**

**FREEMPORT RESOURCES INC.**  
8711 Elsmore Road  
Richmond, B.C. V7C 2A4

**By:**

**Zdenek D. Hora, M.Sc., P.Geo.**  
June 17, 2005

Figure 1. Property Location Map



View of Eaglet from Quesnel Lake, 1980's (Eaglet Mines photo)



## TABLE OF CONTENTS

	<i>Page</i>
1.0 Introduction	1
2.0 History	2
3.0 Geology	5
4.0 Exploration	6
5.0 Geochemical analyses	7
6.0 Drill Hole Logs	8
7.0 Conclusions and recommendations	10
8.0 References	11
9.0 Statement of qualifications	12

### Figures

	<i>After</i>
1. Property Location Map	Title Page
2. Mineral claim map	1
3. Geology map	5
4. Drill hole location map	7

### Appendices

A	Itemized Cost Statement
B	Diamond Drill Hole Pulp & Core Samples Collected from Site for Re-Assay
C	Analytical results Certificates of Analysis
D	Drill Hole Summary 1975 Diamond Drilling 1981-83 Diamond Drilling 1984-1985 Diamond Drilling
E	Logged Drill Hole Coordinates
F	Drill Hole Logs

## 1.0 Introduction

The purpose of this report is to interpret new results derived from assaying drill core and drill core pulps prepared in the mid 1980's by Eaglet Mines, and collected recently from the site by Freeport Resources Inc., the property owners. The results were analyzed in conjunction with review of sixty newly completed drill holes logged by structural geologist Dr. Peter Read.

The Q Claims are located in the Cariboo region of central B.C. (see Figure 1, previous page). They are accessed by road from William's Lake through Horsefly to the south shore of Quesnel Lake, and then by boat to the site. In recent times, logging roads from Likely have been extended and it is believed that construction of one small bridge at the head of the north arm of Quesnel Lake would result in road access to the claims.

Fluorite showings on the Eaglet site have been known since 1947. Extensive exploration took place at the property from 1965 to 1967, and again between 1973 and 1984. This work outlined mineralization over an area of approximately 1500 by 900 metres.

During the course of exploration to 1984, a total of 126 surface drill holes measuring 19,687 metres were completed. In addition, two adits, No. 1 of 292 metres and No. 2 of 373 metres, were driven to test the mineralization underground. Nine flat holes totalling 1,525 metres were drilled from No. 1 Adit.

In spite of early suggestions of similarity to porphyry mineralization systems, it is believed that fluorite and silver were consistently the primary focus in all collected samples and assaying for other minerals with economic potential was either very limited or not recorded. It is known from review of unpublished Eaglet Mines data that a commercial grade molybdenum concentrate was prepared from ore from Adit #2 in 1984. Molybdenite was previously unknown at this property and commercial concentrations were encountered rather unexpectedly during the driving of Adit #2.

**FREEPORT RESOURCES**

TO CLAIM STAKE:  
 LATITUDE: 52°34' 13"  
 LONGITUDE: 121°0' 4"

FOREST REGION: CARIBOO DISTRICT: QUESNEL  
 B.C.G.S. MAP: 53A.055/056

SCALE: 1:120,000

DATUM: NAD 83 ZONE: 10


MAPPED BY: SABREX

DATE DRAWN: 04/26/03 DRAWN BY: S. SHIPLEY

WISN. FILE NO.: SABREX5.DGN

DATE REVISED:

QUESNEL LAKE  
 (NORTH ARM)



Wasiko

Figure 2. Mineral claim map

The property was partly abandoned in 1991, and entirely in 1997. It was restaked in 1994 and then again in 2003 by Freeport Resources Inc. All costs claimed in this Assessment Report represent new previously unreported results, paid for exclusively by Freeport Resources Inc.

## **2.0 History**

The Barrett Creek canyon fluorite showings were staked in 1947 by H Forster of Kamloops. A few scattered shallow trenches were dug in 1965. The following year, Canex Aerial Exploration Ltd, later Placer Development Ltd., optioned the property and conducted trenching, geochemical soil sampling and percussion drilling. The option was cancelled in 1967.

In 1973, Eaglet Mines Limited commenced an exploration program which continued until 1984. In 1985, Eaglet Mines hired Dr. Peter Read to undertake a structural study of the deposit to understand better the distribution of mineralized zones. After completing the fieldwork, the analyses of collected data did not proceed and work remained uncompleted.

Eaglet Mines filed three assessment reports during this period - No. 5639 dated September 26<sup>th</sup>, 1975, No. 9515 dated April 22, 1981, and No. 10,447 dated May 4, 1982. After that time, commodity prices underwent significant changes. From 1984 to 1985, international fluorspar prices decreased almost 50% from US\$ 130-210/metric tonne to US\$ 72-115/metric tonne. Between 1980 and 1983, molybdenum prices decreased from US \$25/kilogram to below US \$10/kilogram. Some claims lapsed in 1990 and 1991, with the remainder in 1997.

The property was first staked by Freeport Resources Inc. from May to July 1994 to cover the exposed fluorite zones. The claims were recently restaked in April 2003, with a digital GPS survey completed.


Current claim holdings are as follows:

Tenure number	Claim name	Date recorded	Owner
401809	Q	April 11, 2003	Freeport Resources Inc.

Since Eaglet Mines abandoned the property, work by Freeport Resources Inc. has been limited in scope. In 1994, a property visit was conducted from September 15 to 17, including selective mapping and sampling of Adit Number 2, by geologist, John Ball.


Physical work at the site has included rehabilitation of the access road, drill core and core racks, and the chip shed storage, as well as shoring the entry portal of Adit #2.

*Construction of new bridge, rehabilitation of core racks*




Two years later, geologists Leo King and Virginia Kuran also visited the site and sampled number 2 Adit, conducted geological surveying, located existing drill hole collars and platforms, and examined and sampled drill core.

*Existing drill collars*



*Rehabilitation of Adit #2*



Freeport has since collected a total of 426 samples from the 1983-1985 drilling programs and which were still stored on the site. These samples were sent for ICP 31 and 34 elements analyses, and also fluorine.



### 3.0 Geology

The mineralization is hosted by Quesnel Lake orthogneiss of Late Devonian to Early Mississippian age near the contact with Late Proterozoic Snowshoe Group metasedimentary rocks (see Figure 3 next page). Some of the mineralization is also found in Snowshoe Group rocks.

The contact of the orthogneiss with metasediments is well exposed in Barrett Creek canyon which crosses the mineralized zone from north-east to south-west. In this outcrop, the intrusive contact where granitic gneiss crosscuts metasediments is clearly visible, with large blocks of metasediments included within the orthogneiss near its margins.

Fluorite occurs as disseminated grains, thin fracture fillings, irregular veins up to 10 centimeters thick and replacement bodies and pots within the gneiss. The mineralization seems to be intimately associated with the late aplitic dikes intruding the orthogneiss body. Small quantities of galena, sphalerite, chalcopyrite, pyrite and locally molybdenite grains are common visible accessories with fluorite mineralization. Celestite was identified in several drill holes and as irregular veins in Adit No.2. ICP analytical results indicate widespread distribution of the celestite throughout the deposit. Earlier reported scheelite and wolframite have not been confirmed. Widespread presence of a salmon pink feldspar in the fluorite mineralized area resembles potassic alteration dominated by K-feldspar commonly associated with porphyry Cu-Mo deposits. Gypsum as a joint filling was identified by Peter Read during the relogging of drill core as well when mapping Adit #2.

The age of muscovite collected by Peter Read from the mineralized zone suggests an age of 127 $\pm$ 4 Ma, with the generally considered less accurate fission track dating of fluorite from the claims suggesting an age of 104.6 $\pm$ 6 Ma. Both cases indicate an Upper Cretaceous age. The important B.C. porphyry molybdenum properties of Endako, Boss Mountain and MAC are also Upper Cretaceous in age.

Figure 3. Geology Map (after Pell & Fontaine, 1989)




Figure 16. Geological map of the Eaglet Prospect. Compiled from Pell and Fontaine, 1989; P. Read, personal communication, 1989 and drill-hole data from Company and Assessment Reports.

The preliminary conclusions of the structural study indicate that mineralized zones are folded by late, broad open upright folds and locally truncated by faulting (P.Read, personal communication, 1989).

#### 4.0 Exploration

Two adits and in total 135 diamond drill holes were completed by previous owners in the course of exploring the property. To process core samples and those from channel sampling from the adit walls, a small laboratory was built on the site. In the lab, the core was split, crushed and ground by laboratory standards. The resulting pulps were shipped for assays to the Kamloops Research & Assay Laboratory Ltd.

A variety of pulp samples were stored inside the lab shed in several cardboard boxes. These include 189 samples processed from the 1983 drill program (DDH 83 to DDH104), as well as 11 boxes of pulp samples from the ribs and rounds of Adit 2. The pulps were stored in small paper bags with approximately 200 grams of ground sample each, preserved in excellent condition.

A shed housing chip samples from most drill holes and the adits is also located on site. These samples are available for further analyses and study.

*Chip Shed*




The available data from previously filed assessment reports is very incomplete and a considerable amount of the information generated by the work remains unavailable. From the 1975 program of 9 drill holes, no logs were filed and only some samples were analyzed for CaF<sub>2</sub> only (Assessment Report 5639). The seven underground drill holes completed in 1980 are not documented at all. In 1981, the company drilled forty vertical drill holes. Only 7 (S1 to S8) have reported logs and CaF<sub>2</sub> and Ag assay results (Assessment Report 9515). Drill holes completed during 1982 and 1983 season have available information on CaF<sub>2</sub> and Ag assay results, but drill hole logs have been reported for nine holes only (S41 to S49). In total, 54 drill holes (S50 to S104) are lacking any core description from geological staff. There is no assay data from Adits #1 and #2, and no Mo data is available in spite of the fact that parts of Adit 2 were driven to outline a Mo mineralized zone and a processing test to produce a commercial grade molybdenum concentrate was carried out in the Kamloops laboratory.

## **5.0 Geochemical Analyses**

In order to obtain a better idea about distribution of other minerals present in the deposit and to help to interpret the deposit type, Freeport Resources Inc. collected the preserved pulp samples from the on-site lab, and sealed them in plastic bags for longer term storage. A number of representative split drill core samples were also collected for analysis. Sample locations are shown on Figure 4 on the following page.

The pulps and core samples were then sent to several commercial laboratories for ICP 31 and 34 elements analyses, REE, gold fire assays and whole rock analyses. In total, 405 samples were assayed by Assayers Canada (Min-En Labs), and 20 by ACME Analytical Laboratories Ltd., and 1 by Chemex Laboratory in Vancouver. The same suite of samples was later analyzed for fluorine to correlate the ICP data with the main exploration target.

Figure 4. Drill Hole Location Map (1984)



Detailed analytical procedures are described on assay sheets from ACME Analytical Laboratories Ltd, and as separate description sheets attached to the certificates of analyses.

## 6.0 Drill Hole Logs

Sixty drill holes were logged in 1984 and 1985 by structural geologist Dr. Peter Read, but his work was never completed. These logs were recently purchased by Freeport Resources, reviewed and compiled into a comprehensive digital database for further work and analysis. The 60 holes are shown on Figure 4, with coordinates and collar elevations listed in Appendix E. The log descriptions are attached as Appendix F. The drill hole logs provide an extensive list and description of veins, as well as rock alteration encountered in the core. Lithologies are described in detail in most of the logs.

From review of the previously unfinished drill core logs, fluorite mineralization seems to be associated with late, mostly unfoliated white and pink aplite sills intruding the Quesnel lake orthogneiss and the schists of the Snowshoe Group. Fluorite mineralization forms sheet-like zones up to 30 meters thick with grade ranging from a few percent to 43% CaF<sub>2</sub>. The best zone of fluorite mineralization is located at the contact between the Snowshoe schist and the orthogneiss and forms a sheet that extends for approximately 100 m east-west and approximately 150 m north-south including the northern arm of Adit #2. Its western boundary coincides with the western arm of Adit #2. The mineralized zone is intercepted in drill holes at depths between 100 to 150 m and is up to 30 m thick with grade of ~10% CaF<sub>2</sub>. This zone seems to taper out outside of the area mentioned above. Other thinner fluorite zones are intercepted in drill holes below and above the main one. A three dimensional map of the mineralized zones intercepted in drill holes would help to visualize the spatial distribution, the grade and volume of the fluorite mineralization.

Several different types of veins were encountered in drill core. In general, veins are more numerous and more varied at depth than at the top of the holes. The main veins seen in drill holes are variant of the following: Quartz ± pyrite ± calcite ± fluorite veins; fluorite ± pyroxene ± K-feldspar ± quartz ± pyrite veins; kaolinite veins; gypsum ± calcite veins; quartz veins; quartz ± Kfeldspar ± pyrite veins. Molybdenite, galena and sphalerite are sometimes seen in some veins, mostly associated with fluorite. In spite of high strontium values identified by ICP analyses, no celestite has been mentioned in these logs. The white, fine-grained nature of this mineral may have been confused with calcite or fluorite.

Slight to moderate, rarely intense, alteration of the host rock is also common in the drill core. The alteration minerals mentioned include K-feldspar, biotite, sericite, muscovite, gypsum, pyroxene, chlorite, and possible clay. Epidote is seldom mentioned as either vein component or rock alteration. It is unknown at this time if the alteration minerals form halos or zones of different intensity and type of alteration as is often seen in porphyry copper and molybdenum systems (i.e. propylitic, argillic, phyllic and potassic alteration zones). It would be interesting to study the spatial distribution of the alteration minerals and vein types encountered in the drill cores to determine whether they show similarity with any porphyry copper and molybdenum deposit model.

## 7.0 Conclusions and recommendations

High values of strontium in all samples indicate the presence of celestite disseminated throughout the whole area mineralized with fluor spar. Potassium is considerably higher than sodium in most of the samples, which points to potentially widespread potassium alteration.

Molybdenum values are elevated in many samples, but independently of fluorine, strontium, lead, copper and zinc. Barium values are less than one tenth of strontium content. Dark purple fluorite is enriched in the thorium and light REE relative to the green and colorless varieties. About 25% of molybdenum ICP values are between 0.001% and 0.02%.


First, available data from published and unpublished sources should be used to establish the most probable deposit type for mineralization at the Q claims. Secondly, based on the conclusion, the mineral target - most probably molybdenum - should be identified and the proposed exploration program developed with the deposit model in mind. Thirdly, samples from Adit #2 should be analyzed for  $\text{MoS}_2$ . Optional use of geophysics to identify potentially higher sulphide concentrations could also be considered.

A three dimensional diagram of alteration and mineralization intercepted in drill holes and Adits would be of assistance to further exploration efforts at this property.

Respectfully submitted.



Z.D. Hora, M.Sc., P. Geo.  
Victoria, B.C.  
June 17, 2005





## 8.0 References

Ball, C.W. (1980): Eaglet Mines Limited. Extension of Geological Report Fluorite Property Eaglet and Eagle Claims; unpublished company report.

Ball, C.W. and Boggaram,G. (1985): Geological investigation of the Eaglet fluorspar deposit; Mining Magazine, June 1985, pages 506 - 509.

McCammon, J.W. (1966): Fluorite: Eaglet Group; B.C. Ministry of Mines, Annual Report 1965, pages 263 - 264.

McDougall, J.J. and Ball, J.C. (1994): Eaglet Fluorite Project. Review of 1973 to 1984 exploration results and proposed exploration program. Unpublished report for Freeport Resources Inc.,11 pages.

Mitchell, J.A. (1975): Eaglet Mines Limited. Diamond Drilling Progress Report; B.C. Ministry of Energy and Mines, Assessment Report 5639.

Pell, J. (1992): Fluorspar and fluorine in British Columbia; B.C. Ministry of Energy, Mines and Petroleum Resources, Open File 1992 - 16, 81 pages.

Robertson, A. (1981): Eaglet Mines Limited. Drilling Report April 22, 1981, B.C. Ministry of Energy and Mines, Assessment Report 9515.

Robertson, A. (1982): Eaglet Mines Limited. Drilling Report May 4, 1982, B.C. Ministry of Energy and Mines, Assessment Report 10447.

Taggart, L.P. (1984): Eaglet Mines Limited, Quesnel Lake Fluorspar Project, Horsefly, British Columbia. Preliminary Feasibility Study; Kilborn Engineering (B.C.) Ltd., Volumes I and II.

## 9.0 Statement of qualifications

I, Zdenek D. Hora, M.Sc., P.Geo., of Victoria, British Columbia, do hereby certify that:

I am a Consulting Geologist and since 1975, a Registered Professional Geoscientist in British Columbia and previously in Alberta, residing at 3657 Doncaster Drive, Victoria, B.C., V8P 3W8.

I graduated from Charles University of Prague, Czechoslovakia with a M.Sc. Degree in geology in 1958. Since graduation, I have been continuously practicing my profession in Europe and overseas, and since 1971 in Canada, namely in Quebec, Alberta, the N.W.T. and British Columbia. My work has largely been focussed on the geology, exploration and evaluation of industrial minerals deposits. From 1978 to 1984, I was the Industrial Minerals Specialist for the British Columbia Ministry of Energy, Mines and Petroleum Resources. From 1984 to 1999, I acted as the Program Manager for industrial minerals inventory and market studies in the province. Since my retirement in 1999, I am consulting in the field of industrial minerals – property assessment and evaluation, tenure aspect of industrial minerals in B.C. and its historical development, aggregate prospecting and deposit models for a wide range of industrial minerals. My professional activities included teaching industrial minerals courses (i.e. University of Victoria -- Economic Geology; B.C. Ministry of Energy, Mines and Petroleum Resources, B.C. and Yukon Chamber of Mines, and Geological Association of Canada – Courses for Prospectors). I have previously served as Chairman of the Industrial Minerals Division of the Canadian Institute of Mining, Metallurgy and Petroleum (CIM), and was the organizer and Co-Chairman of the 27<sup>th</sup> FORUM on Geology of Industrial Minerals and several other symposiums dealing with industrial minerals. From 1995 to 2000, I was part of the CIM Standing Committee on Reserve Definitions representing the CIM Industrial Minerals Division. I am presently a Consulting Geologist and have been so since June, 1999. As a result of my experience and qualifications, I am a Qualified Person as defined in N.P. 43-101.


This report is based on several property visits between 1979 and 1983, published and unpublished reports filed with the B.C. Ministry of Energy and Mines and information provided to me by Freeport Resources Inc.

I have not received, nor do I expect to receive any interest, directly or indirectly, in the properties or securities of Freeport Resources Inc. or any affiliate. I am independent of Freeport Resources Inc. in accordance with the application of Section 1.5 of National Instrument 43-101. I consent to use of this report by the company in submissions for any Regulatory requirements and development opportunities. I am not aware of any material fact or material change which is not reflected in this report. I have read National Instrument 43-101, Form 43-101F1 and this report has been prepared in compliance with NI 43-101 and Form 43-101F1.

Dated in Victoria, B.C., June 17<sup>th</sup>, 2005.



Z.D.Hora, M.Sc., P.Geo.



APPENDIX A  
ITEMIZED COST STATEMENT

**2005 Q PROJECT EXPLORATION COST SUMMARY**

Q Claims (Eaglet Property), submitted June, 2005

<b>1. Consulting Services:</b>			
Dan Hora, Geologist			
	4 days @ \$500	\$ 2,000	
Geotex Consultants Ltd. -- Peter Read, Structural Geologist		\$ 2,000	
	60 diamond drill hole logs		
Gyslaine Gauthier, Geologist			
	73 hrs. @ \$24, enter logs into database,	\$ 1,752	
	maps & copies	\$ 272	
		<b>\$ 6,024</b>	<b>\$ 6,024</b>
<b>2. Sample processing</b>			
<b>Miner</b>	various assays	\$ 3,147	
<b>ACME</b>	various assays	\$ 448	
<b>Chemex</b>	34 ICP Pt, Pd, Au	\$ 72	
<b>Van Pet</b>	10 thin sections, offcuts, k-spar stains	\$ 160	
		<b>\$ 3,826</b>	<b>\$ 3,826</b>
	<b>SUB-TOTAL</b>		<b>\$ 9,850</b>
	<b>TOTAL EXPENDITURES</b>		<b>\$ 9,850</b>
	<b>PAC account (&lt;30% total)</b>		2,950
			<b>\$ 12,800</b>

APPENDIX B

**Diamond Drill Hole Pulp & Core Samples  
Collected from Site for Re-Assay**

**Diamond Drill Hole Pulp Samples  
Collected from Site for Re-Assay**

(Total: 188 Samples, 14 Drill Holes. See Appendix C for assay data)

<b>S92/83</b>		<b>S93/83</b>		<b>S94/83</b>	
<u>Sample No.</u>	<u>Depth (feet)</u>	<u>Sample No.</u>	<u>Depth (feet)</u>	<u>Sample No.</u>	<u>Depth (feet)</u>
18501	A 234-244	18509	A 34-44	18526	A 94-104
18502	B 244-254	18511	B 279-289	18527	B 292-302
18503	C 254-264	18514	C 322-332	18528	C 302-312
18504	D 264-274	18515	D 332-342	18529	D 412-422
18505	E 274-284	18516	E 353-363	18530	E 422-432
18506	F 284-294	18517	F 363-373	18531	F 432-442
18507	G 294-304	18518	G 373-383	18532	G 442-452
		18519	H 383-393	18533	H 462-472
		18520	I 393-403	18534	I 472-482
		18521	J 403-413	18535	J 482-492
		18539	N 413-423	18536	K 452-462
		18522	K 452-462		
		18523	L 473-483		
		18524	M 483-493		
<b>S95/83</b>		<b>S96/83</b>		<b>S97/83</b>	
<u>Sample No.</u>	<u>Depth (feet)</u>	<u>Sample No.</u>	<u>Depth (feet)</u>	<u>Sample No.</u>	<u>Depth (feet)</u>
18537	A 267-277	18556	A0 344-354	18561	A 186-196
18538	B 307-317	18557	A 354-364	18562	B 210-220
18540	D 408-418	18558	B 364-374	18563	C 220-230
18541	E 418-428	18559	C 426-436	18564	D 230-240
18542	F 428-438			18565	E 255-265
18543	G 438-448			18566	F 265-275
18544	H 448-458			18567	G 307-317
18545	I 458-468			18568	H 360-370
18546	J 468-478			18569	I 370-380
18547	K 478-488			18570	J 380-390
18548	L 488-498			18571	K 400-410
18549	M 498-508				
18550	N 508-518				
18551	O 518-528				
18552	P 528-538				
18553	Q 538-548				
18554	R 548-558				
-	S missing				
18555	T 586.5-596.5				

(See Appendix C for assay data)

<b>S98/83</b>		<b>S99/83</b>		<b>S100/83</b>	
<u>Sample No.</u>	<u>Depth (feet)</u>	<u>Sample No.</u>	<u>Depth (feet)</u>	<u>Sample No.</u>	<u>Depth (feet)</u>
18572	A 27-37	18591	A 200.5-210.5	18804	A 51.5- 61.5
18573	B 37-47	18592	B 253-263	18805	B 107-117
18574	C 113.5-123.5	18593	C 275-285	18807	C 266-276
18575	D 123.5-133.5	18594	D 336-346	18808	D 286-296
18576	E 133.5-143.5	18595	E 407-417	18809	E 296-306
18577	F 143.5-153.5	18596	F 417- 427	18810	F 321-331
18578	G 166-176	18597	G 442-452		
18579	H 216-226	18599	H 487-497		
18580	I 265-275	18600	I 497-507		
18581	J 275-285	18801	J 507-517		
18582	K 285-295	18802	K 517-527		
18583	L 357-367	18803	L 527-537		
18584	M 367-377				
18585	N 377-387				
18586	O 411-421				
18587	P 421-431				
18588	Q 541-551				
18589	R 587-597				
18590	S 642-652				

(See Appendix C for assay data)

<b>S101/83</b>			<b>S102/83</b>			<b>S103/83</b>		
<u>Sample No.</u>	<u>Depth (feet)</u>		<u>Sample No.</u>	<u>Depth (feet)</u>		<u>Sample No.</u>	<u>Depth (feet)</u>	
18812	A	120.5-130.5	18828	A	250-260	18855	A	77-87
18813	B	137-147	18829	B	260-270	18856	B	87-97
18814	C	170-180	18830	C	277-287	18857	C	97-107
18815	D	180-190	18831	D	307-317	18858	D	107-117
18816	E	207-217	18832	E	317-327	18859	E	117-127
18817	F	227-237	18833	F	327-337	18860	F	137-147
18818	G	269-279	18834	G	337-347	18861	G	147-157
18819	H	279-289	18835	H	347-357	18863	H	195-205
18820	I	289-299	18836	I	428-438	18864	I	205-215
18821	J	380-390	18837	J	452-462	18865	J	215-225
18822	K	412-422	18838	K	462-472	18866	K	225-235
18823	L	436-446	18839	L	472-482	18867	L	296-306
18824	M	446-456	18840	M	536-546	18868	M	317-327
18825	N	456-466	18841	N	779-789	18869	N	327-337
18826	O	466-476	18842	O	920-930	18870	O	337-347
18827	P	487-497	18843	P	930-940	18871	P	347-357
			18844	Q	975.5-985.5	18872	Q	367-377
			18845	R	1067-1077	18873	R	377-387
			18846	S	1077-1087	18874	S	472-482
			18847	T	1102-1112	18875	T	509-519
			18848	U	1117-1127	18876	U	519-529
			18849	V	1 146-1156	18877	V	555-565
			18850	W	1156-1166	18878	W	565-575
			18851	X	1267-1277	18879	X	575-585
			18852	Y	1277-1287	18880	Y	585-595
			18853	Z	1424.5-	18881	Z	595-605
			1434.5			18882	A1	605-615
			18854	A1	1606-1616	18883	B1	615-625
<b>S104/83</b>			<b>S59/82:</b>					
<u>Sample No.</u>	<u>Depth (feet)</u>		<u>Sample No.</u>	<u>Depth (feet)</u>				
18886	A	97-107	S59/82	A	183-193			
18887	B	127-137						
18892	C	268-278						
18893	D	297-307						
18894	E	307-317						
18895	F	399-409						
18896	G	409-419						
18897	H	426-436						
18898	I	436-446						
18899	J	446-456						
18900	K	456-466						
18904	L	-----						



**Diamond Drill Core Samples  
Collected from Site for Re-Assay**

(Total: 48 Samples, 6 Drill Holes. See Appendix C for assay data)

Hole & Depth	Sample	Comments/ General Description	Tests	Assays
S53-880'	110344	greenish colour from chlorite? Up to 60 @ F10	Thin Section	REE
S110-147'	110322	shear zone		
S110-115'	110323	Up to 40-60 @ F10		Acme REE ICP
S110-323'	110325			
S110-287'	110326	Over 50 @ F10	Thin Section	REE'S
S(110?)-179'	110327	Up to 50-60 @ F10		Acme 35 ICP/Ce
S111-90'	110349	1284ppm Mo, 9308 ppm Sr, 492 ppm Zr @ 95'		Acme 35 ICP
S111-114'	110312			Acme 35 ICP
S111-123.5'	110313	Up to 50-60 @ F10		Acme REE ICP
S111-146'	110314			
S111-180.5'	110315	Up to 50-60 @ F10		Acme REE ICP
S111-201'	110316			Acme REE ICP
S111-301'	110324			
S111-410'	110317			
S111-433'	110318			
S111-449.4'	110319			
S111-450'	110320	anomalous Th, over 100 @ F10	Thin Section	Acme REE ICP
S112-95'	110301			
S112-120'	110302			
S112-117.5'	110303	Up to 50 @ F10		Acme REE ICP
S112-134'	110304			Acme 35 ICP/Ce
S112-136.5'	110305	Up to 40 @ F10, purple fluorite in gneiss		Acme 35 ICP/Ce
S112-164'	110306			Acme 35 ICP
S112-164.1'	110307	Contact zone, friable. Up to 50 @ F10		Acme 35 ICP/Ce
S112-208'	110308			Acme 35 ICP
S112-209'	110309			
S112-237.5'	110310			
S112-326.9'	110311			
S112-33.5'	110321	Up to 50 @ F10		Acme 35 ICP/Ce
S112-372'	110328	Box 15: High grade CaF2		
S112-145.38m.	110329	Biotite/mica. Box 19, Up to 60 @ F10, varies	Thin Section	
S113-57'	110330			Acme 35 ICP
S113-67'	110331	Rhodochrosite	Thin Section	
S113-91'	110332			
S113-122.5'	110333			
S113-180'	110334	Up to 40 @ F10		
S113-238'	110335	Hornfels & Biotite	Thin Section	
S113-298'	110336			
S113-313'	110337			
S113-328'	110338			
S113-399'	110339			
S113-400'	110340			
S113-425'	110341	Mo next to rich CaF2 zone. No Scint (30+/- @ F10)	Thin Section	
S113-417'	110342			

(See Appendix C for assay data)

Hole & Depth	Sample	Comments/ General Description	Tests	Assays
S114-142.5'	110343			
S114-344'	110345	Apatite/ Mo	Thin Section	
S114-365'	110346	Pyrite	Thin Section	
S114-367.5'	110347	Apatite/CaF <sub>2</sub> /Mo this whole sequence	Thin Section	
S114-386.5'	110348	Mo		

APPENDIX C  
**ASSAY DATA**

## ASSAY TESTS & EXPENSES

# of Samples (Lab & Certificate #)	\$/Sample	Total Cost
<b>ASSAYERS CANADA – MINEN</b>		
2 (08-80, 111-85) (Minen 6V-0706-RJ1)	\$20.40	\$40.80
2 (08-80, 111-85 92') (Minen 6V-0706)	\$28.00	\$56.00
10 Au (Minen 6V-1051-PG1)	\$9.50	\$95.00
10 whole rock/RRE-La (Minen)	\$36.00	\$360.00
188 31 ICP (Minen 6V-1051-PJ1-8)	\$6.20	\$1,165.60
188 F (Minen 5V-0153-PG)	\$7.56	\$1,421.28
1 31 ICP (Minen 5V-1053-PJ)	\$7.56	\$7.56
		<b>\$3,146.24</b>
<b>ACME</b>		
11 RRE ICP (Acme)	\$26.00	\$310.60
6 Core Sample Preparation	\$4.10	
10 Multi-Acid Digestion ICP (Acme)	\$8.30	\$137.00
5 Additional CE Multi-Acid Digestion	\$2.60	
10 Core Sample Preparation	\$4.10	
		<b>\$447.60</b>
<b>CHEMEX</b>		
1 ICP-34, Pt, Pd, Au (Chemex VA03021110)	\$72.21	\$72.21
<b>VAN PET</b>		
10 thin sections & k-feldspar stain (Van Pet)		<b>\$159.50</b>
		<b>\$3,825.55</b>



**Assayers Canada**  
8282 Sherbrooke St.  
Vancouver, B.C.  
V5X 4R6  
Tel: (604) 327-3436  
Fax: (604) 327-3423

Quality Assaying for over 25 Years

**Geochemical Analysis Certificate**

5V-0153-PG1

Company: **Freeport Resources**  
Project:  
Attn: **Brenda Clark**

Apr-02-05

We *hereby certify* the following geochemical analysis of 24 pulp samples submitted Mar-14-05

Sample Name	F %	CaF2 %
18501	1.32	2.70
18502	1.90	3.89
18503	1.17	2.41
18504	0.36	0.74
18505	0.66	1.35
18506	0.85	1.75
18507	1.99	4.09
18509	0.84	1.73
18511	1.44	2.95
18514	1.01	2.07
18515	0.99	2.03
18516	1.51	3.10
18517	1.71	3.52
18518	2.34	4.81
18519	1.43	2.94
18520	0.97	1.99
18521	0.70	1.44
18539	1.43	2.93
18522	0.77	1.57
18523	1.07	2.19
18524	0.91	1.88
18526	0.74	1.52
18527	0.94	1.93
18528	0.49	1.00
*DUP 18501	1.34	2.74
*DUP 18514	0.99	2.03
*DUP 18523	0.93	1.90
*STSD-1	0.09	0.18
*BLANK	<0.01	<0.02

Certified by \_\_\_\_\_



**Assayers Canada**  
8282 Sherbrooke St.  
Vancouver, B.C.  
V5X 4R6  
Tel: (604) 327-3436  
Fax: (604) 327-3423

Quality Assaying for over 25 Years

***Geochemical Analysis Certificate***

**5V-0153-PG2**

Company: **Freeport Resources**  
Project:  
Attn: **Brenda Clark**

**Apr-02-05**

We hereby certify the following geochemical analysis of 24 pulp samples submitted Mar-14-05

<b>Sample Name</b>	<b>F %</b>	<b>CaF2 %</b>
18529	1.13	2.32
18530	0.73	1.50
18531	0.84	1.73
18532	0.75	1.53
18533	0.51	1.05
18534	0.96	1.98
18535	0.48	0.99
18536	0.34	0.71
18537	0.61	1.25
18538	0.87	1.79
18540	0.45	0.93
18541	3.26	6.70
18542	1.43	2.95
18543	1.17	2.41
18544	0.21	0.43
18545	0.44	0.90
18546	0.73	1.50
18547	0.54	1.10
18548	1.00	2.05
18549	1.06	2.17
18550	0.53	1.08
18551	1.21	2.48
18552	0.70	1.44
18553	1.31	2.68
*DUP 18529	1.39	2.85
*DUP 18538	0.91	1.87
*DUP 18549	0.94	1.93
*STSD-1	0.09	0.18
*BLANK	<0.01	<0.02

Certified by



Assayers Canada  
8282 Sherbrooke St.  
Vancouver, B.C.  
V5X 4R6  
Tel: (604) 327-3436  
Fax: (604) 327-3423

Quality Assaying for over 25 Years

**Geochemical Analysis Certificate**

5V-0153-PG3

Company: **Freeport Resources**  
Project:  
Attn: **Brenda Clark**

Apr-02-05

We hereby certify the following geochemical analysis of 24 pulp samples submitted Mar-14-05

Sample Name	F %	CaF2 %
18554	0.76	1.57
18554 M	0.12	0.25
18555	1.34	2.75
18556	1.10	2.25
18557	0.85	1.75
18558	1.28	2.62
18559	1.03	2.12
18561	1.16	2.38
18562	0.75	1.54
18563	0.61	1.25
18564	0.92	1.89
18565	0.56	1.14
18566	1.18	2.43
18567	0.45	0.93
18568	0.69	1.42
18569	0.40	0.82
18570	0.51	1.05
18571	0.26	0.54
18572	0.90	1.85
18573	0.31	0.65
18574	1.18	2.43
18575	0.92	1.90
18576	0.80	1.64
18577	0.26	0.54
*DUP 18554	0.89	1.83
*DUP 18563	0.75	1.55
*DUP 18573	0.47	0.96
*STSD-1	0.12	0.31
*BLANK	<0.01	<0.02

Certified by \_\_\_\_\_



**Assayers Canada**  
8282 Sherbrooke St.  
Vancouver, B.C.  
V5X 4R6  
Tel: (604) 327-3436  
Fax: (604) 327-3423

Quality Assaying for over 25 Years

***Geochemical Analysis Certificate***

**5V-0153-PG4**

Company: **Freeport Resources**  
Project:  
Attn: **Brenda Clark**

**Apr-02-05**

We hereby certify the following geochemical analysis of 24 pulp samples submitted Mar-14-05

<b>Sample Name</b>	<b>F %</b>	<b>CaF2 %</b>
18578	0.29	0.60
18579	0.91	1.87
18580	0.78	1.60
18581	0.72	1.48
18582	0.53	1.09
18583	0.31	0.63
18584	0.85	1.74
18585	1.63	3.36
18586	1.74	3.58
18587	0.26	0.53
18588	0.55	1.13
18589	1.37	2.82
18590	1.84	3.77
18591	0.51	1.04
18592	1.17	2.40
18593	0.28	0.57
18594	0.99	2.03
18595	0.37	0.75
18596	0.54	1.11
18597	0.57	1.17
18599	0.86	1.77
18600	0.70	1.45
18801	1.37	2.81
18802	N/S	N/S
*DUP 18578	0.25	0.51
*DUP 18587	0.25	0.51
*DUP 18597	0.66	1.36
*STSD-1	0.10	0.20
*BLANK	<0.01	<0.02

Certified by \_\_\_\_\_





Assayers Canada  
8282 Sherbrooke St.  
Vancouver, B.C.  
V5X 4R6  
Tel: (604) 327-3436  
Fax: (604) 327-3423

Quality Assaying for over 25 Years

**Geochemical Analysis Certificate**

5V-0153-PG5

Company: **Freeport Resources**  
Project:  
Attn: **Brenda Clark**

Apr-02-05

We hereby certify the following geochemical analysis of 24 pulp samples submitted Mar-14-05

Sample Name	F %	CaF2 %
18803	0.82	1.69
18804	0.58	1.19
18805	0.74	1.52
18807	1.99	4.02
18808	0.48	0.99
18809	0.80	1.63
18810	0.88	1.81
18812	1.81	3.71
18813	1.49	3.05
18814	2.13	4.38
18815	0.47	0.96
18816	1.54	3.16
18817	0.42	0.87
18818	0.40	0.83
18819	0.89	1.83
18820	1.02	2.09
18821	1.21	2.48
18822	1.29	2.66
18823	1.19	2.45
18824	1.42	2.92
18825	1.62	3.33
18826	1.05	2.16
18827	0.88	1.80
18828	1.28	2.63
*DUP 18803	0.84	1.72
*DUP 18814	1.98	4.07
*DUP 18824	1.25	2.57
*STSD-1	0.11	0.23
•BLANK	<0.01	<0.02

Certified by \_\_\_\_\_



Assayers Canada  
8282 Sherbrooke St.  
Vancouver, B.C.  
V5X 4R6  
Tel: (604) 327-3436  
Fax: (604) 327-3423

Quality Assaying for over 25 Years

**Geochemical Analysis Certificate**

5V-0153-PG6

Company: **Freeport Resources**  
Project:  
Attn: **Brenda Clark**

Apr-02-05

We hereby certify the following geochemical analysis of 24 pulp samples submitted Mar-14-05

Sample Name	F %	CaF2 %
18829	0.42	0.86
18830	0.77	1.59
18831	0.42	0.86
18832	0.59	1.20
18833	2.12	4.36
18834	0.61	1.25
18835	0.73	1.50
18836	1.49	3.05
18837	1.76	3.61
18838	1.53	3.15
18839	1.86	3.83
18840	1.32	2.71
18841	1.38	2.83
18842	0.27	0.55
18843	0.47	0.96
18844	1.72	3.53
18845	0.72	1.47
18846	0.35	0.72
18847	0.47	0.97
18848	0.68	1.41
18849	0.90	1.85
18850	0.78	1.59
18851	2.13	4.38
18852	1.00	2.06
*DUP 18829	0.52	1.07
*DUP 18838	1.44	2.97
*DUP 18848	0.64	1.31
*STSD-1	0.10	0.21
•BLANK	<0.01	<0.02

Certified by \_\_\_\_\_



Assayers Canada  
8282 Sherbrooke St.  
Vancouver, B.C.  
V5X 4R6  
Tel: (604) 327-3436  
Fax: (604) 327-3423

Quality Assaying for over 25 Years

**Geochemical Analysis Certificate**

**5V-0153-PG7**

Company: **Freeport Resources**  
Project:  
Attn: **Brenda Clark**

Apr-02-05

We hereby certify the following geochemical analysis of 24 pulp samples submitted Mar-14-05

Sample Name	F %	CaF2 %
18853	0.98	2.02
18854	1.86	3.82
18855	1.21	2.49
18856	0.84	1.72
18857	0.86	1.78
18858	0.99	2.04
18859	1.15	2.37
18860	1.64	3.37
18861	0.80	1.65
18863	1.70	3.49
18864	0.93	1.90
18865	1.76	3.62
18866	0.62	1.28
18867	0.77	1.59
18868	0.60	1.23
18869	1.02	2.10
18870	1.07	2.20
18871	0.62	1.26
18872	1.90	3.90
18873	0.71	1.45
18874	0.36	0.73
18875	0.13	0.27
18876	0.54	1.11
18877	0.26	0.54
*DUP 18853	0.85	1.74
*DUP 18863	1.49	3.05
*DUP 18873	0.69	1.42
*STSD-1	0.12	0.25
•BLANK	<0.01	<0.02

Certified by \_\_\_\_\_



Assayers Canada  
8282 Sherbrooke St.  
Vancouver, B.C.  
V5X 4R6  
Tel: (604) 327-3436  
Fax: (604) 327-3423

Quality Assaying for over 25 Years

***Geochemical Analysis Certificate***

**5V-0153-PG8**

Company: **Freeport Resources**  
Project:  
Attn: **Brenda Clark**

**Apr-02-05**

We hereby certify the following geochemical analysis of 19 pulp samples submitted Mar-14-05

<b>Sample Name</b>	<b>F %</b>	<b>CaF2 %</b>
18878	2.98	6.12
18879	1.91	3.93
18880	0.89	1.83
18881	2.81	5.77
18882	1.26	2.59
18883	2.93	6.02
18886	0.72	1.46
18887	0.55	1.13
18892	0.69	1.42
18893	1.31	2.69
18894	2.47	5.08
18895	1.21	2.48
18896	1.09	2.24
18897	1.47	3.02
18898	2.25	4.62
18899	2.23	4.58
18900	0.79	1.61
18904	2.10	4.31
S59/82 183-193	6.20	12.74
*DUP 18878	2.65	5.44
*DUP 18893	1.40	2.87
*STSD-1	0.11	0.23
*BLANK	<0.01	<0.02

Certified by \_\_\_\_\_

**Freeport Resources**

Attention: Brenda Clark

Project:

Sample: pulp

**Assayers Canada**

8282 Sherbrooke St., Vancouver, B.C., V5X 4R6

Tel: (604) 327-3436 Fax: (604) 327-3423

Report No : 5V0153 PJ

Date : Apr-02-05

**MULTI-ELEMENT ICP ANALYSIS**

Aqua Regia Digestion

Sample Number	Ag ppm	Al %	As ppm	Ba ppm	Be ppm	Bi ppm	Ca %	Cd ppm	Co ppm	Cr ppm	Cu ppm	Fe %	K %	Mg %	Mn ppm	Mo ppm	Na %	Ni ppm	P ppm	Pb ppm	Sb ppm	Sc ppm	Sn ppm	Sr ppm	Ti %	V ppm	W ppm	Y ppm	Zn ppm	Zr ppm
S59/82 183-193	8.7	0.85	7	568	<0.5	<5	13.61	>100	2	47	26	1.69	0.65	0.04	1091	3	0.23	24	76	4955	15	<1	<10	4686	<0.01	7	97	20	6435	6

A .5 gm sample is digested with 5 ml 3:1 HCl/HNO3  
at 95c for 2 hours and diluted to 25ml with D.I.H2O.

**Freeport Resources**  
Attention: BRENDA CLARK  
Project: HUTTON  
Sample:

**Assayers Canada**  
8282 Sherbrooke St., Vancouver, B.C., V5X 4R6  
Tel: (604) 327-3436 Fax: (604) 327-3423

Report No : 6V1051 PJ  
Date : Dec-20-96

**ICP Report**  
Aqua Regia Digestion

Sample Number	Ag ppm	Al %	As ppm	Ba ppm	Be ppm	Bi ppm	Ca %	Cd ppm	Co ppm	Cr ppm	Cu ppm	Fe %	K %	Mg %	Mn ppm	Mo ppm	Na %	Ni ppm	P ppm	Pb ppm	Sb ppm	Sn ppm	Sr ppm	Ti %	V ppm	W ppm	Zn ppm
DDH S92-501	0.7	0.69	<5	148	<0.5	<5	2.96	<1	2	5	10	1.00	0.27	0.03	323	65	0.27	6	119	19	7	<10	2553	<0.01	15	<10	28
DDH S92-502	1.5	1.03	36	741	<0.5	<5	11.30	<1	3	6	9	0.65	0.59	0.10	1089	31	0.17	9	1173	47	12	<10	>10000	<0.01	16	<10	26
DDH S92-503	1.0	0.97	<5	394	<0.5	<5	4.68	<1	2	5	9	0.70	0.34	0.08	231	5	0.32	6	445	21	8	<10	8345	<0.01	17	<10	27
DDH S92-504	0.8	0.93	<5	324	<0.5	<5	4.20	<1	2	6	9	1.19	0.25	0.09	322	5	0.35	8	414	17	7	<10	6652	<0.01	44	<10	44
DDH S92-505	0.9	1.18	<5	206	<0.5	<5	3.97	<1	2	6	11	1.03	0.13	0.06	332	7	0.30	6	281	22	9	<10	5066	<0.01	22	<10	16
DDH S92-506	0.8	0.91	<5	339	<0.5	<5	4.18	<1	2	6	8	0.89	0.10	0.05	386	5	0.30	6	168	14	6	<10	6299	<0.01	24	<10	27
DDH S92-507	0.9	1.02	<5	175	<0.5	<5	5.11	<1	2	7	14	0.91	0.26	0.04	368	5	0.25	7	603	22	8	<10	4284	<0.01	23	<10	21
DDH S93-509	1.1	1.49	18	25	0.6	<5	4.33	<1	3	6	10	1.44	0.13	0.07	583	10	<0.01	10	156	76	9	<10	279	<0.01	7	<10	63
DDH S93-511	1.9	1.85	83	385	0.9	<5	6.69	<1	3	5	9	0.94	0.54	0.16	975	8	0.02	9	598	57	15	<10	7348	<0.01	22	<10	68
DDH S93-514	1.3	1.70	67	580	<0.5	<5	6.34	<1	3	5	7	0.79	0.68	0.20	975	25	0.06	8	1032	33	12	<10	>10000	<0.01	14	<10	39
DDH S93-515	0.9	1.36	<5	409	<0.5	<5	3.76	<1	4	5	11	0.95	0.28	0.06	480	153	0.11	7	219	25	9	<10	6779	<0.01	24	<10	27
DDH S93-516	1.4	2.08	53	923	<0.5	<5	8.69	<1	2	5	5	0.72	0.58	0.07	678	36	0.41	7	672	46	16	<10	>10000	<0.01	23	<10	23
DDH S93-517	1.6	1.60	44	884	<0.5	<5	9.29	<1	2	5	5	0.41	0.67	0.03	1014	9	0.53	7	348	52	15	<10	>10000	<0.01	9	<10	8
DDH S93-518	1.7	1.34	48	1145	<0.5	5	11.69	<1	2	6	6	0.42	0.33	0.04	1686	7	0.51	10	1033	56	13	<10	>10000	<0.01	11	<10	9
DDH S93-519	1.6	2.72	89	889	0.6	<5	11.14	<1	3	7	8	0.89	0.24	0.11	899	9	0.08	9	2233	52	17	<10	>10000	<0.01	37	<10	29
DDH S93-520	1.6	1.86	76	1182	<0.5	5	11.45	<1	2	6	6	0.44	0.29	0.04	1426	7	0.12	9	1577	54	15	<10	>10000	<0.01	18	<10	11
DDH S93-521	6.5	2.82	77	476	0.6	12	10.61	<1	2	6	8	0.64	0.07	0.07	1069	52	0.04	9	634	314	17	<10	8878	<0.01	21	<10	16
DDH S93-522	1.2	1.03	42	236	0.6	<5	6.13	<1	4	8	14	1.15	0.68	0.19	850	74	0.25	10	161	27	8	<10	2711	<0.01	29	<10	29
DDH S93-523	1.4	0.84	<5	797	<0.5	<5	5.17	<1	2	6	10	0.73	0.26	0.09	608	7	0.42	8	156	31	6	<10	>10000	<0.01	17	<10	13
DDH S93-524	0.8	0.93	<5	203	<0.5	<5	3.91	<1	2	7	11	0.88	0.30	0.15	485	9	0.25	7	145	16	6	<10	2913	<0.01	25	<10	21
DDH S93-539	16.5	2.08	40	493	<0.5	26	7.33	<1	3	7	14	1.14	0.22	0.09	877	41	0.03	8	301	568	13	<10	8976	<0.01	27	<10	23
DDH S94-526	0.8	1.48	35	96	<0.5	<5	2.59	<1	2	6	10	1.06	0.28	0.18	671	6	0.01	9	135	60	7	<10	219	<0.01	8	<10	35
DDH S94-527	0.9	0.92	<5	112	<0.5	<5	3.09	<1	2	4	8	1.08	0.55	0.06	484	8	0.19	7	296	49	7	<10	2401	<0.01	11	<10	67
DDH S94-528	0.9	0.69	<5	42	<0.5	<5	2.82	<1	2	4	7	1.15	0.20	0.15	677	6	0.07	8	141	32	5	<10	723	<0.01	14	<10	48
DDH S94-529	1.5	2.06	72	705	<0.5	<5	10.06	<1	2	6	9	0.78	0.86	0.07	949	22	0.02	9	364	53	15	<10	>10000	<0.01	19	<10	79
DDH S94-530	1.0	1.57	41	247	<0.5	<5	5.34	<1	2	4	6	0.86	0.33	0.10	779	22	0.04	8	179	27	11	<10	6733	<0.01	15	<10	53
DDH S94-531	1.2	1.17	11	372	<0.5	<5	5.34	<1	2	4	8	0.78	0.33	0.06	727	10	0.03	6	93	27	9	<10	>10000	<0.01	14	<10	44
DDH S94-532	0.9	0.97	<5	220	<0.5	<5	3.53	<1	1	4	7	0.64	0.36	0.04	432	6	0.12	5	122	25	8	<10	6683	<0.01	10	<10	17
DDH S94-533	1.0	0.87	<5	135	<0.5	<5	3.89	<1	2	4	8	0.97	0.48	0.11	736	4	0.20	8	145	26	7	<10	1904	<0.01	12	<10	60
DDH S94-534	1.2	1.11	8	214	<0.5	<5	4.79	<1	2	5	8	0.84	0.50	0.07	774	5	0.16	7	333	120	10	<10	4318	<0.01	8	<10	299

A .5 gm sample is digested with 5 ml 3:1 HCl/HNO3 at 95c for 90 min and diluted to 25ml with D.I.H2O.



Freeport Resources  
 Attention: BRENDA CLARK  
 Project: HUTTON

Assayers Canada  
 8282 Sherbrooke St., Vancouver, B.C., V5X 4R6  
 Tel: (604) 327-3436 Fax: (604) 327-3423

Report No : 6V1051 PJ  
 Date : Dec-20-96

Sample:

ICP Report  
 Aqua Regia Digestion

Sample Number	Ag ppm	Al %	As ppm	Ba ppm	Be ppm	Bi ppm	Ca %	Cd ppm	Co ppm	Cr ppm	Cu ppm	Fe %	K %	Mg %	Mn ppm	Mo ppm	Na %	Ni ppm	P ppm	Pb ppm	Sb ppm	Sn ppm	Sr ppm	Ti %	V ppm	W ppm	Zn ppm
DDH S94-535	6.3	0.86	<5	202	<0.5	<5	3.05	13	2	3	7	0.96	0.34	0.04	489	4	0.13	6	49	4071	14	<10	3105	<0.01	5	<10	1322
DDH S94-536	0.7	0.74	<5	105	<0.5	<5	2.65	<1	1	5	9	0.80	0.37	0.02	403	5	0.18	6	207	54	7	<10	1081	<0.01	4	<10	22
DDH S95-537	0.8	0.95	<5	89	0.8	<5	3.25	<1	2	5	8	0.86	0.52	0.09	536	9	0.20	7	88	17	6	<10	1713	<0.01	17	<10	36
DDH S95-538	1.3	1.02	15	103	<0.5	<5	4.84	<1	2	6	7	0.83	0.63	0.10	959	5	0.21	8	125	42	10	<10	1254	<0.01	13	<10	120
DDH S95-540	0.9	0.63	25	59	1.3	<5	3.28	<1	3	10	11	1.52	0.23	0.28	1579	12	0.06	14	443	19	<5	<10	1740	<0.01	52	<10	152
DDH S95-541	1.3	1.64	31	676	<0.5	<5	6.45	<1	4	17	26	1.40	0.74	0.04	305	20	0.05	16	687	44	13	<10	>10000	<0.01	12	<10	50
DDH S95-542	1.5	1.32	53	439	<0.5	<5	8.02	<1	3	11	12	1.05	0.57	0.11	886	7	0.03	12	587	36	16	<10	>10000	<0.01	16	<10	37
DDH S95-543	1.6	1.51	64	299	1.0	<5	7.96	<1	5	13	15	1.72	0.55	0.36	1369	15	0.03	19	1016	26	10	<10	8227	0.02	48	<10	133
DDH S95-544	1.0	0.57	<5	122	<0.5	<5	3.36	<1	2	8	10	0.72	0.16	0.05	673	4	0.04	9	70	28	6	<10	3176	<0.01	5	<10	40
DDH S95-545	1.1	0.80	<5	120	<0.5	<5	3.68	<1	2	8	10	1.01	0.17	0.04	638	5	0.04	9	58	114	8	<10	2169	<0.01	5	<10	129
DDH S95-546	1.5	0.83	<5	96	<0.5	<5	3.50	<1	2	8	10	1.08	0.44	0.03	537	7	0.18	9	220	273	9	<10	1028	<0.01	4	<10	579
DDH S95-547	1.6	0.98	<5	130	<0.5	<5	4.47	<1	2	7	9	0.87	0.47	0.04	600	5	0.16	8	398	411	10	<10	1765	<0.01	5	<10	316
DDH S95-548	1.2	1.13	12	298	<0.5	<5	6.02	<1	2	6	7	0.76	0.45	0.07	948	10	0.08	8	363	59	10	<10	6176	<0.01	7	<10	32
DDH S95-549	1.8	1.94	57	261	0.5	<5	4.81	<1	3	6	7	0.88	0.74	0.12	398	7	0.08	8	268	59	13	<10	4298	<0.01	16	<10	72
DDH S95-550	1.5	2.07	72	419	0.7	<5	5.23	<1	2	5	6	0.64	0.87	0.08	618	7	0.22	7	162	46	14	<10	6523	<0.01	13	<10	58
DDH S95-551	1.7	1.54	48	176	<0.5	<5	5.65	<1	5	18	13	1.87	0.58	0.33	969	9	0.04	18	1098	39	12	<10	3660	0.02	24	<10	160
DDH S95-552	1.4	0.85	<5	152	<0.5	<5	5.10	<1	2	9	11	0.98	0.46	0.05	1029	5	0.08	10	144	45	11	<10	3212	<0.01	8	<10	28
DDH S95-553	2.0	1.03	64	296	<0.5	<5	6.45	<1	3	9	15	1.16	0.63	0.12	1078	7	0.09	11	414	40	14	<10	5098	<0.01	21	<10	55
DDH S95-554	2.5	1.55	77	384	0.8	5	6.64	<1	3	8	14	1.10	0.65	0.09	954	9	0.11	8	262	52	15	<10	8429	<0.01	20	<10	42
DDH S95-555	2.3	1.39	23	667	<0.5	<5	6.39	<1	2	7	11	0.60	0.97	0.03	726	8	0.24	7	151	212	13	<10	>10000	<0.01	6	<10	66
DDH S96-556	1.6	2.10	66	263	<0.5	<5	10.80	<1	3	9	12	1.10	0.64	0.06	1177	10	0.03	10	391	154	19	<10	4826	<0.01	7	<10	24
DDH S96-557	1.4	2.19	54	225	0.9	<5	6.01	<1	2	5	9	0.93	0.57	0.08	658	6	0.06	7	92	163	16	<10	3199	<0.01	12	<10	119
DDH S96-558	1.8	2.14	70	716	0.6	<5	6.26	<1	2	5	8	0.69	0.99	0.08	709	32	0.04	6	305	45	17	<10	>10000	<0.01	22	<10	44
DDH S96-559	1.7	1.77	70	670	<0.5	<5	7.88	<1	2	5	5	0.45	0.55	0.05	1494	128	0.28	7	509	118	15	<10	9110	<0.01	11	<10	13
DDH S97-561	3.3	1.21	<5	198	<0.5	<5	4.64	<1	2	7	12	1.17	0.77	0.03	669	9	0.19	7	54	849	10	<10	2182	<0.01	5	<10	341
DDH S97-562	1.9	0.93	68	157	<0.5	<5	14.82	<1	3	7	8	1.01	0.59	0.12	3165	6	0.21	14	304	81	11	<10	>10000	<0.01	9	<10	55
DDH S97-563	1.8	1.71	82	668	0.6	<5	11.19	<1	3	6	7	0.68	0.77	0.09	1110	8	0.18	8	641	56	15	<10	>10000	<0.01	12	<10	47
DDH S97-564	1.2	1.42	29	365	<0.5	<5	6.28	<1	2	6	7	0.69	0.69	0.14	555	6	0.22	6	649	41	13	<10	>10000	<0.01	14	<10	44
DDH S97-565	1.6	1.85	76	173	<0.5	<5	14.24	<1	3	6	8	0.75	0.47	0.12	1128	7	0.03	8	175	43	14	<10	>10000	<0.01	14	<10	35
DDH S97-566	1.5	1.70	65	464	<0.5	<5	9.20	<1	2	6	7	0.60	0.61	0.07	1136	7	0.10	8	212	43	14	<10	>10000	<0.01	11	<10	26

A .5 gm sample is digested with 5 ml 3:1 HCl/HNO3 at 95c for 90 min and diluted to 25ml with D.I.H2O.



Freeport Resources

Attention: BRENDA CLARK

Project: HUTTON

Sample:

Assayers Canada

8282 Sherbrooke St., Vancouver, B.C., V5X 4R6

Tel: (604) 327-3436 Fax: (604) 327-3423

Report No : 6V1051 PJ

Date : Dec-20-96

ICP Report  
Aqua Regia Digestion

Sample Number	Ag ppm	Al %	As ppm	Ba ppm	Be ppm	Bi ppm	Ca %	Cd ppm	Co ppm	Cr ppm	Cu ppm	Fe %	K %	Mg %	Mn ppm	Mo ppm	Na %	Ni ppm	P ppm	Pb ppm	Sb ppm	Sn ppm	Sr ppm	Ti %	V ppm	W ppm	Zn ppm
DDH S97-567	1.0	0.87	<5	159	<0.5	<5	4.58	<1	2	6	8	0.75	0.54	0.05	761	4	0.21	7	140	52	9	<10	2593	<0.01	7	<10	89
DDH S97-568	1.2	1.05	<5	163	<0.5	<5	4.24	<1	2	6	8	0.87	0.67	0.05	611	7	0.18	6	248	75	10	<10	2395	<0.01	11	<10	85
DDH S97-569	0.9	0.53	<5	95	<0.5	<5	2.44	<1	2	5	10	1.00	0.34	0.03	490	5	0.13	6	65	183	7	<10	1054	<0.01	7	<10	399
DDH S97-570	1.5	0.94	<5	144	<0.5	<5	5.17	<1	2	6	9	1.05	0.28	0.05	1134	5	0.07	8	96	194	9	<10	2168	<0.01	9	<10	134
DDH S97-571	2.0	0.86	<5	79	<0.5	<5	3.99	<1	2	5	9	1.06	0.28	0.09	777	5	0.10	7	102	169	9	<10	1220	<0.01	7	<10	298
DDH S98-572	1.4	1.26	<5	163	<0.5	<5	7.02	<1	2	8	12	1.01	0.71	0.02	1022	6	0.29	9	55	194	10	<10	662	<0.01	5	<10	43
DDH S98-573	0.7	0.60	<5	108	<0.5	<5	2.22	<1	2	6	9	0.93	0.33	0.04	395	4	0.18	6	124	18	6	<10	287	<0.01	9	<10	29
DDH S98-574	1.6	1.27	<5	407	<0.5	<5	5.41	<1	2	6	8	0.90	0.69	0.05	766	7	0.15	8	243	283	11	<10	9514	<0.01	11	<10	590
DDH S98-575	1.4	1.16	36	343	<0.5	<5	4.82	<1	2	5	9	1.14	0.71	0.08	724	5	0.12	7	263	68	12	<10	7501	<0.01	21	<10	413
DDH S98-576	1.5	1.51	50	278	0.6	<5	8.13	<1	2	5	5	0.97	0.69	0.10	1571	6	0.12	9	199	68	12	<10	6693	<0.01	24	<10	125
DDH S98-577	1.3	1.12	<5	234	<0.5	<5	4.44	<1	2	5	8	0.99	0.43	0.09	703	5	0.11	6	225	53	10	<10	5082	<0.01	11	<10	79
DDH S98-578	1.3	0.86	29	185	<0.5	<5	8.48	<1	3	7	11	1.23	0.07	0.11	1216	6	<0.01	11	234	40	8	<10	7724	<0.01	11	<10	54
DDH S98-579	1.4	1.04	30	104	<0.5	<5	6.25	<1	2	5	10	1.05	0.46	0.08	1394	5	0.13	8	114	246	10	<10	2141	<0.01	11	<10	117
DDH S98-580	1.6	0.99	50	210	<0.5	<5	9.78	<1	2	5	8	0.83	0.67	0.06	1252	7	0.27	7	401	127	11	<10	>10000	<0.01	8	<10	62
DDH S98-581	2.0	1.01	11	242	<0.5	<5	5.27	<1	3	6	11	1.49	0.56	0.06	707	8	0.23	8	110	246	9	<10	5521	<0.01	15	<10	947
DDH S98-582	1.3	0.98	8	195	<0.5	<5	3.85	<1	2	5	11	1.46	0.28	0.08	731	5	0.08	10	318	146	9	<10	4007	<0.01	14	<10	428
DDH S98-583	14.5	1.76	58	504	<0.5	25	4.51	32	2	5	10	0.78	1.02	0.04	438	6	0.20	5	579	1157	20	<10	>10000	<0.01	12	21	1763
DDH S98-584	7.3	1.81	56	457	<0.5	11	6.40	<1	2	4	7	1.04	1.11	0.09	818	6	0.10	6	315	753	17	<10	>10000	<0.01	16	<10	61
DDH S98-585	2.2	2.43	113	605	0.5	6	14.15	<1	2	5	8	0.45	1.21	0.06	1544	6	0.14	8	1031	142	26	<10	>10000	<0.01	8	<10	37
DDH S98-586	1.5	1.05	32	198	<0.5	<5	5.45	<1	2	6	13	1.18	0.73	0.04	1044	5	0.17	8	115	76	13	<10	2356	<0.01	9	<10	118
DDH S98-587	1.4	0.85	45	132	<0.5	<5	11.43	<1	3	8	24	1.25	0.28	0.11	2412	6	0.04	14	176	68	13	<10	>10000	<0.01	12	<10	83
DDH S98-588	36.6	2.60	86	285	0.7	56	7.91	11	2	4	12	0.60	1.43	0.05	768	8	0.40	5	257	3494	27	<10	4147	<0.01	12	12	703
DDH S98-589	7.0	1.77	66	366	<0.5	11	7.96	<1	2	6	14	1.03	1.58	0.04	1005	7	0.23	6	177	667	20	<10	3269	<0.01	10	<10	196
DDH S98-590	2.0	1.76	51	156	0.7	<5	5.21	<1	2	5	11	0.89	0.73	0.05	790	16	0.20	5	190	219	15	<10	2479	<0.01	10	<10	242
DDH S99-591	1.3	2.10	59	117	0.6	<5	6.82	<1	2	6	11	0.96	0.41	0.07	1038	7	0.02	9	292	49	17	<10	2107	<0.01	15	<10	46
DDH S99-592	0.9	1.33	25	71	<0.5	<5	4.24	<1	2	6	12	1.07	0.50	0.04	472	5	0.14	5	98	66	11	<10	631	<0.01	9	<10	115
DDH S99-593	0.6	0.87	<5	87	<0.5	<5	1.92	<1	1	6	11	0.73	0.40	<0.01	121	4	0.18	5	41	34	11	<10	903	<0.01	3	<10	49
DDH S99-594	0.6	1.08	18	210	<0.5	<5	3.95	<1	2	6	11	0.87	0.48	0.04	554	11	0.18	7	85	37	10	<10	3698	<0.01	15	<10	39
DDH S99-595	0.8	0.92	22	377	<0.5	<5	2.83	<1	2	5	26	1.17	0.12	0.06	568	6	0.03	5	65	36	13	<10	7277	<0.01	14	<10	55
DDH S99-596	7.0	0.98	52	496	<0.5	8	6.45	<1	2	5	14	0.75	0.27	0.03	541	5	0.12	5	60	1663	16	<10	>10000	<0.01	6	<10	75

A .5 gm sample is digested with 5 ml 3:1 HCl/HNO<sub>3</sub> at 95c for 90 min and diluted to 25ml with D.I.H<sub>2</sub>O.



**Freeport Resources**  
 Attention: BRENDA CLARK  
 Project: HUTTON  
 Sample:

**Assayers Canada**  
 8282 Sherbrooke St., Vancouver, B.C., V5X 4R6  
 Tel: (604) 327-3436 Fax: (604) 327-3423

Report No : 6V1051 PJ  
 Date : Dec-20-96

**ICP Report**  
 Aqua Regia Digestion

Sample Number	Ag ppm	Al %	As ppm	Ba ppm	Be ppm	Bi ppm	Ca %	Cd ppm	Co ppm	Cr ppm	Cu ppm	Fe %	K %	Mg %	Mn ppm	Mo ppm	Na %	Ni ppm	P ppm	Pb ppm	Sb ppm	Sn ppm	Sr ppm	Ti %	V ppm	W ppm	Zn ppm
DDH S99-597	0.7	0.90	10	153	<0.5	<5	3.71	<1	2	7	12	1.19	0.38	0.04	411	84	0.32	7	48	47	9	<10	2703	<0.01	15	<10	35
DDH S99-599	1.3	1.41	75	832	<0.5	<5	9.69	<1	2	6	10	0.51	0.88	0.04	920	103	0.29	6	1959	81	19	<10	>10000	<0.01	5	<10	19
DDH S99-600	0.6	0.76	<5	301	<0.5	<5	3.34	<1	2	6	13	1.44	0.35	0.06	265	24	0.23	7	294	17	8	<10	7217	<0.01	45	<10	39
DDH S99-801	0.8	0.93	21	203	<0.5	<5	6.44	<1	2	5	10	1.15	0.31	0.03	533	9	0.41	6	157	42	9	<10	6294	<0.01	19	<10	23
DDH S99-802	0.9	0.87	32	364	<0.5	<5	7.69	<1	3	8	12	1.41	0.45	0.11	926	28	0.20	10	565	31	9	<10	>10000	<0.01	40	<10	27
DDH S99-803	0.9	0.65	32	241	<0.5	<5	8.18	<1	5	17	25	1.98	0.42	0.32	1440	51	0.13	16	837	25	6	<10	8477	0.01	66	<10	52
DDH S100-804	0.7	0.96	22	72	<0.5	<5	2.21	<1	2	7	10	0.92	0.34	0.02	236	7	0.05	5	287	163	25	<10	700	<0.01	13	<10	62
DDH S100-805	1.0	0.53	<5	120	<0.5	<5	3.52	<1	2	9	13	1.30	0.30	0.05	784	5	0.09	10	62	166	9	<10	2090	<0.01	8	<10	492
DDH S100-807	1.0	1.30	38	382	<0.5	<5	4.73	<1	2	11	12	0.82	0.91	0.03	452	6	0.26	9	91	40	16	<10	2050	<0.01	11	<10	48
DDH S100-808	0.5	0.65	<5	61	<0.5	<5	1.73	<1	1	5	9	0.95	0.30	0.02	353	4	0.15	5	53	32	7	<10	554	<0.01	8	<10	56
DDH S100-809	0.8	1.49	36	47	0.7	<5	2.80	<1	2	6	8	0.93	0.27	0.04	492	5	0.08	5	84	43	15	<10	677	<0.01	14	<10	76
DDH S100-810	0.7	1.10	24	52	<0.5	<5	2.95	<1	1	5	10	0.78	0.41	0.02	385	4	0.24	5	83	47	11	<10	526	<0.01	5	<10	85
DDH S101-812	2.5	1.48	29	100	<0.5	<5	4.60	<1	2	16	13	1.26	0.70	0.02	593	7	0.49	13	170	357	18	<10	918	<0.01	6	<10	278
DDH S101-813	1.2	1.14	18	160	0.6	<5	3.38	<1	2	18	14	1.16	0.73	0.02	502	6	0.34	12	141	125	14	<10	1243	<0.01	9	<10	135
DDH S101-814	0.8	1.13	24	93	2.3	<5	5.21	<1	3	22	14	1.49	0.87	0.20	905	7	0.34	16	620	42	10	<10	1198	0.01	18	<10	132
DDH S101-815	5.2	0.54	<5	85	<0.5	<5	2.71	<1	3	28	21	1.83	0.36	0.02	615	7	0.12	17	89	835	8	<10	1185	<0.01	9	<10	1360
DDH S101-816	1.2	2.19	53	668	0.6	<5	4.35	<1	2	14	10	0.87	0.80	0.09	325	15	0.17	9	173	116	20	<10	>10000	<0.01	9	<10	114
DDH S101-817	0.7	0.59	<5	65	<0.5	<5	2.10	<1	2	15	12	1.06	0.30	0.02	415	13	0.14	9	72	48	7	<10	1067	<0.01	5	<10	89
DDH S101-818	0.6	0.45	<5	86	<0.5	<5	2.57	<1	2	15	11	0.92	0.27	0.02	602	7	0.14	10	211	30	6	<10	1154	<0.01	6	<10	37
DDH S101-819	0.7	0.69	<5	125	<0.5	<5	3.08	<1	1	12	8	0.78	0.39	0.03	618	4	0.16	9	74	43	9	<10	1952	<0.01	7	<10	42
DDH S101-820	0.8	0.86	26	421	0.8	<5	4.64	<1	2	13	10	1.07	0.52	0.09	1117	8	0.14	11	99	34	11	<10	7666	<0.01	34	<10	66
DDH S101-821	0.9	2.24	33	174	0.6	<5	3.61	<1	2	11	9	1.36	0.22	0.04	409	20	0.04	9	144	66	19	<10	2661	<0.01	15	<10	96
DDH S101-822	0.8	2.17	53	156	<0.5	<5	5.63	<1	2	11	8	0.81	0.30	0.03	914	7	0.08	9	241	44	18	<10	1851	<0.01	6	<10	17
DDH S101-823	0.6	1.46	41	422	<0.5	<5	4.33	<1	1	7	7	0.55	0.35	0.03	388	29	0.15	4	189	25	14	<10	4845	<0.01	12	<10	10
DDH S101-824	0.9	2.04	54	585	<0.5	<5	5.61	<1	2	11	10	0.71	0.39	0.03	248	72	0.08	8	916	38	18	<10	9417	<0.01	14	<10	11
DDH S101-825	0.6	1.25	12	315	<0.5	<5	3.93	<1	2	17	13	1.40	0.40	0.05	278	75	0.41	10	264	20	11	<10	5455	<0.01	39	<10	17
DDH S101-826	0.6	0.98	6	304	<0.5	<5	4.79	<1	3	15	37	1.58	0.37	0.15	860	76	0.11	13	173	23	8	<10	4629	<0.01	55	<10	42
DDH S101-827	0.3	1.39	<5	169	<0.5	<5	2.95	<1	4	18	13	2.14	0.24	0.13	1080	26	0.02	15	293	40	10	<10	2622	<0.01	107	<10	119
DDH S102-828	0.7	1.16	26	127	<0.5	<5	2.90	<1	1	8	6	0.76	0.51	0.04	339	4	0.23	5	73	28	11	<10	2454	<0.01	9	<10	31
DDH S102-829	1.1	0.87	25	106	<0.5	<5	2.80	<1	1	7	7	0.80	0.33	0.04	612	4	0.08	5	41	573	11	<10	1867	<0.01	8	<10	81

A .5 gm sample is digested with 5 ml 3:1 HCl/HNO3 at 95c for 90 min and diluted to 25ml with D.I.H2O.

## Assayers Canada

8282 Sherbrooke St., Vancouver, B.C., V5X 4R6

Tel: (604) 327-3436 Fax: (604) 327-3423

Report No : 6V1051 PJ

Date : Dec-20-96

### Freeport Resources

Attention: BRENDA CLARK

Project: HUTTON

Sample:

### ICP Report Aqua Regia Digestion

Sample Number	Ag ppm	Al %	As ppm	Ba ppm	Be ppm	Bi ppm	Ca %	Cd ppm	Co ppm	Cr ppm	Cu ppm	Fe %	K %	Mg %	Mn ppm	Mo ppm	Na %	Ni ppm	P ppm	Pb ppm	Sb ppm	Sn ppm	Sr ppm	Ti %	V ppm	W ppm	Zn ppm
DDH S102-830	1.5	1.00	6	19	<0.5	<5	1.86	<1	2	16	11	1.08	0.20	0.02	281	5	0.02	10	74	813	8	<10	145	<0.01	4	<10	412
DDH S102-831	5.8	0.47	<5	29	<0.5	<5	2.10	<1	2	14	10	1.20	0.18	0.09	672	5	0.03	11	213	3134	10	<10	580	<0.01	10	<10	158
DDH S102-832	0.8	0.46	<5	57	<0.5	<5	1.93	<1	2	13	11	0.98	0.29	0.03	416	10	0.12	10	182	316	7	<10	1044	<0.01	4	<10	114
DDH S102-833	0.9	0.97	24	98	<0.5	<5	3.40	<1	2	14	12	1.24	0.75	0.08	458	5	0.24	10	88	51	11	<10	633	<0.01	12	<10	53
DDH S102-834	0.5	0.44	<5	41	<0.5	<5	1.33	<1	1	11	8	0.83	0.25	0.04	297	4	0.15	7	33	20	6	<10	444	<0.01	8	<10	17
DDH S102-835	1.0	0.51	7	63	<0.5	<5	2.08	<1	2	13	11	1.28	0.31	0.06	479	5	0.15	10	43	242	6	<10	668	<0.01	13	<10	72
DDH S102-836	0.6	0.78	20	64	2.6	<5	4.62	<1	7	16	14	1.89	0.65	0.38	2017	8	0.14	19	675	18	6	<10	375	<0.01	48	<10	160
DDH S102-837	0.8	1.00	28	134	1.4	<5	4.33	<1	3	13	9	1.41	0.67	0.19	985	6	0.27	13	129	21	11	<10	2443	<0.01	30	<10	87
DDH S102-838	0.9	0.89	30	121	<0.5	<5	4.68	<1	2	11	8	0.97	0.73	0.08	883	5	0.20	11	1078	24	10	<10	2032	<0.01	6	<10	35
DDH S102-839	0.6	1.03	23	72	2.6	<5	4.42	<1	3	11	8	1.60	0.76	0.30	1600	6	0.20	15	298	20	8	<10	1311	<0.01	31	<10	132
DDH S102-840	1.0	0.89	26	111	<0.5	<5	4.47	<1	2	11	9	0.98	0.57	0.04	765	4	0.26	9	59	226	11	<10	850	<0.01	6	<10	128
DDH S102-841	1.0	1.22	29	134	0.9	<5	3.68	<1	2	10	7	1.04	0.71	0.06	465	5	0.28	8	39	45	15	<10	2012	<0.01	12	<10	78
DDH S102-842	1.0	1.23	37	322	0.7	<5	6.05	<1	2	14	10	0.87	0.65	0.06	710	20	0.26	10	142	76	12	<10	8467	<0.01	14	<10	50
DDH S102-843	1.1	0.58	20	75	<0.5	<5	7.43	<1	2	13	10	1.05	0.25	0.05	2099	9	0.11	14	56	171	8	<10	3686	<0.01	6	<10	121
DDH S102-844	1.1	1.16	36	316	0.7	<5	5.97	<1	2	14	10	0.86	0.61	0.06	702	18	0.24	11	148	67	11	<10	8354	<0.01	13	<10	48
DDH S102-845	1.7	0.50	12	172	<0.5	<5	4.34	<1	2	18	13	0.99	0.36	0.04	728	17	0.14	13	339	428	9	<10	4952	<0.01	5	<10	287
DDH S102-846	1.5	0.28	<5	75	<0.5	<5	2.54	<1	2	19	12	1.33	0.19	0.04	634	40	0.08	13	108	589	<5	<10	1572	<0.01	6	<10	273
DDH S102-847	1.1	0.86	8	100	<0.5	<5	2.75	<1	2	11	9	1.14	0.31	0.03	422	10	0.18	7	60	170	11	<10	995	<0.01	4	<10	231
DDH S102-848	1.4	0.55	<5	116	<0.5	<5	2.79	<1	2	14	11	1.28	0.24	0.03	600	7	0.21	11	54	446	8	<10	1818	<0.01	5	<10	82
DDH S102-849	1.0	0.78	39	184	<0.5	<5	6.53	<1	2	12	8	0.68	0.25	0.04	1000	10	0.15	10	660	55	12	<10	4434	<0.01	4	<10	34
DDH S102-850	0.7	0.61	16	126	<0.5	<5	2.94	<1	2	14	11	1.06	0.33	0.06	384	17	0.19	10	107	18	10	<10	2494	<0.01	29	<10	44
DDH S102-851	1.2	0.88	36	282	<0.5	<5	10.00	<1	3	14	10	1.23	0.47	0.08	1353	112	0.36	13	205	92	10	<10	7161	<0.01	37	<10	59
DDH S102-852	0.7	0.61	<5	186	<0.5	<5	4.92	<1	4	15	12	2.22	0.31	0.20	763	31	0.26	14	301	95	5	<10	4026	<0.01	73	<10	225
DDH S102-853	1.0	0.87	36	295	1.0	<5	5.31	<1	3	13	12	1.15	0.56	0.20	598	7	0.24	10	912	23	7	<10	6507	0.02	36	<10	39
DDH S102-854	1.1	1.16	48	1079	1.1	<5	7.48	<1	4	15	14	1.61	0.75	0.19	795	14	0.30	13	1095	33	11	<10	>10000	0.02	81	<10	58
DDH S103-855	1.1	1.71	47	166	<0.5	<5	5.15	<1	2	8	9	1.00	0.80	0.05	752	7	0.17	9	266	233	17	<10	1203	<0.01	14	<10	204
DDH S103-856	1.3	2.02	78	230	<0.5	<5	5.56	<1	2	6	7	0.68	1.03	0.04	725	6	0.19	5	112	106	21	<10	1559	<0.01	9	<10	41
DDH S103-857	1.6	0.71	9	125	<0.5	<5	2.28	<1	1	8	8	0.63	0.55	0.02	328	3	0.13	6	128	728	11	<10	672	<0.01	5	<10	228
DDH S103-858	0.9	1.08	36	118	<0.5	<5	2.90	<1	2	9	9	0.86	0.61	0.07	467	5	0.13	7	244	193	13	<10	1102	<0.01	11	<10	79
DDH S103-859	0.9	1.06	44	213	<0.5	<5	4.30	<1	3	8	8	1.09	0.82	0.12	824	7	0.14	9	284	34	11	<10	2923	0.01	20	<10	108

A .5 gm sample is digested with 5 ml 3:1 HCl/HNO3 at 95c for 90 min and diluted to 25ml with D.I.H2O.



## Assayers Canada

8282 Sherbrooke St., Vancouver, B.C., V5X 4R6  
Tel: (604) 327-3436 Fax: (604) 327-3423

Report No : 6V1051 PJ  
Date : Dec-20-96

Freeport Resources  
Attention: BRENDA CLARK  
Project: HUTTON  
Sample:

### ICP Report Aqua Regia Digestion

Sample Number	Ag ppm	Al %	As ppm	Ba ppm	Be ppm	Bi ppm	Ca %	Cd ppm	Co ppm	Cr ppm	Cu ppm	Fe %	K %	Mg %	Mn ppm	Mo ppm	Na %	Ni ppm	P ppm	Pb ppm	Sb ppm	Sn ppm	Sr ppm	Ti %	V ppm	W ppm	Zn ppm
DDH S103-860	1.1	1.30	55	199	0.7	<5	5.01	<1	2	9	7	0.94	0.79	0.05	759	6	0.19	7	630	34	18	<10	3097	<0.01	28	<10	86
DDH S103-861	1.2	2.08	70	238	1.3	<5	5.39	<1	2	7	6	0.68	0.57	0.05	752	6	0.13	6	188	45	20	<10	3378	<0.01	16	<10	37
DDH S103-863	1.3	1.04	68	400	<0.5	<5	7.02	<1	2	8	6	0.51	0.74	0.03	1203	11	0.14	7	382	219	14	<10	5727	<0.01	4	<10	70
DDH S103-864	1.0	1.03	40	414	<0.5	<5	5.19	<1	1	6	5	0.52	0.55	0.03	675	4	0.12	5	331	33	14	<10	5028	<0.01	5	<10	22
DDH S103-865	1.4	1.61	98	497	<0.5	<5	11.42	<1	2	6	5	0.52	1.16	0.04	1451	5	0.17	7	354	94	20	<10	4917	<0.01	6	<10	45
DDH S103-866	1.4	0.89	32	219	<0.5	<5	4.47	<1	2	9	8	1.25	0.55	0.08	872	5	0.09	9	94	309	9	<10	2265	<0.01	12	<10	225
DDH S103-867	1.0	0.88	36	99	<0.5	<5	2.40	<1	1	9	9	0.80	0.45	0.02	389	4	0.26	6	63	335	11	<10	629	<0.01	6	<10	246
DDH S103-868	1.2	0.61	29	54	<0.5	<5	4.86	<1	3	12	11	1.61	0.33	0.17	1481	6	0.08	13	146	230	8	<10	1675	<0.01	17	<10	448
DDH S103-869	1.2	1.13	32	92	<0.5	<5	3.55	<1	2	8	7	1.11	0.35	0.06	702	5	0.12	8	172	131	12	<10	1190	<0.01	8	<10	100
DDH S103-870	1.0	1.24	45	90	<0.5	<5	3.91	<1	2	6	8	0.84	0.44	0.02	571	5	0.16	6	170	92	16	<10	1009	<0.01	6	<10	106
DDH S103-871	0.7	1.42	41	84	<0.5	<5	3.54	<1	1	7	7	0.70	0.15	0.02	845	5	0.01	6	74	107	17	<10	1529	<0.01	7	<10	70
DDH S103-872	0.9	1.94	55	168	0.5	<5	7.09	<1	1	5	5	0.59	0.87	0.03	838	5	0.18	6	67	71	17	<10	1549	<0.01	6	<10	75
DDH S103-873	0.5	0.75	<5	53	<0.5	<5	3.36	<1	1	9	7	0.78	0.24	0.03	793	4	0.07	8	103	32	6	<10	1047	<0.01	10	<10	39
DDH S103-874	3.7	0.33	<5	35	<0.5	<5	2.10	>100	2	12	11	0.94	0.13	0.01	550	5	0.07	9	77	2361	10	<10	677	<0.01	4	39	>10000
DDH S103-875	0.9	0.31	<5	62	<0.5	<5	1.90	<1	1	7	7	0.87	0.06	0.01	396	3	0.05	5	13	238	<5	<10	1370	<0.01	3	<10	188
DDH S103-876	1.1	0.89	<5	116	<0.5	<5	3.09	<1	1	6	6	0.63	0.21	0.02	478	5	0.10	5	27	254	11	<10	1909	<0.01	3	<10	207
DDH S103-877	0.6	1.17	23	128	<0.5	<5	2.70	<1	1	8	6	0.80	0.09	0.02	471	136	<0.01	6	43	45	7	<10	2404	<0.01	21	<10	34
DDH S103-878	1.2	1.76	86	837	<0.5	<5	10.82	<1	1	5	4	0.29	1.28	0.04	998	178	0.13	6	855	67	18	<10	>10000	<0.01	6	<10	30
DDH S103-879	1.2	1.17	16	395	<0.5	<5	7.29	<1	2	5	4	0.54	0.65	0.05	981	32	0.15	6	307	53	11	<10	7781	<0.01	6	<10	30
DDH S103-880	0.7	0.66	<5	171	<0.5	<5	3.46	<1	1	6	5	0.35	0.25	0.02	516	4	0.06	4	38	19	6	<10	2371	<0.01	3	<10	45
DDH S103-881	1.4	1.19	26	708	<0.5	<5	8.87	<1	2	8	8	0.59	0.53	0.05	1226	62	0.26	8	440	47	13	<10	>10000	<0.01	11	<10	21
DDH S103-882	1.3	1.42	24	515	<0.5	<5	6.68	<1	2	8	8	0.52	0.60	0.02	620	27	0.34	7	195	48	16	<10	9903	<0.01	6	<10	15
DDH S103-883	1.0	1.80	19	349	<0.5	<5	5.21	<1	1	5	6	0.39	0.57	0.02	308	116	0.45	4	413	39	14	<10	6899	<0.01	8	<10	11
DDH S104-886	1.1	0.68	<5	52	0.8	<5	2.72	<1	2	16	10	1.27	0.13	0.06	786	6	0.04	13	75	127	6	<10	884	<0.01	14	<10	92
DDH S104-887	0.8	0.45	<5	33	<0.5	<5	1.69	<1	2	14	11	1.24	0.16	0.03	502	9	0.10	11	51	69	<5	<10	506	<0.01	5	<10	68
DDH S104-892	0.9	0.55	<5	75	<0.5	<5	2.26	<1	1	12	8	0.78	0.22	0.01	560	4	0.12	8	42	57	8	<10	1108	<0.01	3	<10	43
DDH S104-893	2.2	1.42	52	107	<0.5	<5	11.20	<1	2	12	8	0.85	0.18	0.06	1514	43	0.02	12	194	555	12	<10	6256	<0.01	6	<10	338
DDH S104-894	1.0	1.85	25	287	<0.5	<5	5.19	<1	2	10	7	0.87	0.49	0.01	374	9	0.02	8	82	69	10	<10	5487	<0.01	9	<10	51
DDH S104-895	0.6	1.25	<5	80	<0.5	<5	2.56	<1	2	18	11	1.44	0.25	0.02	259	38	0.05	14	172	30	8	<10	1367	<0.01	12	<10	52
DDH S104-896	0.8	1.69	27	341	<0.5	<5	3.48	<1	2	12	8	1.02	0.39	0.04	191	61	0.05	8	84	33	11	<10	5555	<0.01	17	<10	34

A .5 gm sample is digested with 5 ml 3:1 HCl/HNO3 at 95c for 90 min and diluted to 25ml with D.I.H2O.



**Freeport Resources**  
 Attention: BRENDA CLARK  
 Project: HUTTON  
 Sample:

**Assayers Canada**  
 8282 Sherbrooke St., Vancouver, B.C., V5X 4R6  
 Tel: (604) 327-3436 Fax: (604) 327-3423

Report No : 6V1051 PJ  
 Date : Dec-20-96

**ICP Report**  
 Aqua Regia Digestion

Sample Number	Ag ppm	Al %	As ppm	Ba ppm	Be ppm	Bi ppm	Ca %	Cd ppm	Co ppm	Cr ppm	Cu ppm	Fe %	K %	Mg %	Mn ppm	Mo ppm	Na %	Ni ppm	P ppm	Pb ppm	Sb ppm	Sn ppm	Sr ppm	Ti %	V ppm	W ppm	Zn ppm
DDH S104-897	1.2	0.90	12	306	<0.5	<5	8.55	<1	2	10	7	0.70	0.28	0.04	1219	8	0.12	9	141	53	10	<10	5884	<0.01	5	<10	4
DDH S104-898	1.2	1.66	50	307	<0.5	<5	7.44	<1	2	10	7	0.76	0.27	0.07	984	11	0.04	9	432	36	13	<10	4440	<0.01	17	<10	22
DDH S104-899	1.4	1.79	56	319	<0.5	<5	6.96	<1	2	7	6	0.67	0.23	0.05	769	7	0.03	6	522	35	12	<10	5935	<0.01	8	<10	18
DDH S104-900	0.8	0.83	<5	283	<0.5	<5	2.37	<1	2	7	7	0.80	0.16	0.04	249	52	0.06	6	59	15	8	<10	4767	<0.01	15	<10	11
DDH S104-904	0.7	1.22	17	265	<0.5	<5	5.95	<1	6	12	27	1.54	0.28	0.11	608	82	0.13	12	78	18	8	<10	4501	<0.01	73	<10	34
DDH S59-82	9.1	0.89	69	485	<0.5	8	14.18	>100	4	39	27	1.34	0.57	0.04	1084	11	0.18	26	149	5070	28	<10	5966	<0.01	7	17	5629

A .5 gm sample is digested with 5 ml 3:1 HCl/HNO3 at 95c for 90 min and diluted to 25ml with D.I.H2O.



Assayers Canada  
8282 Sherbrooke St.  
Vancouver, B.C.  
V5X 4R6  
Tel: (604) 327-3436  
Fax: (604) 327-3423

Quality Assaying for over 25 Years

**Assay Certificate**

**6V-1051-PA9**  
Page 1 of 2  
Dec-20-96

Company: **Freeport Resources**  
Project: **HUTTON**  
Attn: **BRENDA CLARK**

We hereby certify the following assay of 10 pulp samples submitted Nov-02-96 by Brenda Clark.

Sample Name	Al <sub>2</sub> O <sub>3</sub> %	Ba %	CaO %	Fe <sub>2</sub> O <sub>3</sub> %	K <sub>2</sub> O %	MgO %	MnO %	Na <sub>2</sub> O %
S92-502	12.75	0.370	17.37	1.38	5.14	0.19	0.15	3.45
S93-518	13.16	0.275	17.76	0.88	1.39	0.09	0.23	6.34
S93-519	12.69	0.240	17.57	1.65	2.54	0.20	0.12	1.61
S93-520	12.18	0.465	20.21	0.92	2.90	0.09	0.20	2.61
S102-854	10.28	0.345	11.09	3.42	3.73	0.34	0.11	3.07
S59-82	1.93	0.090	26.98	2.95	0.85	0.12	0.17	0.09
S97-562	9.74	0.080	20.14	2.01	3.72	0.22	0.44	2.66
S98-585	12.01	0.265	22.60	1.19	5.16	0.15	0.23	1.12
S103-874	14.03	0.045	2.72	1.84	4.42	0.06	0.06	4.45
S103-878	12.96	0.370	17.21	0.76	7.57	0.10	0.15	1.17

\*Sample was fused with Lithium metaborate and dissolved in dilute HNO<sub>3</sub>.

Certified by \_\_\_\_\_



Assayers Canada  
8282 Sherbrooke St.  
Vancouver, B.C.  
V5X 4R6  
Tel: (604) 327-3436  
Fax: (604) 327-3423

Quality Assaying for over 25 Years

**Assay Certificate**

6V-1051-PA9  
Page 2 of 2  
Dec-20-96

Company: **Freeport Resources**  
Project: **HUTTON**  
Attn: **BRENDA CLARK**

We hereby certify the following assay of 10 pulp samples submitted Nov-02-96 by Brenda Clark.

Sample Name	P2O5 %	SiO2 %	Sr %	TiO2 %
S92-502	0.29	44.03	2.825	0.06
S93-518	0.20	44.83	2.175	0.21
S93-519	0.67	46.78	1.895	0.11
S93-520	0.51	41.46	3.985	0.12
S102-854	0.16	54.65	2.330	0.12
S59-82	0.01	50.96	0.610	0.02
S97-562	0.01	45.60	0.960	0.08
S98-585	0.33	38.70	1.900	0.03
S103-874	0.01	66.81	0.120	0.04
S103-878	0.24	45.49	2.190	0.02

\*Sample was fused with Lithium metaborate and dissolved in dilute HNO3.

Certified by \_\_\_\_\_ 



**Assayers Canada**  
 8282 Sherbrooke St.  
 Vancouver, B.C.  
 V5X 4R6  
 Tel: (604) 327-3436  
 Fax: (604) 327-3423

Quality Assaying for over 25 Years

**Geochemical Analysis Certificate**

**6V-1051-PG1**

Company: **Freeport Resources**  
 Project: **HUTTON**  
 Attn: **BRENDA CLARK**

Dec-20-96

We hereby certify the following geochemical analysis of 20 pulp samples submitted Nov-02-96 by Brenda Clark.

Sample Name	Au-fire PPB	NB PPM	Y PPM	ZR PPM	LA PPM
S59-82	2				
S93-539	14				
S94-535	12				
S98-583	4				
S98-584	13				
S98-588	2				
S98-589	1				
S99-596	1				
S102-831	1				
S103-874	1				
S92-502		11	<10	497	23
S93-518		12	<10	309	45
S93-519		<10	<10	165	81
S93-520		<10	<10	132	118
S102-854		17	<10	296	147
S59-82		<10	28	89	382
S97-562		<10	29	521	415
S98-585		<10	20	394	228
S103-874		<10	13	514	132
S103-878		<10	23	855	200

Certified by



Assayers Canada  
8282 Sherbrooke St.  
Vancouver, B.C.  
V5X 4R6  
Tel: (604) 327-3436  
Fax: (604) 327-3423

Quality Assaying for over 25 Years

***Assay Certificate***

6V-0706-PA3

Page 1 of 2

Sep-27-96

Company: **Freeport Resources**  
Project: **HUTTON**  
Attn: **BRENDA CLARK**

We hereby certify the following assay of 2 pulp samples  
submitted Mar-07-05

Sample Name	Al <sub>2</sub> O <sub>3</sub> %	Ba %	CaO %	Fe <sub>2</sub> O <sub>3</sub> %	K <sub>2</sub> O %	MgO %	MnO %	Na <sub>2</sub> O %
08-80 END OF HOLE	4.01	0.045	6.17	18.29	0.82	0.74	0.14	0.01
111-85 92	16.29	0.075	7.32	3.11	0.26	0.36	0.07	8.59

\*Sample was fused with Lithium metaborate and dissolved in dilute HNO<sub>3</sub>.

Certified by \_\_\_\_\_





Assayers Canada  
8282 Sherbrooke St.  
Vancouver, B.C.  
V5K 4R6  
Tel: (604) 327-3436  
Fax: (604) 327-3423

Quality Assaying for over 25 Years

**Assay Certificate**

**6V-0706-PA3**  
Page 2 of 2  
Sep-27-96

Company: **Freeport Resources**  
Project: **HUTTON**  
Attn: **BRENDA CLARK**

We hereby certify the following assay of 2 pulp samples  
submitted Mar-07-05

Sample Name	P2O5 %	SiO2 %	Sr %	TiO2 %
08-80 END OF HOLE	0.01	68.83	0.025	0.17
111-85 92	0.01	62.00	0.705	0.06

\*Sample was fused with Lithium metaborate and dissolved in dilute HNO3.

Certified by \_\_\_\_\_



Quality Assaying for over 25 Years

**Geochemical Analysis Certificate**

**6V-0706-RG1**

Company: **Freeport Resources**  
Project: **HUTTON**  
Attn: **BRENDA CLARK**

**Sep-27-96**

We hereby certify the following geochemical analysis of 2 rock samples  
submitted Mar-07-05

<b>Sample Name</b>	<b>AU-FIRE PPB</b>
08-80 END OF HOLE	12
111-85 92'	1

Certified by \_\_\_\_\_



**Assayers Canada**  
8282 Sherbrooke St.  
Vancouver, B.C.  
V5X 4R6  
Tel: (604) 327-3436  
Fax: (604) 327-3423

Quality Assaying for over 25 Years

**Assay Certificate**

**6V-0706-PA2**

Company: **Freeport Resources**  
Project: **HUTTON**  
Attn: **BRENDA CLARK**

Sep-27-96

We *hereby certify* the following assay of 2 pulp samples  
submitted Mar-07-05

<b>Sample Name</b>	<b>Nb PPM</b>	<b>Y PPM</b>	<b>Zr PPM</b>
08-80 END OF HOLE	6	14	20
111-85 92	3	22	492

Certified by \_\_\_\_\_

**Freeport Resources**  
 Attention: BRENDA CLARK  
 Project: HUTTON  
 Sample:

**Assayers Canada**  
 8282 Sherbrooke St., Vancouver, B.C., V5X 4R6  
 Tel: (604) 327-3436 Fax: (604) 327-3423

Report No : 6V0706 RJ  
 Date : Sep-27-96

**ICP Report**  
 Aqua Regia Digestion

Sample Number	Ag ppm	Al %	As ppm	Ba ppm	Be ppm	Bi ppm	Ca %	Cd ppm	Co ppm	Cr ppm	Cu ppm	Fe %	K %	Mg %	Mn ppm	Mo ppm	Na %	Ni ppm	P ppm	Pb ppm	Sb ppm	Sn ppm	Sr ppm	Ti %	V ppm	W ppm	Zn ppm
08-80 END OF HOLE	<0.2	0.89	<5	82	<0.5	<5	3.80	<1	39	285	245	9.09	0.27	0.33	897	81	0.01	70	<10	<2	<5	<10	157	0.01	9	<10	10
111-85 92'	0.4	1.59	30	529	<0.5	<5	4.80	<1	3	52	2	1.38	0.09	0.20	525	1284	0.67	11	246	15	16	<10	9308	<0.01	70	<10	43

Y

A .5 gm sample is digested with 5 ml 3:1 HCl/HNO3 at 95c for 90 min and diluted to 25ml with D.I.H2O.

Signed: \_\_\_\_\_ 



## GEOCHEMICAL ANALYSIS CERTIFICATE



Freeport Resources Inc. PROJECT HUTTON File # 96-6592 Page 1  
8711 Elsmore Road, Richmond BC V7C 2A4

SAMPLE#	Mo	Cu	Pb	Zn	Ag	Ni	Co	Mn	Fe	As	U	Au	Th	Sr	Cd	Sb	Bi	V	Ca	P	La	Cr	Mg	Ba	Ti	Al	Na	K	W	Zr	Sn	Y	Nb	Be	Sc	Ce		
	ppm	ppm	ppm	ppm	ppm	ppm	ppm	ppm	%	ppm	ppm	ppm	ppm	ppm	ppm	ppm	ppm	ppm	%	%	ppm	ppm	%	ppm	%	%	%	%	ppm	ppm	ppm	ppm	ppm	ppm	ppm	ppm	ppm	ppm
B 110304	3	37	25	444	<.5	5	<2	3001	1.88	<5	60	<4	363	531	.6	<5	<5	45	6.75	.013	63	2	.09	281	.05	6.51	2.88	3.19	4	29	4	21	178	23	3	105		
B 110305	<2	6	88	174	<.5	<2	<2	1514	1.47	<5	105	<4	121	1966	<.4	<5	<5	34	5.60	.030	513	3	.12	581	.04	8.25	3.01	5.24	<4	6	3	20	121	5	3	761		
B 110307	3	8	71	196	<.5	5	<2	392	1.16	<5	45	<4	25	1113	.4	<5	<5	16	.71	<.002	32	<2	.08	309	.04	11.43	2.90	3.71	<4	6	<2	3	155	8	<1	49		
B 110321	6	7	11	69	<.5	<2	<2	1584	.97	<5	26	<4	104	1706	.5	6	<5	72	11.24	<.002	10	<2	.04	202	.03	7.03	4.86	.54	<4	6	2	22	76	5	2	22		
B 110327	3	13	78	116	<.5	4	<2	506	1.86	<5	105	<4	165	1898	<.4	<5	<5	64	1.23	.003	91	7	.14	1141	.09	8.33	3.34	4.66	<4	45	4	9	837	14	2	194		
RE B 110327	4	15	86	121	<.5	7	<2	517	1.89	<5	120	<4	166	1916	<.4	<5	<5	64	1.26	.002	92	7	.14	1153	.09	8.51	3.41	4.79	5	49	3	9	861	15	1	198		

ICP - .250 GRAM SAMPLE IS DIGESTED WITH 10ML HClO<sub>4</sub>-HNO<sub>3</sub>-HCL-HF AT 200 DEG. C TO FUMING AND IS DILUTED TO 10 ML WITH DILUTED AQUA REGIA. THIS LEACH IS PARTIAL FOR MAGNETITE, CHROMITE, BARITE, OXIDES OF AL, ZR & MN AND MASSIVE SULFIDE SAMPLES. AS, CR, SB, AU SUBJECT TO LOSS BY VOLATILIZATION DURING HClO<sub>4</sub> FUMING.

- SAMPLE TYPE: CORE Samples beginning 'RE' are Reruns and 'RRE' are Reject Reruns.

DATE RECEIVED: DEC 13 1996

DATE REPORT MAILED: *Dec 20/96*

SIGNED BY: *C. Leong* D. TOYE, C. LEONG, J. WANG; CERTIFIED B.C. ASSAYERS



SAMPLE#	Mo	Cu	Pb	Zn	Ag	Ni	Co	Mn	Fe	As	U	Au	Th	Sr	Cd	Sb	Bi	V	Ca	P	La	Cr	Mg	Ba	Ti	Al	Na	K	W	Zr	Sn	Y	Nb	Be	Sc	
	ppm	ppm	ppm	ppm	ppm	ppm	ppm	ppm	%	ppm	ppm	ppm	ppm	ppm	ppm	ppm	ppm	ppm	%	%	ppm	ppm	%	ppm	%	%	%	ppm	ppm	ppm	ppm	ppm	ppm	ppm	ppm	ppm
B 110306	2	9	21	268	<.5	3	<2	1728	1.09	<5	35	<4	115	397	<.4	<5	<5	34	7.92	.002	129	6	.05	95	.06	5.85	.43	4.02	<4	6	3	22	86	13	3	
B 110308	161	5	26	48	<.5	5	<2	589	.98	<5	45	<4	43	3809	<.4	<5	<5	35	2.15	.003	145	5	.12	593	.04	6.19	1.51	3.49	<4	8	3	13	64	8	3	
B 110312	3	5	15	59	<.5	2	<2	1223	.68	<5	25	<4	56	303	<.4	<5	<5	7	2.77	<.002	148	<2	.04	142	.07	6.42	.15	3.25	<4	31	3	14	101	15	2	
B 110330	4	7	27	27	<.5	8	<2	485	.53	<5	<10	<4	5	425	<.4	<5	<5	15	8.74	<.002	2	5	.03	74	.02	3.33	1.07	2.51	<4	8	<2	18	21	4	<1	
B 110349	1235	6	14	73	<.5	<2	<2	704	2.56	<5	30	<4	5	6684	<.4	<5	<5	176	3.46	.035	10	2	.09	916	.05	6.53	5.23	.10	<4	5	5	4	63	4	7	
RE B 110349	1196	7	14	72	<.5	3	<2	710	2.60	<5	39	<4	4	6760	.5	<5	<5	180	3.50	.034	11	<2	.09	948	.05	6.62	5.19	.10	<4	5	5	4	67	4	8	
STANDARD CT2	19	58	45	143	7.0	79	32	1228	4.45	26	22	6	40	238	16.3	20	16	128	1.12	.115	43	113	1.17	910	.31	7.17	1.60	1.73	16	45	21	11	10	2	15	

Sample type: CORE. Samples beginning 'RE' are Reruns and 'RRE' are Reject Reruns.



## WHOLE ROCK ICP-MS ANALYSIS



Freeport Resources Inc. PROJECT HUTTON File # 96-6593 Page 1

8711 Elsmore Road, Richmond BC V7C 2A4

SAMPLE#	Y ppm	La ppm	Ce ppm	Pr ppm	Nd ppm	Sm ppm	Eu ppm	Gd ppm	Tb ppm	Dy ppm	Ho ppm	Er ppm	Tm ppm	Yb ppm	Lu ppm
B 110303	26.3	124.5	212.1	14.0	14.5	2.2	1.6	2.5	1.4	2.6	1.2	10.1	<.2	3.3	1.2
B 110313	4.8	45.2	79.9	5.7	5.5	1.6	.5	1.3	1.0	<.8	<.3	4.3	<.2	.6	.3
B 110315	26.4	11.0	22.0	.5	4.9	<.5	.4	1.3	.3	<.8	.6	<.8	<.2	1.4	.3
B 110316	7.9	41.2	73.8	4.4	7.2	1.6	.6	1.5	.6	<.8	<.3	4.4	<.2	.8	.2
B 110320	44.5	105.1	243.3	16.3	31.6	10.4	3.9	6.5	2.0	4.3	1.8	14.9	<.2	5.9	1.3
B 110323	14.2	31.7	77.3	6.3	7.6	1.5	.9	1.4	.9	<.8	.7	1.5	<.2	2.7	.7
RE B 110323	13.9	33.7	79.0	6.5	8.3	1.9	.9	2.1	.9	1.3	.3	5.0	<.2	2.6	.6
STANDARD SY-3	700.4	1215.0	2126.0	223.0	670.0	110.9	19.1	103.1	22.4	110.4	32.7	50.2	7.6	61.6	11.1

.200 GRAM SAMPLE FUSED WITH 1.2 GM LIBO2 AND IS DISSOLVED AND DILUTED TO 100 ML WITH 5% HNO3. RARE EARTH ELEMENTS PRE-CONCENTRATED AND SEPARATED FROM MAJOR ELEMENTS, ANALYSED BY ICP -

- SAMPLE TYPE: P1 CORE P2 PULP Samples beginning 'RE' are Reruns and 'RRE' are Reject Reruns.

DATE RECEIVED: DEC 13 1996

DATE REPORT MAILED:

Dec 23/96

SIGNED BY.....D.TOYE, C.LEONG, J.WANG; CERTIFIED B.C. ASSAYERS



SAMPLE#	Y ppm	La ppm	Ce ppm	Pr ppm	Nd ppm	Sm ppm	Eu ppm	Gd ppm	Tb ppm	Dy ppm	Ho ppm	Er ppm	Tm ppm	Yb ppm	Lu ppm
S92-502	18.4	35.8	62.5	4.4	5.7	<.5	1.2	2.2	.2	<.8	<.3	.8	<.2	1.5	.3
S93-520	21.7	37.3	54.8	5.0	4.6	<.5	1.3	2.3	<.2	<.8	<.3	1.3	<.2	1.8	.5
S97-562	27.7	206.3	449.4	18.8	24.0	.9	3.1	3.1	.6	1.2	.4	16.5	.3	2.3	.7
S98-585	22.1	103.1	173.4	10.2	12.6	<.5	1.5	2.1	<.2	<.8	<.3	2.2	<.2	1.9	.5
S102-854	23.1	64.5	109.5	6.4	11.8	1.9	1.9	2.7	.4	1.5	.5	4.6	<.2	2.2	.5
RE S102-854	20.7	58.6	101.5	6.9	10.7	1.8	1.6	2.7	1.0	1.9	.5	4.9	<.2	1.9	.4

Sample type: PULP. Samples beginning 'RE' are Reruns and 'RRE' are Reject Reruns.





**ALS Chemex**  
**EXCELLENCE IN ANALYTICAL CHEMISTRY**  
ALS Canada Ltd.  
212 Brooksbank Avenue  
North Vancouver BC V7J 2C1 Canada  
Phone: 604 984 0221 Fax: 604 984 0218

FREEPORT RESOURCES INC.  
8711 ELSMORE ROAD  
RICHMOND BC V7C 2A4

Page #: 1  
Date: 23-Jun-2003  
Account: FREEPO

**CERTIFICATE VA03021110**

Project :  
P.O. No:  
This report is for 1 SOIL sample submitted to our lab in North Vancouver, BC, Canada on 17-Jun-2003.

The following have access to data associated with this certificate:

BRENDA CLARK

**SAMPLE PREPARATION**

ALS CODE	DESCRIPTION
WEI-21	Received Sample Weight
LOG-22	Sample login - Rcd w/o BarCode
SCR-41	Screen to -180um and save both

**ANALYTICAL PROCEDURES**

ALS CODE	DESCRIPTION	INSTRUMENT
PGM-ICP23	Pt, Pd, Au 30g FA ICP	ICP-AES
ME-ICP41	34 Element Aqua Regia ICP-AES	ICP-AES

To: **FREEPORT RESOURCES INC.**  
**ATTN: BRENDA CLARK**  
**8711 ELSMORE ROAD**  
**RICHMOND BC V7C 2A4**

This is the Final Report and supersedes any preliminary report with this certificate number. Results apply to sample as submitted. All pages of this report have been checked and approved for release.

Signature:



# ALS Chemex

**EXCELLENCE IN ANALYTICAL CHEMISTRY**

ALS Canada Ltd.  
 212 Brooksbank Avenue  
 North Vancouver BC V7J 2C1 Canada  
 Phone: 604 984 0221 Fax: 604 984 0218

: FREEPORT RESOURCES INC.  
 8711 ELSMORE ROAD  
 RICHMOND BC V7C 2A4

age #: 2 - A  
 Total # of pages : 2 (A - C)  
 Date : 23-Jun-2003  
 Account: FREEPO

## CERTIFICATE OF ANALYSIS VA03021110

Method Analyte Units	WEI-21 Recvd Wt kg	PGM-ICP23 Au ppm	PGM-ICP23 Pt ppm	PGM-ICP23 Pd ppm	ME-ICP41 Ag ppm	ME-ICP41 Al %	ME-ICP41 As ppm	ME-ICP41 B ppm	ME-ICP41 Ba ppm	ME-ICP41 Be ppm	ME-ICP41 Bi ppm	ME-ICP41 Ca %	ME-ICP41 Cd ppm	ME-ICP41 Co ppm	ME-ICP41 Cr ppm
Sample Description	LOR														
EAGLET STREAM	0.98	<0.001	0.007	<0.001	<0.2	1.03	<2	<10	80	<0.5	<2	0.33	<0.5	12	24



**ALS Chemex**  
**EXCELLENCE IN ANALYTICAL CHEMISTRY**

ALS Canada Ltd.  
 212 Brooksbank Avenue  
 North Vancouver BC V7J 2C1 Canada  
 Phone: 604 984 0221 Fax: 604 984 0218

: FREEPORT RESOURCES INC.  
 8711 ELSMORE ROAD  
 RICHMOND BC V7C 2A4

age #: 2 - B  
 Total # of pages : 2 (A - C)  
 Date : 23-Jun-2003  
 Account: FREEPO

**CERTIFICATE OF ANALYSIS      VA03021110**

Sample Description	Method	ME-ICP41	ME-ICP41	ME-ICP41	ME-ICP41	ME-ICP41	ME-ICP41	ME-ICP41	ME-ICP41	ME-ICP41	ME-ICP41	ME-ICP41	ME-ICP41	ME-ICP41	ME-ICP41	
	Analyte	Cu	Fe	Ga	Hg	K	La	Mg	Mn	Mo	Na	Ni	P	Pb	S	
Units		ppm	%	ppm	ppm	%	ppm	%	ppm	ppm	%	ppm	ppm	ppm	%	
LOR		1	0.01	10	1	0.01	10	0.01	5	1	0.01	1	10	2	0.01	
EAGLET STREAM		18	2.10	<10	1	0.16	30	0.76	303	<1	0.02	40	680	51	0.09	<2



# ALS Chemex

**EXCELLENCE IN ANALYTICAL CHEMISTRY**

ALS Canada Ltd.

212 Brooksbank Avenue

North Vancouver BC V7J 2C1 Canada

Phone: 604 984 0221 Fax: 604 984 0218

: FREEPORT RESOURCES INC.

8711 ELSMORE ROAD

RICHMOND BC V7C 2A4

Page #: 2 - C

Total # of pages : 2 (A - C)

Date : 23-Jun-2003

Account: FREEPO

## CERTIFICATE OF ANALYSIS

VA03021110

Sample Description	Method	ME-ICP41	ME-ICP41	ME-ICP41	ME-ICP41	ME-ICP41	ME-ICP41	ME-ICP41	ME-ICP41
	Analyte	Sc	Sr	Ti	Ti	U	V	W	Zn
Units		ppm	ppm	%	ppm	ppm	ppm	ppm	ppm
LOR		1	1	0.01	10	10	1	10	2
EAGLET STREAM		2	48	0.09	<10	<10	23	<10	53

APPENDIX D

**DRILL HOLE SUMMARY**

1975 Diamond Drilling  
1981-83 Diamond Drilling  
1984-85 Diamond Drilling

## Drill Hole Summary

### 1975 Diamond Drilling

Drill Hole #	Depth (M)	Intercept (from - to)	Width	Assay CaF2%	Ag Assays	DDH Log	X Section
1-75	124.7	39.3 - 41.7 79.9 - 82.9	2.4 3.0	5.2 5.0	-----	YES (unfiled)	-----
2-75	39.0	13.4 - 16.5	3.1	5.3	-----	YES (unfiled)	-----
3-75	152.44	67.2 - 67.8 112.2 - 115.2 115.4 - 117.3 149.4 - 150.1	0.6 3.0 2.0 1.5	26.3 6.0 27.0 41.0	----- -----	YES (unfiled)	----- -----
4-75	157.0	80.2 - 82.9 82.9 - 84.5	2.7 1.6	6.5 18.0	-----	YES (unfiled)	-----
5-75	116.16	89.6 - 93.3	3.7	3.5	-----	YES (unfiled)	-----
6-75	106.7	52.4 - 55.2 55.2 - 58.0 81.4 - 82.0 83.5 - 84.7 89.6 - 92.3	2.8 2.8 0.6 1.2 2.7	13.9 6.7 17.7 10.3 10.0	----- ----- ----- ----- -----	YES (unfiled)	----- ----- ----- ----- -----
7-75	106.1	42.7 - 45.7 57.6 - 60.1 86.9 - 88.6	3.0 2.5 1.7	12.0 6.0 7.1	----- ----- -----	YES (unfiled)	----- ----- -----
8-75	80.5	42.1 - 44.8	2.7	5.5	-----	YES (unfiled)	-----
9-75	30.8	19.2 - 19.5 57.0 - 60.1	0.3 3.1	6.2 7.7	----- -----	----- -----	----- -----

### 1981-1983 Diamond Drilling

(see Appendix C for assay data)

Drill Hole #	Depth (M)	Intercept (from - to)	Width	Assay CaF2%	Ag Assays	DDH Log	X Section
S1-81	187.4	36.5 - 39.6	3.0	11.69	.05	YES	YES
		94.5 - 97.5	3.0	7.79	.05	Structural	
		106.6 - 161.5	54.9	10.80	.058		
S2-81	125.9	88.3 - 97.4	9.1	9.40	.06	YES Structural	YES
S3-81	133.8	48.7 - 51.8	3.0	9.74	.09	YES	YES
		97.5 - 100.5	3.0	4.87	.09		
		118.8 - 121.8	3.0	9.93	.10		
		131.0 - 133.1	2.1	6.04	.04		
S4-81	190.5	76.2 - 82.2	6.0	4.49	.095	YES	----
		106.6 - 109.6	3.0	4.01	.06		
		167.6 - 170.6	3.0	4.60	.13		
S5-81	215.5	39.6 - 42.6	3.0	5.47		YES	----
		82.0 - 85.0	3.0	15.39	.11		
		143.2 - 146.2	3.0	5.36	.06		
S6-81	185.0	143.2 - 146.2	3.0	4.27	.09	YES Structural	----
S7-81	224.3	131.0 - 134.0	3.0	19.09	.14	YES	----
		198.0 - 204.0	3.0	8.86	---	Structural	
S8-81	170.0	30.4 - 334.0	3.0	6.62	---	YES	----
		149.3 - 155.4	6.0	9.90	----	Structural	
S9-81	111.9	73.0 - 79.0	6.0	7.10	---	Structural	YES
S10-81	60.0	45.7 - 48.7	3.0	3.25	---	Structural	YES
S11-81	44.8	18.0 - 21.0	3.0	2.84	----	Structural	----
S12-81	96.6	70.0 - 73.0	3.0	9.16	----	Structural	YES
S13-81	69.0	48.7 - 51.7	3.0	5.26	----	Structural	----
S14-81	209.0	176.7 - 179.7	3.0	13.44	----	----	----
S15-81	133.0	62.4 - 65.4	3.0	6.39	----	----	----
S16-81	127.0	46.3 - 49.3	3.0	4.23	----	----	----
S17-81	130.0	86.5 - 89.5	3.0	7.17	----	Structural	YES
S18-81	145.0	82.0 - 85.0	3.0	9.19	----	Structural	----
		122.0 - 125.0	3.0	12.50	----		
S19-81	135.6	79.0 - 82.0	3.0	4.48	----	Structural	----
S20-81	66.0	39.6 - 42.6	3.0	5.45	----	Structural	----
S21-81	38.7	-----		----	----	Structural	----
S22-81	41.8	12.0 - 18.0	6.0	5.36	----	Structural	----
		24.0 - 27.0	3.0	5.26	----		
S23-81	32.6	15.0 - 21.0	6.0	8.86	----	Structural	----
S24-81	170.0	128.0 - 134.0	6.0	9.83	----	Structural	----
S25-81	80.1	52.0 - 58.0	6.0	4.44	----	Structural	----
S26-81	92.7	61.0 - 64.0	6.0	5.75	----	Structural	----

### 1981-1983 Diamond Drilling

(see Appendix C for assay data)

Drill Hole #	Depth (M)	Intercept (from - to)	Width	Assay CaF2%	Ag Assays	DDH Log	Cross Section
S27-81	255.0	204.0 - 207.0	3.0	14.42	-----	Structural	----
		234.0 - 237.0	3.0	5.26	-----		
S28-81	153.6	76.0 - 79.0	3.0	7.40	-----	Structural	----
		128.0 - 131.0	3.0	5.70	-----		
S29-81	221.5	102.4 - 103.6	1.2	11.50	-----	Structural	YES
S30-81	209.0	149.3 - 152.3	3.0	3.25	-----	Structural	----
S31-81	236.5	176.7 - 179.7	3.0	10.50	-----	Structural	----
S32-81	230.7	170.0 - 173.0	3.0	5.16	-----	Structural	----
S33-81	220.0	170.0 - 173.0	3.0	6.76	-----	Structural	----
		204.0 - 207.0	3.0	5.01	-----		
S34-81	179.0	137.0 - 146.0	9.0	10.10	-----	Structural	----
S35-81	155.4	56.0 - 62.0	6.0	4.10	-----	Structural	----
S36-81	169.7	137.0 - 140.0	3.0	6.04	-----	Structural	----
S37-81	175.8	90.0 - 93.0	3.0	11.30	-----	Structural	----
		121.0 - 122.8	1.8	15.20	-----		
S38-81	172.8	92.0 - 95.0	3.0	5.64	-----	Structural	----
		134.7 - 137.7	3.0	4.94	-----		
S39-81	163.6	30.4 - 33.4	3.0	4.89	-----	Structural	----
S40-81	91.7	15.8 - 18.8	3.0	7.08	-----	Structural	----
		80.0 - 83.0	3.0	5.57	-----		
S41-81	160.6	127.0 - 130.0	3.0	3.76	< .03	YES Structural	----
S42-81	175.8	84.0 - 87.0	3.0	4.40	.06	YES	----
						Structural	
S43-81	185.0	79.8 - 82.8	3.0	5.80	.44	YES	YES
		134.7 - 140.8	6.0	6.55	.045	Structural	
		147.0 - 162.0	15.0	9.22	.066		
S44-81	145.0	49.0 - 52.0	3.0	7.40	.06	YES	YES
		92.0 - 95.0	3.0	7.10	.06	Structural	
		98.0 - 101.0	3.0	7.20	.06		
		104.0 - 134.0	30.0	9.20	.05		
S45-81	175.5	68.0 - 71.0	3.0	5.70	.03	YES	YES
		120.0 - 123.0	3.0	5.60	.06	Structural	
		126.0 - 138.0	12.0	15.00	.075		
		141.0 - 150.0	9.0	6.30	.04		
S46-81	132.8	52.0 - 58.0	6.0	5.44	.03	YES	YES
		116.0 - 119.0	3.0	8.39	.03	Structural	
S47-81	142.0	16.7 - 25.9	9.0	7.93	.04	YES	YES
		83.8 - 86.8	3.0	7.06	.06	Structural	
		105.0 - 135.0	30.0	10.70	.07		
S48-81	148.4	31.4 - 34.0	3.0	5.33	.09	YES	YES
		98.0 - 122.0	24.0	17.50	.106	Structural	
S49-81	139.0	32.0 - 35.0	3.0	7.90	.09	YES	YES
		99.0 - 123.0	24.0	8.20	.078	Structural	
S50-82	153.3	123.7 - 132.8	9.0	6.60	-----	Structural	----
S51-82	160.0	79.0 - 82.0	3.0	13.70	-----	----	----
		91.0 - 94.0	3.0	8.30	-----		
S52-82	180.8	88.0 - 91.0	3.0	21.40	-----	----	YES
		170.0 - 173.0	3.0	8.12	-----		



### 1981-1983 Diamond Drilling

(see Appendix C for assay data)

Drill Hole #	Depth (M)	Intercept (from - to)	Width	Assay CaF2%	Ag Assays	DDH Log	Cross Section
S53-82	185.0	83.5 - 86.5 132.0 - 141.0 175.0 - 178.0	3.0 9.0 3.0	13.20 17.72 13.60	----- ----- -----	----	YES
S54-82	220.0	83.5 - 86.5 100.5 - 103.5 120.0 - 141.0 149.0 - 150.2 197.0 - 200.0 218.8 - 220.0	3.0 3.0 21.0 1.2 3.0 1.2	12.00 6.50 10.60 7.20 7.20 15.60	----- ----- ----- ----- ----- -----	----	YES
S55-82	142.0	33.3 - 34.5 35.8 - 37.6 53.6 - 56.6 75.0 - 79.8	1.2 1.8 3.0 4.8	8.60 9.00 8.00 10.93	----- ----- ----- -----	----	YES
S56-82	111.5	39.6 - 45.7 87.8 - 90.7	6.0 3.0	15.35 5.70	----- -----	----	YES
S57-82	97.8	38.0 - 47.0	9.0	10.70	-----	----	YES
S58-82	203.0	174.6 - 177.6 184.7 - 196.9	3.0 12.0	12.10 8.90	----- -----	----	YES
S59-82	117.6	55.7 - 58.8 72.5 - 75.5 115.5 - 114.5	3.0 3.0 3.0	25.60 13.20 15.80	----- ----- -----	Structural	----
S60-82	120.7	105.5 - 108.5 115.5 - 114.5	3.0 3.0	6.40 7.10	----- -----	Structural	YES
S61-82	130.0	117.0 - 120.0	3.0	3.86	-----	Structural	----
S62-82	160.0	53.6 - 56.6	3.0	7.30	-----	Structural	----
S63-82	157.0	45.0 - 47.3 102.7 - 108.8	2.3 6.0	43.00 6.75	----- -----	Structural	YES
S64-82	145.0	114.6 - 117.6 120.6 - 123.7	3.0 3.0	12.90 8.10	----- -----	Structural	----
S65-82	172.5	144.7 - 147.7 154.0 - 157.0	3.0 3.0	7.70 6.00	----- -----	Structural	----
S66-82	190.8	169.0 - 175.0	6.0	5.10	-----	Structural	YES
S67-82	195.0	174.0 - 177.0 189.0 - 192.0	3.0 3.0	16.30 14.70	----- -----	Structural	----
S68-82	267.0	255.0 - 258.0	3.0	13.10	-----	----	----
S69-82	157.0	123.0 - 126.0	3.0	4.80	-----	----	----
S70-82	166.0	126.0 - 129.0 132.0 - 135.0 145.0 - 148.0	3.0 3.0 3.0	11.00 6.90 6.90	----- ----- -----	----	YES
S71-82	120.7	82.6 - 105.0	22.5	11.00	-----	----	----
S72-82	248.7	229.0 - 232.0	3.0	4.80	-----	----	----
S73-82	190.8	-----	----	-----	-----	----	----
S74-82	65.8	50.0 - 53.0	3.0	3.90	-----	----	YES
S75-82	59.7	-----	-----	-----	-----	----	YES
S76-82	114.6	70.0 - 73.0	3.0	6.60	-----	----	----
S77-82	86.8	31.0 - 40.0	9.0	12.50	-----	----	YES
S78-82	108.5	42.6 - 45.6 86.0 - 95.0	3.0 9.0	12.10 11.60	----- -----	----	----
S79-82	102.4	-----	-----	-----	-----	----	----
S80-82	102.4	-----	-----	-----	-----	----	YES

### 1981-1983 Diamond Drilling

(see Appendix C for assay data)

Dill Hole #	Depth (M)	Intercept (from - to)	Width	Assay CaF2%	Ag Assays	DDH Log	Cross Section
S81-82	61.2	-----	----	-----	-----	----	----
S82-82	111.5	-----	----	-----	-----	Structural	----
S83-82	248.7	81.0 - 84.0 201.0 - 204.0	3.0 3.0	7.22 9.55	----- -----	----	----
S84-82	266.7	-----	----	-----	-----	----	----
S85-82	111.5	15.0 - 18.0	3.0	5.80	-----	Structural	----
S86-82	120.7	-----	----	-----	-----	Structural	----
S87-82	153	35.0 - 41.0 117.0 - 126.0 132.0 - 135.0	6.0 9.0 3.0	9.55 11.50 6.40	----- ----- -----	Structural	----
S88-82	136	50.0 - 53.0	3.0	6.90	-----	Structural	----
S89-82	139	-----	----	-----	-----	Structural	----
S90-82	-----	-----	----	-----	-----	----	----
S91-82	-----	-----	----	-----	-----	----	----
S92-83		74.0 - 89.0	6.0	9.40	-----	----	----
S93-83	-----	108 -129.0	21.0	14.17	-----	----	----
S94-83	-----	126 - 129.0	3.0	9.42	-----	----	----
S95-83	-----	127 - 133.0	6.5	11.06	-----	----	----
S96-83	Abandoned	-----	----	-----	-----	----	----
S97-83	-----	78.0 - 84.0	6.0	10.3	-----	----	----
S98-83	-----	115.0 - 118.0	3.0	11.8	-----	----	----
S99-83	-----	148.0 -151.0	3.0	8.63	-----	----	----
S100-83	-----	76.0 - 79.0	3.0	9.3	-----	----	----
S101-83	-----	136.0 - 139.0	3.0	11.8	-----	----	----
S102-83	-----	-----	----	-----	-----	----	----
S103-83	-----	-----	----	-----	-----	----	----
S104-83	-----	-----	----	-----	-----	----	----

### 1984-1985 Diamond Drilling

(see Appendix C for assay data)

S110-85	-----	-----	----	-----	-----	----	----
S111-85	-----	-----	----	-----	-----	Structural	----
S112-85	-----	-----	----	-----	-----	----	----
S113-85	-----	-----	----	-----	-----	----	----
S114-85	-----	-----	----	-----	-----	----	----
S115-85	-----	-----	----	-----	-----	----	----

APPENDIX E  
**LOGGED DRILL HOLE COORDINATES**

## Location of Logged Drill Holes

Holes	Coordinates (m)		Coordinates (m)		Collar elevation (m)
S1-81	1.8	North	258.9	West	908.61
S2-81	107.6	South	254.8	West	864.72
S6-81	2.0	South	402.0	West	899.61
S7-81	23.3	South	550.9	West	892.15
S8-81	110.0	South	405.0	West	874.18
S9-81	264.1	South	491.7	West	799.84
S10-81	347.3	South	350.8	West	800.90
S11-81	301.8	South	433.0	West	798.87
S12-81	347.7	South	580.6	West	765.80
S13-81	372.2	South	550.0	West	766.74
S17-81	172.5	South	421.9	West	841.31
S18-81*	54.0	South	325.0	West	894.0
S19-81	191.4	South	255.4	West	880.54
S20-81	248.2	South	384.7	West	824.03
S21-81	264.3	South	344.2	West	825.31
S22-81	268.2	South	376.7	West	818.43
S23-81	276.1	South	387.2	West	806.60
S24-81	247.0	South	692.0	West	778.39
S25-81	196.0	South	329.0	West	856.80
S26-81	200.0	South	394.0	West	840.35
S27-81	61.0	North	260.0	West	927.80
S28-81	1.4	North	200.1	West	914.36
S29-81	70.0	North	186.0	West	933.75
S30-81	117.0	North	128.0	West	940.07
S31-21	136.0	North	202.0	West	942.56
S32-81	192.0	North	100.0	West	947.23
S33-81	57.6	North	320.9	West	925.88
S34-81	0.6	South	321.9	West	905.89
S35-81	82.1	South	540.0	West	874.43
S36-81	191.2	South	555.2	West	819.90
S37-81	129.7	South	770.5	West	820.02
S38-81	77.3	South	860.9	West	823.82
S39-81	36.5	South	769.2	West	860.10
S40-81	30.8	South	679.4	West	873.74
S41-81	60.0	North	780.0	West	876.00
S42-81	40.0	South	600.0	West	888.00
S43-81	1.0	North	293.0	West	907.50
S44-81	33.0	South	259.0	West	904.00
S45-81	1.0	North	223.0	West	911.60
S46-81	30.0	South	330.0	West	900.10
S47-81	65.5	South	269.5	West	895.50
S48-81	73.6	South	241.4	West	896.90
S49-81	54.5	South	303.4	West	893.90
S50-82	32.0	North	258.6	West	918.67
S59-82	103.8	South	284.2	West	881.66
S60-82	64.5	South	338.8	West	886.19

\* coordinates derived from property map

Holes	Coordinates (m)		Coordinates (m)		Collar elevation (m)
S61-82	60.4	South	368.4	West	888.44
S62-82	3.7	North	348.7	West	905.62
S63-82	90.9	South	371.6	West	878.80
S64-82	93.1	South	397.5	West	876.19
S65-82	85.5	South	457.7	West	873.80
S66-82	31.7	South	299.0	West	921.04
S67-82	36.1	North	292.0	West	919.15
S82-82	156.5	South	301.3	West	869.90
S85-82	323.7	South	696.0	West	756.60
S86-82	290.3	South	753.3	West	759.30
S87-82	261.4	South	808.2	West	757.50
S88-82	226.0	South	857.4	West	759.50
S89-82	197.7	South	901.0	West	758.50
S111-85*	Near S63-82				

\*Location approximate

APPENDIX F  
DRILL HOLE LOGS

<b>Project</b>	<b>Client</b>	Eaglet Mines Ltd.		<b>Collar Elevation</b>	908.61
<b>Drill hole</b>	Structural analysis			<b>Inclination</b>	90
<b>Coordinates</b>	S1-81			<b>Length</b>	187.4 m
	1.75 m North			<b>Logged by:</b>	P.B.Read
	258.9 m West				

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
0.0	30.0	0.0	9.1		Overburden				
30.0	86.9	9.1	26.5		Medium grained (2 mm) biotite muscovite schist	34.0	10.4	Foliation	5
						50.0	15.2	Foliation	17
						55.0	16.8	Foliation	15
						58.0	17.7	Foliation	10
						66.0	20.1	Foliation	10
						76.0	23.2	Foliation	20
						77.0	23.5	to 80.0' (24.4 m) heavily muscovitized and possibly faulted	
						80.0	24.4	First appearance of gypsum, potassium feldspar veins and galena	
						82.0	25.0	First appearance of coarse pyrite	
						85.0	25.9	Foliation	20
						85.0	25.9	Pyrite, potassium feldspar, quartz and calcite vein. 010/75SE	
86.9	87.2	26.5	26.6		Medium grained (2 mm) pink muscovite gneiss sill				
87.2	97.0	26.6	29.6		Medium grey, medium grained (1 mm) biotite- muscovite schist	90.0	27.4	Foliation	5
						92.0	28.0	Vein (15 mm) with vuggy quartz, calcite and K-feldspar parallel to foliation.	
						92.0	28.0	Foliation	20
97.0	102.0	29.6	31.1		Medium grained (1 mm) pink aplitic gneiss sill and thin slivers of Snowshoe schist. Sample S5-81-101. Alteration: pyroxene ?				
102.0	109.0	31.1	33.2		Medium grained (2 mm) biotite muscovite schist	108.0	32.9	Foliation	13
109.0	115	33.2	35.1		Medium grained (1 mm) biotite (10%) aplitic gneiss	112.0	34.1	Foliation	25
115.0	118	35.1	36.0		Medium grained (1-2 mm) biotite muscovite schist	117.0	35.7	Foliation	0
118.0	123.0	36.0	37.5		Pink aplitic sill and gneissic aplite upper margin 2 feet thick.	121.0	36.9	to 123.0 (37.5 m) coarse pegmatitic calcite quartz fluorite zone. First appearance of fluorite	

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
123.0	151.0	37.5	46.0		White aplite sill with disseminated fluorite and local gneissic aplitic layers.	142.0	43.3	Foliation	10
151.0	175.0	46.0	53.3		Medium grained (1-2 mm) biotite, muscovite schist (Snowshoe)	152.0	46.3	Foliation	21
						154.0	46.9	Foliation	38
175.0	177.0	53.3	54.0		Pink aplitic sill				
177.0	188.0	54.0	57.3		Fault gouge and breccia. Sericitized biotite schist fragments.			Base of Snowshoe	
188.0	193.0	57.3	58.8		Medium grained (1-2 mm) altered biotite schist.	190.0	57.9	Foliation. Vein (2 mm) of gypsum; 090/vertical	43
193.0	196.0	58.8	59.7		Medium grained (2-3 mm) leuco granodioritic gneiss.	193.0	58.8	Foliation	31
196.0	207.0	59.7	63.1		Pink, slightly gneissic aplite and disseminated fluorite.				
207.0	213.0	63.1	64.9		Medium grained (2-4 mm) leucogranodioritic gneiss.				
213.0	241.5	64.9	73.6		Pink gneissic aplite and disseminated fluorite and muscovite-bearing.	220.0	67.1	Foliation	41
						237.0	72.2	Foliation	34
						237.5	72.4	Foliation	41
						239.0	72.8	Foliation	
						239.0	72.8	Vein (1 mm) with K-feldspar. 090/37S.	
						241.0	73.5	Foliation	48
241.5	249.0	73.6	75.9		Pink to white aplite with disseminated fluorite and pyrite.	244.0	74.4	Disseminated fluorite and pyrite vein parallel to foliation.	45
						244.0	74.4	Vein (2 mm) of quartz- K-feldspar and galena. 090/45S	
249.0	262.0	75.9	79.9		Cream to pink muscovite-bearing aplitic medium grained (1-2 mm) gneiss and many gypsum veins.	256.0	78.0	Foliation	35
						261.0	79.6	Foliation	50
						261.0	79.6	Vein (3 mm) of gypsum. 300/41SW.	
262.0	296.0	79.9	90.2		White aplite with disseminated fluorite and pyrite	269.0	82.0	First appearance of pyroxene alteration for certain.	
						288.0	87.8	Foliation	90
						288.0	87.8	Vein of hematite- k-feldspar- pyrite-galena and quartz. 010/30SE	
						293.0	89.3	Foliation	90



<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
296.0	305.0	90.2	93.0		Pink, medium grained (1 mm) muscovite bearing aplitic gneiss.	303.0	92.4	Foliation	46
						311.0	94.8	Disseminated fluorite parallel to foliation	55
305.0	319.0	93.0	97.2		Pink, medium grained (2 mm) aplite and disseminated fluorite.				
319.0	324.0	97.2	98.8		Medium grained (1-3 mm) biotite (6%) gneiss. Coarse fluorite.	319.0	97.2	Foliation	51
						322.0	98.1	Foliation	55
						326.0	99.4	Foliation	
						330.0	100.6	Disseminated fluorite parallel to foliation	46
						349.0	106.4	Foliation	66
334.0	364.0	101.8	110.9		Pink to locally white aplite and biotite aplitic gneiss.				
364.0	368.0	110.9	112.2		Pink, medium grained (2 mm) biotite- muscovite gneiss.				
368.0	413.0	112.2	125.9		Pink pegmatitic aplite and disseminated fluorite	413.0	125.9	Fault	37
413.0	438.0	125.9	133.5		Medium grained leucogranodioritic gneiss cut by many gypsum veinlets. The gneiss has less than 7% pyroxene.	435.0	132.6	Foliation	41
						437.0	133.2	Foliation	45
438.0	441.0	133.5	134.4		White aplite.				
441.0	447.0	134.4	136.2		Medium grained (1-3 mm) pyroxene (5%) leucogranodioritic gneiss.	445.0	135.6	Foliation	90
						446.0	135.9	Foliation	68
447.0	451.0	136.2	137.5		Fine grained (1 mm) biotite (5%) aplitic gneiss.	448.0	136.6	Foliation	73
451.0	452.0	137.5	137.8		Medium grained (2-4 mm) pink leucogranodioritic gneiss.				
452.0	455.0	137.8	138.7		Medium grained (1 mm) biotite-pyroxene (20%) aplitic gneiss.				
455.0	457.0	138.7	139.3		White aplite.				
457.0	461.0	139.3	140.5		Medium grained (1-3 mm) biotite-pyroxene (10%) granodioritic gneiss.	45.0	13.7	foliation	40
461.0	528.0	140.5	160.9		White, locally pink aplite with disseminated fluorite. Sample S1-81-505. Sample for thin section.	472.0	143.9	Vein of disseminated fluorite parallel to foliation.	36
						473.0	144.2	Foliation	62

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
528.0	607.0	160.9	185.0		Medium grained (2-3 mm) biotite (5%) leucogranodioritic gneiss with many gypsum veinlets.	520.0	158.5	Foliation	60
						534.0	162.8	Foliation	23
						536.0	163.4	Foliation	90
						541.0	164.9	Foliation	90
						559.0	170.4	Foliation	59
						562.0	171.3	Foliation	58
						575.0	175.3	Vein (3 mm) of gypsum parallel to foliation	51
						576.0	175.6	Foliation	64
						579.0	176.5	Foliation	70
						599.0	182.6	Foliation	50
						604.0	184.1	Foliation	38
607.0	611.0	185.0	186.2		Pink aplite with heavy gypsum alteration.	611.0	186.2	Foliation	45
611.0	615.0	186.2	187.5		Medium grained (2-4mm) leucogranodioritic gneiss				
	615.0		187.5		End of hole.				

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>	
				<b>Client</b>	Eaglet Mines Ltd.					
<b>Project</b>	Structural analysis								<b>Collar Elevation</b>	884.72
<b>Drill hole</b>	S2-81								<b>Inclination</b>	90
<b>Coordinates</b>	107.6 m South								<b>Length</b>	125.9 m
	254.8 m West								<b>Logged by:</b>	JAF

Note: Assumed strike of foliation ~ 090. Core has been completely split.

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
0.0	4.0	0.0	1.2		Overburden				
4.0		1.2		QLG	Pink to cream coloured aplite and aplite gneiss with muscovite zones. Disseminated fluorite common.				
	14.0				Biotite- K-feldspar alteration zone. Randomly oriented biotite and muscovite abundant. K-feldspar and minor fluorite veins (small- 1-2 mm present form top of hole). Gypsum abundant in several fracture orientations.	22.0	6.7	Foliation	60
						27.0	8.2	Foliation	40
						31.0	9.4	Foliation	40
						31.0	9.4	Vein (2mm) of gypsum. 010/55NW.	
						33.0	10.1	Foliation	55
						37.0	11.3	Foliation	45
						40.0	12.2	Foliation	20
						43.0	13.1	Foliation	15
						47.0	14.3	Vein (1mm) with pyrite-fluorite- K-feldspar, parallel to foliation	25
						47.0	14.3	Vein of galena- fluorite-pyrite and K-feldspar. 085/47S	
						51.0	15.5	Foliation and zone of biotite-fluorite and pyrite, 3 to 4 cm wide.	45
						51.0	15.5	Vein of quartz-fluorite and pyrite, 2 mm. 160/42 SW.	
						60.0	18.3	Foliation	60
						65.0	19.8	Foliation	60
						70.0	21.3	Foliation	55
						73.0	22.3	Foliation	45
						73.0	22.3	Vein of K-feldspar-fluorite-pyrite and galena. 165/48SW.	
						77.0	23.5	Foliation	45

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						80.0	24.4	Foliation	40
						80.0	24.4	Vein with K-feldspar-fluorite and pyrite. 085/25S	
						82.0	25.0	Foliation	30
						82.0	25.0	Vein of gypsum, 2 mm in fracture 090/83N.	
						90.0	27.4	Foliation	40
						93.0	28.3	Foliation	35
						93.0	28.3	Vein with K-feldspar-calcite and pyrite, 2 mm. 090/40S.	
						97.0	29.6	Foliation	50
						97.0	29.6	Vein of gypsum, 1 mm in fracture 145/83SW.	
110	170.0	33.5	51.8		Medium grained pyroxene (1%) granodioritic gneiss with muscovite-aplite gneiss still in abundance. Pyroxene as alteration also present in conjunction with fluorite.	103.0	31.4	Foliation	40
						121.0	36.9	Foliation	45
						124.0	37.8	Foliation	25
						124.0	37.8	Quartz vein with slicks (55-> SW). 175/60 (vein).	
						130.0	39.6	Foliation	10
						140.0	42.7	Foliation	15
						145.0	44.2	Foliation	30
						147.0	44.8	Foliation	25
						155.0	47.2	Foliation	30
						160.0	48.8	Foliation	30
						163.0	49.7	Foliation	30
						163.0	49.7	Vein of gypsum in fracture. 1mm. 045/90.	
						167.0	50.9	Foliation	Horizontal
170	194.0	51.8	59.1			Biotite and muscovite abundant in pink aplite and aplite gneiss	172.0	52.4	Foliation
						188.0	57.3	Foliation	Horizontal
194	210.0	59.1	64.0		Granodiorite gneiss and lesser pink aplite with local pyroxene (1%) and cream coloured fine grained aplite.	194.0	59.1	Foliation	40
						203.0	61.9	Foliation	25
						203.0	61.9	Vein of gypsum in fracture. 3 mm. 015/84 NW.	
						206.0	62.8	Foliation	25

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
210.0	303.0	64.0	92.4		Coarse grained granodiorite gneiss with minor aplite and muscovite- aplite.	210.0	64.0	Foliation	30
						215.0	65.5	Foliation	30
						215.0	65.5	Vein (2 mm) of gypsum with slicks. Vein: 108/38SW. Slicks: 30-> W.	
						220.0	67.1	Foliation	35
						225.0	68.6	Foliation	30
						236.0	71.9	Foliation	Horizontal
						245.0	74.7	Foliation	48
						250.0	76.2	Vein (4 mm) of fluorite and K-feldspar. Parallel to foliation.	45
						260.0	79.2	Foliation	30
						268.0	81.7	Vein-fracture (3 mm) with gypsum. 110/68 SW.	
						275.0	83.8	Foliation	25
						280.0	85.3	Vein (2 cm) of fluorite, calcite and pyrite parallel to foliation.	40
						285.0	86.9	Foliation	40
						291.0	88.7	Foliation	30
303.0	327.0	92.4	99.7		Cream and white aplite with granodiorite gneiss. Local fluorite ore in pockets.	303.0	92.4	Foliation	40
						306.0	93.3	Foliation	75
						319.0	97.2	Foliation	30
327.0	413.0	99.7	125.9		Medium grained granodiorite gneiss with fine grained areas, pyroxene up to 5%.	327.0	99.7	Foliation	45
						327.0	99.7	Vein of quartz and pyrite. 2 mm. 105/40 SW.	
						334.0	101.8	Foliation	45
						334.0	101.8	Vein of fluorite and K-feldspar, 1 cm wide. 010/74 NW.	
						340.0	103.6	Foliation	40
						345.0	105.2	Foliation	50
						353.0	107.6	Foliation	45
						357.0	108.8	Foliation	20
						362.0	110.3	Foliation	45
						362.0	110.3	Vein of gypsum. 1-2 mm. 095/65SE,	
						377.0	114.9	Vein of quartz- K-feldspar- fluorite and galena parallel to foliation.	50
						382.0	116.4	Vein of gypsum in fracture. 1 mm. 065/65SE.	

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						382.0	116.4	Foliation	45
						385.0	117.3	Foliation	45
						385.0	117.3	Vein of gypsum and chlorite in fracture. 2 mm. 018/58 SE.	
						392.0	119.5	Foliation	40
						397.0	121.0	Vein (6 mm) of quartz and fluorite parallel to foliation.	45
						402.0	122.5	Foliation	35
						402.0	122.5	Vein (3 mm) of quartz and minor pyrite. 087/54S.	
						409.0	124.7	Foliation	50
	413.0		125.9		End of hole				

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
<b>Project</b>	<b>Client</b>			Eaglet Mines Ltd.				<b>Collar Elevation</b>	899.61 m
<b>Drill hole</b>	Structural analysis							<b>Inclination</b>	90
<b>Coordinates</b>	S6-81							<b>Length</b>	185.0 m
	2 m South							<b>Logged by:</b>	P.B. Read
	402 m West								
<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
0.0	20.0	0.0	6.1		Overburden				
20.0	122.0	6.1	37.2		Medium grey medium grained (2 mm) biotite ± muscovite schist. 66.5 to 67.5' (20.3-20.6 m) grey biotite aplite.	26.0	7.9	Foliation	21
						33.0	10.1	Foliation	17
						52.0	15.8	Foliation	11
						63.0	19.2	Foliation	5
						63.0	19.2	Vein (3 mm) of quartz and calcite. Unknown attitude.	
						66.0	20.1	Vein (3 mm) of quartz and pyrite. Unknown attitude.	
						70.0	21.3	Vein (3 mm) of K-feldspar. Unknown attitude.	
						70.0	21.3	First appearance of K-feldspar.	
						72.0	21.9	Foliation	5
						85.0	25.9	Foliation	15
						100.0	30.5	Foliation	19
122.0	125.0	37.2	38.1		Fine grained (1 mm) pink aplitic gneiss.	122.0	37.2	Contact broken and probably faulted	
125.0	130.0	38.1	39.6		Medium grained (2 mm) leuco-granodiorite gneiss with disseminated fluorite. Sample S2-81-130'.	125.0	38.1	First appearance of fluorite.	
						126.0	38.4	First appearance of pyroxene (?).	
						127.0	38.7	First appearance of gypsum.	
130.0	136.0	39.6	41.5		White and pink aplite. Medium grained (2 mm) pink muscovite gneiss.	130.0	39.6	First appearance of certain pyroxene.	
136.0	142.0	41.5	43.3		White to pink aplite with disseminated fluorite and pyrite.	136.0	41.5	Foliation	20
						140.0	42.7	Foliation	29
142.0	152.0	43.3	46.3		Medium grained (2-3 mm) pyroxene (5%) leuco-granodiorite gneiss.	142.0	43.3	Foliation	33
152.0	156.0	46.3	47.5		Pink to white aplite with disseminated fluorite	155.0	47.2	Disseminated fluorite parallel to foliation	27

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
156.0	160.0	47.5	48.8		Pink medium grained (2 mm) muscovite gneiss.	159.0	48.5	Foliation	29
160.0	161.0	48.8	49.1		White aplite.				
161.0	162.0	49.1	49.4		Pink medium grained (2 mm) muscovite gneiss.	161.5	49.2	Foliation	30
162.0	163.0	49.4	49.7		Pink to white aplite.	163.0	49.7	Contact between aplite and gneiss.	
163.0	175.0	49.7	53.3		Medium grained (2-4 mm) pyroxene (2-10%) leuco-granodiorite gneiss and pyroxene alteration zones.	165.0	50.3	Foliation	33
175.0	181.0	53.3	55.2		Pink medium grained (2 mm) muscovite gneiss.	176.0	53.6	Foliation	24
181.0	184.0	55.2	56.1		Medium grained (1-3 mm) pyroxene (3%) leuco-granodiorite gneiss.	182.0	55.5	Foliation	30
184.0	190.0	56.1	57.9		Heavily altered (gypsum) muscovite gneiss aplite.				
190.0	213.0	57.9	64.9		Medium grained leuco-granodiorite gneiss altered by gypsum.				
213.0	219.0	64.9	66.8		Pink medium grained (2-3 mm) muscovite gneiss heavily altered by gypsum				
219.0	221.0	66.8	67.4		Medium grained (2-34 mm) pinkish leuco granodiorite gneiss.	220.0	67.1	Foliation	43
						220.0	67.1	Vein (2 mm) of gypsum. 305/55SW.	
221.0	226.0	67.4	68.9		White to pink aplite sill with disseminated fluorite and pyrite.	221.0	67.4	Contact between gneiss and aplite parallel to foliation.	54
226.0	252.0	68.9	76.8		Medium grained (2-4 mm) pyroxene (2 %) leuco-granodiorite gneiss.	227.0	69.2	Foliation	46
						239.0	72.8	Foliation	45
						245.0	74.7	Foliation	54
252.0	266.0	76.8	81.1		Pink aplite and scarcely disseminated fluorite and pyrite.	252.0	76.8	Foliation	52
						265.0	80.8	Foliation	53
266.0	275.0	81.1	83.8		Medium grained (2-4 mm) pyroxene (5 %) leuco-granodiorite gneiss. Sample S2-81-268', typical granodiorite gneiss; thin section.	268.0	81.7	Foliation	39
275.0	287.0	83.8	87.5		Pink aplite.				



<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
287.0	331.0	87.5	100.9		Medium grained (2-4 mm) pyroxene (2 %) leuco-granodiorite gneiss. Sample S2-81-302' Typical thin section.	292.0	89.0	Foliation	35
						326.0	99.4	Foliation	47
331.0	336.0	100.9	102.4		Pink aplite.	336.0	102.4	Disseminated fluorite parallel to foliation	51
336.0	358.0	102.4	109.1		Medium grained (2-4 mm) pyroxene (5 %) leuco-granodiorite slightly gneissic.	349.0	106.4	Disseminated fluorite parallel to foliation	42
						357.0	108.8	Foliation	45
358.0	391.0	109.1	119.2		Pink aplite with disseminated fluorite. Locally gneissic.	359.0	109.4	Foliation	48
						372.0	113.4	Foliation	35
						380.0	115.8	Foliation	38
391.0	396.0	119.2	120.7		Medium grained (2-4 mm) leuco-granodiorite gneiss.	395.0	120.4	Foliation	53
						395.0	120.4	Vein(1 mm) of gypsum. 032/60SE.	
396.0	401.0	120.7	122.2		Pink aplite and medium grained (2 mm) pink muscovite gneiss.	400.0	121.9	Foliation	50
						400.0	121.9	Vein (1 mm) of K-feldspar. 090/40S.	
401.0	404.0	122.2	123.1		Medium grained (2-4 mm) pyroxene (2 %) leuco-granodiorite gneiss.	402.0	122.5	Disseminated fluorite parallel to foliation	51
404.0	423.0	123.1	128.9		Pink fine grained (1 mm) aplite and gneissic aplite. Sample S2-81-416'. Typical pink gneissic aplite.	408.0	124.4	Disseminated fluorite parallel to foliation	46
						408.0	124.4	Vein (2 mm) of quartz. 028/74NW.	
						412.0	125.6	Foliation	51
						413.0	125.9	Foliation	55
						413.0	125.9	Vein(2 mm) of gypsum. 090/90..	
						422.0	128.6	Foliation	50
423.0	426.0	128.9	129.8		Medium grained (2-4 mm) pink leuco-granodiorite gneiss.	423.0	128.9	Foliation	55
426.0	446.0	129.8	135.9		Pink aplite and gneissic aplite.	439.0	133.8	Foliation	38
						439.0	133.8	Vein (4 mm) of pyrite, quartz and fluorite. 005/37SE.	
446.0	448.0	135.9	136.6		Medium grained (2-3 mm) pink muscovite gneiss.	447.0	136.2	Foliation	43
448.0	467.0	136.6	142.3		Pink aplite and gneissic aplite.	452.0	137.8	Foliation	50
						457.0	139.3	Foliation	43

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						462.0	140.8	Foliation	53
467.0	473.0	142.3	144.2		Medium grained (2-3 mm) pink leuco-granodiorite gneiss.	472.0	143.9	Foliation	43
473.0	478.0	144.2	145.7		Pink aplite with disseminated fluorite.	478.0	145.7	Foliation	62
478.0	493.0	145.7	150.3		Medium grained (2-4 mm) leuco-granodiorite gneiss.	482.0	146.9	Foliation	67
493.0	503.0	150.3	153.3		Pink to white aplite with disseminated fluorite and pyrite.	501.0	152.7	Foliation	47
						502.0	153.0	Foliation	47
						502.0	153.0	Vein (2 mm) of quartz and pyrite. 090/43S.	
503.0	508.0	153.3	154.8		Grey pyroxene (15%) granodiorite gneiss.	504.0	153.6	Foliation	47
508.0	517.0	154.8	157.6		Pink aplite with disseminated fluorite and gneissic aplite.	509.0	155.1	Foliation	55
						511.0	155.8	Foliation	36
						514.0	156.7	Foliation	52
517.0	522.0	157.6	159.1		Medium grained (2-4 mm) pink pyroxene (3%) leuco-granodiorite gneiss.	517.0	157.6	Foliation	43
522.0	551.0	159.1	167.9		Pink to white aplite. Sample S6-81-532' for opaques. Pink muscovite gneiss slivers present.	543.0	165.5	Foliation	45
						544.0	165.8	to 602' (183.5 m). Numerous gypsum veins	
551.0	553.0	167.9	168.6		Medium grained leuco-granodiorite gneiss heavily altered by gypsum.	553.0	168.6	Foliation	40
553.0	561.0	168.6	171.0		Fine grained (1 mm) biotite (30%) gneiss heavily altered by gypsum.	560.0	170.7	Gypsum veinlets parallel to foliation.	45
561.0	567.0	171.0	172.8		Pink aplite.				
567.0	574.0	172.8	175.0		Fine grained (1 mm) biotite - pyroxene aplitic gneiss.	569.0	173.4	Foliation	50
574.0	605.0	175.0	184.4		Medium grained (1-3 mm) pyroxene (10 %) leuco-granodiorite gneiss.	575.0	175.3	Foliation	45
						593.0	180.7	Foliation	57
						603.0	183.8	Foliation	52
605.0	607.0	184.4	185.0		Medium grained (1 mm) biotite (40%) schist.				
	607.0		185.0		End of hole.				

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>	
				<b>Client</b>	Eaglet Mines Ltd.					
<b>Project</b>	Structural analysis								<b>Collar Elevation</b>	892.15 m
<b>Drill hole</b>	S7-81								<b>Inclination</b>	90
<b>Coordinates</b>	23.3 m South								<b>Length</b>	224.2 m
	550.9 m West								<b>Logged by:</b>	JAF

Note: Assumed strike of foliation ~ 054°.Core was completely split.

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
0.0	20.0	0.0	6.1		Overburden				
20.0	42.0	6.1	12.8	QLG	Pink aplite and altered muscovite aplite gneiss (muscovite up to 15%).	22.0	6.7	Foliation	55
						42.0	12.8	Foliation	50
						47.0	14.3	Foliation	55
						63.0	19.2	Foliation	45
						63.0	19.2	Vein(2 mm) of barren quartz. 160/75SW.	
						78.0	23.8	Foliation	45
						78.0	23.8	Vein (2 cm) of quartz (90%) and fluorite. 140/35NE.	
						82.0	25.0	Foliation	50
						82.0	25.0	Vein (3 mm) of quartz. 055/45SE.	
						90.0	27.4	Foliation	45
						90.0	27.4	Vein (8 cm) of fluorite, quartz and pyrite. 055/50NW.	
						107.0	32.6	Foliation	
						107.0	32.6	Vein (3 cm) of quartz and fluorite parallel to S1. Foliation	55
						127.0	38.7	Foliation	45
						137.0	41.8	Foliation	45
						141.0	43.0	Foliation	40
						141.0	43.0	Vein (5 mm) of quartz, fluorite and pyrite. 056/90	
						153.0	46.6	Foliation	30
						177.0	54.0	Foliation	50
						177.0	54.0	Vein (2 mm) of quartz, fluorite and pyrite. 10/50NE.	
						187.0	57.0	Foliation	45
						187.0	57.0	Vein (3 mm) of K-feldspar and pyrite. 055/50SE.	
						201.0	61.3	Foliation	0
						223.0	68.0	Foliation	35
						232.0	70.7	Foliation	40

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						239.0	72.8	Vein (3 cm) of quartz, fluorite, pyrite and pyroxene parallel to foliation.	30
						251.0	76.5	Vein (2-3 mm) of pyroxene and fluorite parallel to foliation	55
						255.0	77.7	Vein (1 cm) of calcite and disseminated fluorite parallel to foliation.	35
						276.0	84.1	Foliation	45
						276.0	84.1	Vein (3 mm) of barren quartz. 055/50SE.	
						279.0	85.0	Foliation	50
						279.0	85.0	Vein (2 mm) of fluorite and quartz.	
						284.0	86.6	Foliation	50
						298.0	90.8	Foliation	40
						298.0	90.8	Vein (3 mm) of quartz, fluorite and pyrite. 055/60SE.	
						303.0	92.4	Foliation	55
						303.0	92.4	Vein (4 mm) of fluorite and minor quartz. 040/45SE.	
						312.0	95.1	Foliation	35
						328.0	100.0	Vein (2 cm) of fluorite and pyroxene parallel to foliation	45
						334.0	101.8	Foliation	50
						345.0	105.2	Disseminations of fluorite ~ 3 mm wide parallel to foliation.	45
						352.0	107.3	Foliation	50
						352.0	107.3	Vein (4 mm) of fluorite, sphalerite and quartz. 110/60NE.	
						362.0	110.3	Foliation	60
						362.0	110.3	Vein of quartz and fluorite. 055/30SE.	
						372.0	113.4	Disseminations of fluorite and pyroxene ~ 3 cm wide parallel to foliation.	50
						375.0	114.3	Foliation	50
						375.0	114.3	Vein-fracture (1 mm) with kaolinite. 170/50SE.	
						387.0	118.0	Foliation	40
						387.0	118.0	Vein-fracture (2 mm) with kaolinite. 150/65NE.	
						405.0	123.4	Foliation	50
						411.0	125.3	Foliation	50
						420.0	128.0	Foliation	55

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						432.0	131.7	Broad zone of fluorite + pyrite mineralization ~ 20 cm wide parallel to foliation.	50
						442.0	134.7	Foliation	45
						442.0	134.7	Vein of K-feldspar and pyrite. 135/10SW.	
						450.0	137.2	Foliation	40
						450.0	137.2	Vein of K-feldspar, quartz, fluorite and pyrite. 032/52SE.	
						455.0	138.7	Foliation	45
						455.0	138.7	Vein of quartz, fluorite, galena and sphalerite. 080/50SE.	
						461.0	140.5	Vein (5 mm) of fluorite parallel to foliation	45
						463.0	141.1	Foliation	50
						465.0	141.7		55
						465.0	141.7	Vein (1 mm) of quartz, pyrite and molybdenite. 090/50S.	
						474.0	144.5	Vein (2-3 mm) of fluorite parallel to foliation.	45
						486.0	148.1	Foliation	35
						507.0	154.5	Foliation	60
						510.0	155.4	Foliation	35
						510.0	155.4	Vein (5 mm) of quartz and minor fluorite, pyrite and molybdenite. 030/80SE.	
						522.0	159.1	Vein (5 mm) of purple and pink fluorite parallel to foliation.	45
						526.0	160.3	Foliation	50
						531.0	161.9	Foliation	50
						544.0	165.8	Foliation	45
						549.0	167.3	Foliation	50
						549.0	167.3	Fracture (1-2 mm) with kaolinite. 035/80NW.	
						567.0	172.8	Foliation	55
						567.0	172.8	Vein-fracture. 040/80SE.	
						567.0	172.8	Slickensides 70/180	
						576.0	175.6	Foliation	60
						578.0	176.2	Foliation	60
						578.0	176.2	Vein (5 mm) of pure fluorite. 160/88NE.	
						578.0	176.2	Vein (2-3 cm) of quartz, pyroxene and minor fluorite. 010/60E.	
						581.0	177.1	Foliation	50
						583.0	177.7	Foliation	50

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						583.0	177.7	Three veins (5 mm wide) of barren quartz and minor fluorite. 035/45SE.	
						595.0	181.4	Foliation	50
						601.0	183.2	Foliation	50
						603.0	183.8	Foliation	45
						612.0	186.5	Vein (2 mm) of fluorite and quartz parallel to foliation.	45
						617.0	188.1	Vein (1 cm) of fluorite parallel to foliation.	55
						627.0	191.1	Foliation	45
						640.0	195.1	Foliation	50
						640.0	195.1	Vein-zone (3 cm wide) of fluorite, pyroxene and K-feldspar. 055/55NW.	
						654.0	199.3	Vein (2 cm) of K-feldspar, fluorite, pyroxene and pyrite parallel to foliation	45
						656.0	200.0	Foliation	60
						656.0	200.0	Vein of fluorite, quartz, pyrite and molybdenite. 070/42SE.	
						663.0	202.1	Foliation	55
						673.0	205.1	Foliation	60
						673.0	205.1	Vein of quartz, fluorite, pyrite and galena. 055/35SE.	
						687.0	209.4	Foliation	50
						690.0	210.3	Vein (5 mm) of fluorite and pyrite parallel to foliation.	50
						692.0	210.9	Foliation	55
						692.0	210.9	Vein-fracture (1-2 mm) with kaolinite. 035/48SE.	
						693.0	211.2	Foliation	50
						703.0	214.3	Foliation	40
						703.0	214.3	Vein (3 mm) of gypsum with pyrite at margins. 050/85SE.	
						707.0	215.5	Foliation	50
						707.0	215.5	Vein (2 mm) of quartz, galena and pyrite. 055/48SE.	
						712.0	217.0	Foliation	50
						712.0	217.0	Vein (1 mm) of K-feldspar and pyrite. 090/70S.	
						720.0	219.5	Foliation	60
	735.7		224.2		End of hole.				

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>	
				<b>Client</b>	Eaglet Mines Ltd.					
<b>Project</b>	Structural analysis								<b>Collar Elevation</b>	874.18 m
<b>Drill hole</b>	S8-81								<b>Inclination</b>	90
<b>Coordinates</b>	110 m south 405 m west								<b>Length</b>	173.7 m
								<b>Logged by:</b>	JAF	

Note: Assumed strike of foliation ~ 054°. Core was completely split. Core is at level of adit #2 at 278'

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
0.0	40.0	0.0	12.2		Overburden				
40.0	68.0	12.2	20.7	Snow-shoe	Quartz-feldspar-biotite schist with local layer and augen of K-feldspar (alteration). Dark grey to black granular schist.	50.0	15.2	Foliation	20
						63.0	19.2	Foliation	10
						67.0	20.4	Foliation	30
68.0	89.0	20.7	27.1	QLG	Pink to cream aplite with zones of altered fine grained muscovite aplite gneiss well foliated for first 100' (30.49 m) Calcite and gypsum alteration noted at 75 feet (22.9 m).	83.0	25.3	Foliation	60
89.0	123.0	27.1	37.5		Pink and local white aplite. Strong pyroxene/fluorite alteration. Muscovite absent. Pink aplite with no muscovite.	93.0	28.3	Disseminated fluorite and pyroxene in zones of 1 cm parallel to foliation.	50
						100.0	30.5	Foliation	65
						106.0	32.3	Foliation	40
						121.0	36.9	Foliation	35
						123.0	37.5	Foliation	65
						129.0	39.3	Three veins with fluorite and pyrite parallel to foliation	40
						137.0	41.8	Foliation	50
						143.0	43.6	Foliation	65
						143.0	43.6	Vein (5mm) of pure fluorite. 180/90.	
						144.0	43.9	Foliation	55
						145.0	44.2	Vein (1 cm) of fluorite and vein (1.5 cm) of calcite parallel to foliation.	55
						177.0	54.0	Foliation	65
						177.0	54.0	Vein (1 mm) of pyrite. Horizontal	
						185.0	56.4	Foliation	70

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						189.0	57.6	Foliation	75
						193.0	58.8	Vein-fracture with kaolinite coating. 150/90.	
						200.0	61.0	Foliation	70
						212.0	64.6	Foliation	90
						217.0	66.1	Foliation	90
						223.0	68.0	Foliation	70
						224.0	68.3	Foliation	80
						229.0	69.8	Foliation	80
						229.0	69.8	Vein (2 mm) with slickensides. Vein: 065/77NW; slick: 65 ->285.	
						237.0	72.2	Foliation	45
						241.0	73.5	Foliation	90
						245.0	74.7	Foliation	80
						247.0	75.3	Foliation	65
						247.0	75.3	Vein (2 mm) of gypsum 015/90; slicks: 45 ->195.	
						254.0	77.4	Foliation	50
						263.0	80.2	Foliation	50
						270.0	82.3	Foliation	70
						270.0	82.3	Vein-fracture (1-2 mm) with gypsum. 055/75NW.	
						272.0	82.9	Foliation	40
						271.0	82.6	Vein (1 mm) of pyroxene (?). 180/55W.	
278.0		84.7			Medium grained granodiorite gneiss in contact with pink coarse grained muscovite gneiss at 283' (86.3 m) Granodiorite extends upward for 25 more feet (i.e. 253' - 77.1 m at least).	279.0	85.0	Vein (2 mm) of K-feldspar. 055/40SE.	
						283.0	86.3	Foliation	68
						291.0	88.7	Foliation	45
						292.0	89.0	Foliation	55
						298.0	90.8	Disseminated fluorite parallel to foliation.	50
						313.0	95.4	Foliation	45
						313.0	95.4	Vein (2 mm) of quartz. 055/45SE.	
						323.0	98.5	Foliation	55
						327.0	99.7	Foliation	45
						327.0	99.7	Vein (2 mm) of quartz. 055/40SE.	
						327.0	99.7	Vein (3 mm) of gypsum with slicks. 340/50SW; slicks: 278/53NW.	



<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						330.0	100.6	Disseminated fluorite parallel to foliation.	45
						346.0	105.5	Foliation	40
						371.0	113.1	Vein (10 cm) of fluorite and quartz parallel to foliation.	50
						380.0	115.8	Foliation	45
						398.0	121.3	Foliation	60
						414.0	126.2	Foliation	50
						419.0	127.7	Disseminated fluorite parallel to foliation.	50
						432.0	131.7	Foliation	60
						467.0	142.3	Foliation	60
						490.0	149.4	Foliation	55
						507.0	154.5	Foliation	30
						516.0	157.3	Foliation	45
						529.0	161.2	Foliation	35
						534.0	162.8	Foliation	65
						534.0	162.8	Vein (2 mm) of quartz. 055/30SE.	
						538.0	164.0	Vein (2 mm) of gypsum. 078/52SE; slicks: 50 W.	
						538.0	164.0	Vein (2 mm) of gypsum. 055/40SE.	
						541.0	164.9	Foliation	90
						550.0	167.6	Foliation	40
						550.0	167.6	Fault.	
	569.9		173.7		End of hole.				

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
					Eaglet Mines Ltd.				
<b>Project</b>	<b>Client</b> Structural analysis							<b>Collar Elevation</b>	799.84
<b>Drill hole</b>	S9-81							<b>Inclination</b>	90
<b>Coordinates</b>	264.1 m South							<b>Length</b>	111.9 m
	491.7 m West							<b>Logged by:</b>	P.B. Read

Note: Assumed strike of foliation ~ 054°. Core was completely split.

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
0.0	27.0	0.0	8.2		Overburden.				
27.0		8.2			Pink fluorite bearing aplite with cream medium grained, 2 mm zone of muscovite gneiss.	29.0	8.8	Foliation	20
						46.0	14.0	Foliation	55
						68.0	20.7	Foliation	65
						87.0	26.5	Foliation	35
						96.0	29.3	Foliation	55
						102.0	31.1	Foliation	40
						121.0	36.9	Foliation	60
						131.0	39.9	Vein (5 mm) of fluorite and K-feldspar parallel to foliation.	65
						137.0	41.8	Foliation	70
						137.5	41.9	Vein (10 cm) of fluorite and K-feldspar. 015/40SE.	70
						140.0	42.7	Foliation	70
						151.0	46.0	Disseminated fluorite over 20 cm. Parallel to foliation.	40
						160.0	48.8	Foliation	55
						171.0	52.1	Foliation	40
						171.0	52.1	Vein (2 mm) of quartz. 083/45 SE.	
						179.0	54.6	Disseminated fluorite and pyroxene parallel to foliation.	45
						186.0	56.7	Foliation	40
						186.0	56.7	Vein (1 cm) of fluorite and K-feldspar. 054/50SE.	
						190.0	57.9	Disseminated fluorite parallel to foliation.	55
						206.0	62.8	Foliation	80
						208.0	63.4	Vein (3 cm) of fluorite and K-feldspar parallel to foliation.	50
						217.0	66.1	Foliation	55
						230.0	70.1	Foliation	50
						230.0	70.1	Vein (1-2 mm) of fluorite and pyroxene. 054/70SE.	

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						242.0	73.8	Disseminated fluorite and K-feldspar parallel to foliation.	55
						253.0	77.1	Foliation	25
						269.0	82.0	Disseminated fluorite, quartz and K-feldspar parallel to foliation.	50
						272.0	82.9	Foliation	60
						272.0	82.9	Vein of fluorite and K-feldspar. 090/60N.	
						275.0	83.8	Disseminated fluorite parallel to foliation.	45
						282.0	86.0	Foliation	55
						285.0	86.9	Foliation	45
						293.0	89.3	Foliation	40
						297.0	90.5	Foliation	45
						300.0	91.4	Foliation	55
						303.0	92.4	Foliation	45
						303.0	92.4	Vein of fluorite, pyrite and K-feldspar. 335/vertical.	
						328.0	100.0	Foliation	45
						330.0	100.6	Foliation	45
						330.0	100.6	Vein (2 mm) with kaolinite. Vein: 045/85SE. Slicks: 60SW.	
						330.0	100.6	Vein of quartz and pyrite. 302/85SW.	
						339.0	103.3	Vein (1 cm) of fluorite, quartz, pyroxene and K-feldspar. Parallel to foliation.	50
						339.0	103.3	Vein (2 mm) of gypsum. 054/60 SE. Slicks: 075/38NE.	
						343.0	104.5	Disseminated pyroxene and fluorite parallel to foliation	55
						348.0	106.1	Foliation	45
						348.0	106.1	Vein (5 mm) of quartz and pyrite. 345/90.	
						356.0	108.5	Vein (1cm) of fluorite parallel to foliation.	55
	366.0		111.6		End of hole.	366.0	111.6	Foliation	55

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>	
				<b>Client</b>	Eaglet Mines Ltd.					
<b>Project</b>	Structural analysis								<b>Collar Elevation</b>	800.9
<b>Drill hole</b>	S10-81								<b>Inclination</b>	90
<b>Coordinates</b>	347.3 m South								<b>Length</b>	60 m
	350.8 m West								<b>Logged by:</b>	JAF

Note: Assumed strike of foliation ~ 054°. Core was completely split.

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
0.0	50.0	0.0	15.2		Overburden				
50.0	70.0	15.2	21.3		Pink to cream coloured aplite gneiss with less than 5% pyroxene. Some muscovite altered zones with disseminated fluorite.	64.0	19.5	Foliation	65
70.0	197.0	21.3	60.0		Medium grained leucogranodiorite gneiss with less than 1% of pyroxene.	74.0	22.6	Foliation	40
						75.0	22.9	Foliation	50
						80.0	24.4	Foliation	45
						87.0	26.5	Vein (2 mm) of gypsum. 150/48 NE.	
						87.0	26.5	Foliation	65
						100.0	30.5	Foliation	50
						115.0	35.1	Foliation	35
						118.0	36.0	Foliation	40
						132.0	40.2	Foliation	40
						138.0	42.1	Foliation	35
						138.0	42.1	Vein (1 cm) of barren quartz. 033/43 SE.	
						138.0	42.1	Vein-fracture filled with quartz and chlorite. 048/43 NW.	
						145.0	44.2	Foliation	60
						147.0	44.8	Vein (3 mm) of biotite and fluorite parallel to foliation.	55
						153.0	46.6	Foliation	60
						153.0	46.6	Fracture coated with chlorite and quartz. 165/90.	
						162.0	49.4	Foliation	50
						167.0	50.9	Foliation. Shows some movement within S1 plane: slicks 58 ->015	65
						171.0	52.1	Foliation	45
						178.0	54.3	Vein (1 cm) of quartz. Parallel to S1	60

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						188.0	57.3	Foliation	40
						194.0	59.1	Foliation	45
						197.0	60.0	Foliation	50
	197.0		60.0		End of hole.	197.0	60.0	Vein (1 mm) of pyrite and chlorite. 055/60 SE..	

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>	
				<b>Client</b>	Eaglet Mines Ltd.					
<b>Project</b>	Structural analysis								<b>Collar Elevation</b>	798.87
<b>Drill hole</b>	S11-81								<b>Inclination</b>	90
<b>Coordinates</b>	301.8 m South								<b>Length</b>	44.8
	433.0 m West								<b>Logged by:</b>	P.B.Read

Note: Assumed strike of foliation ~ 054°. Core was completely split.

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
0.0	36.0	0.0	11.0		Overburden				
36.0	46.0	11.0	14.0		Medium grained (2-3 mm) muscovite gneiss with thin zones of medium grey biotite- muscovite gneiss and cream coloured gneissic aplite.	41.0	12.5	Foliation	45
46.0		14.0			Cream coloured aplite gneiss.	45.0	13.7	Disseminated fluorite parallel to foliation.	55
						58.0	17.7	Foliation	55
						73.0	22.3	Disseminated hematite parallel to foliation.	45
						81.0	24.7	Foliation	45
						97.0	29.6	Foliation	45
						100.0	30.5	Foliation	45
						105.0	32.0	Foliation	40
						110.0	33.5	Disseminated fluorite and K-feldspar parallel to foliation	45
						116.0	35.4	Disseminated fluorite parallel to foliation.	40
						118.0	36.0	Foliation	40
						118.0	36.0	Vein ( 2mm) of quartz and pyrite. 292/70SW.	
						126.0	38.4	Foliation	65
						128.0	39.0	Foliation	45
						128.0	39.0	Vein (1 mm) of quartz. 055.50 SE.	
						138.0	42.1	Disseminated fluorite up to 5 mm thick parallel to foliation.	45
						140.0	42.7	Foliation	40
	147.0		44.8		End of hole.				

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>	
				<b>Client</b>	Eaglet Mines Ltd.					
<b>Project</b>	Structural analysis								<b>Collar Elevation</b>	765.8
<b>Drill hole</b>	S12-81								<b>Inclination</b>	90
<b>Coordinates</b>	347.7m South								<b>Length</b>	96.6
	580.6 m West								<b>Logged by:</b>	JAF

Note: Assumed strike of foliation ~ 054°. Core was completely split.

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
0.0	14.0	0.0	4.3		Overburden				
14.0	317.0	4.3	96.6		Pink fluorite-bearing aplite with minor cream coloured medium grained (2 mm) muscovite gneiss zones.	14.0	4.3	Minor disseminated fluorite and quartz parallel to foliation.	40
						16.0	4.9	Vein (1-2 mm) of pyrite and fluorite parallel to foliation	50
						27.0	8.2	Foliation	50
						27.0	8.2	Two veins (3-4 mm) of fluorite and quartz. 060/52 SE.	
						32.0	9.8	Foliation	50
						46.0	14.0	Foliation	50
						55.0	16.8	Foliation and vein ( 4 cm) of fluorite and minor quartz parallel to S1	45
						67.0	20.4	Foliation	50
						72.0	21.9	Foliation	55
						72.0	21.9	Vein (2 mm) of quartz. 170/58W.	
						84.0	25.6	Foliation	60
						93.0	28.3	Foliation	50
						93.0	28.3	Fracture- filled with quartz and minor kaolinite. 175/65 E.	
						107.0	32.6	Foliation	65
						114.0	34.7	Foliation	55
						124.0	37.8	Foliation with minor disseminated fluorite parallel to S1.	45
						134.0	40.8	Foliation	45
						149.0	45.4	Foliation	55
						167.0	50.9	Zones of disseminated fluorite a few millimetre wide. Parallel to foliation	65
						176.0	53.6	Foliation	80
						176.0	53.6	Vein(4-5 mm) of pyroxene (99%) with minor fluorite. 070/60 SE	45

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						182.0	55.5	Zones of pyroxene, fluorite and quartz approximately 3 cm wide. Parallel to foliation	45
						197.0	60.0	Foliation	30
						202.0	61.6	Foliation	50
						215.0	65.5	Foliation	50
						223.0	68.0	Foliation	60
						223.0	68.0	Fracture- filled with kaolinite (2mm). 065/90.	60
						225.0	68.6	Vein (1 cm) of fluorite parallel to foliation.	65
						244.0	74.4	Foliation	50
						244.0	74.4	Vein (2 mm) pyrite and quartz. 045/58SE.	
						247.0	75.3	Foliation	35
						267.0	81.4	Foliation	90
						270.0	82.3	Vein (2 mm) of pyroxene, chlorite and pyrite. Parallel to foliation.	90
						280.0	85.3	Foliation	60
						280.0	85.3	Vein (2 mm) of chlorite, quartz and minor pyrite. 055/88 SE with vertical slickensides.	
						291.0	88.7	Foliation	60
						295.0	89.9	Foliation	60
						302.0	92.1	Vein (1.5 cm) of fluorite and quartz parallel to foliation.	65
						312.0	95.1	Foliation	65
						316.0	96.3	Foliation	65
	317		96.6		End of hole.				



<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
					Eaglet Mines Ltd.				
<b>Project</b>	Structural analysis							<b>Collar Elevation</b>	766.74 m
<b>Drill hole</b>	S13-81							<b>Inclination</b>	90
<b>Coordinates</b>	372.2m South							<b>Length</b>	69.1
	550.0 m West							<b>Logged by:</b>	JAF

Note: Assumed strike of foliation ~ 054°. Core was completely split.

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
0.0	34.0	0.0	10.4		Overburden				
34.0	227.0	10.4	69.2		Fine grained (1 mm) white to pink aplite with thin zones of cream coloured medium grained (2 mm) muscovite gneiss.	37.0	11.3	Thin and discontinuous zones of disseminated fluorite parallel to foliation.	35
						55.0	16.8	Foliation	55
						55.0	16.8	Vein (1 mm) of pyrite. 135/60 SW.	
						60.0	18.3	Foliation	45
						72.0	21.9	Foliation	45
						72.0	21.9	Discontinuous vein (0-3 mm) of fluorite, pyrite and quartz. 055/40 SE.	
						92.0	28.0	Foliation	50
						92.0	28.0	Vein (1 cm) of fluorite. 100/78 N.	
						104.0	31.7	Foliation	45
						107.0	32.6	Foliation	50
						107.0	32.6	Fracture-filled vein (1-2 mm) of kaolinite. 030/54 SE.	
						111.0	33.8	Foliation	90
						127.0	38.7	Foliation with local disseminated fluorite parallel to S1.	60
						137.0	41.8	Foliation	45
						145.0	44.2	Foliation with discontinuous vein (1-1.5 cm) of fluorite parallel to S1.	45
						153.0	46.6	Vein (5 mm) of pyroxene, fluorite and quartz parallel to foliation.	60
						157.0	47.9	Vein (2 cm) of barren quartz. Parallel to foliation.	70
						172.0	52.4	Foliation with disseminated molybdenum patches in zone of 2 mm. Parallel to S1.	45
						177.0	54.0	Foliation	40
						188.0	57.3	Foliation	40
						196.0	59.7	Foliation	50
						198.0	60.4	Foliation	50

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						198.0	60.4	Vein (3-4 mm) of quartz and pyrite. 040/30SE.	
						203.0	61.9	Foliation with parallel zone (4 mm) of fluorite and pyroxene	60
						219.0	66.8	Foliation	75
						219.0	66.8	Vein, kaolinite in fracture (1-2 mm). 055/90	
						226.0	68.9	Foliation	70
	227.0		69.2		End of hole.				

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
				<b>Client</b> Eaglet Mines Ltd.					
<b>Project</b>	Structural analysis						<b>Collar Elevation</b>	841.31 m	
<b>Drill hole</b>	S17-81						<b>Inclination</b>	90	
<b>Coordinates</b>	172.5 m South						<b>Length</b>	130.2	
	421.9 m West						<b>Logged by:</b>	JAF	

Note: Assumed strike of foliation ~ 090°. Core is completely split.

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
0.0	44.0	0.0	13.4		Overburden				
44.0	76.0	13.4	23.2	QLG	Pink aplite and muscovite-aplite gneiss with lesser granodiorite gneiss. Alteration at the top of the hole is gypsum and pyroxene. Pyroxene is reddish weathering (+pyrite?). Disseminated fluorite and local veins of fluorite common (disseminated parallel to S1).			First appearance of gypsum and pyroxene(?)	
76.0	86.0	23.2	26.2		Pink aplite and muscovite aplite. Granodiorite gneiss absent. Fine grained pyroxene (1-2%) in aplite.	52.0	15.8	Foliation	50
						56.0	17.1	Foliation	55
86.0	97.0	26.2	29.6		Pink aplite, muscovite-aplite gneiss and medium grained granodiorite gneiss (pyroxene less than 4-5%).	65.0	19.8	Foliation	50
97.0	130.0	29.6	39.6		Pink aplite, muscovite aplite gneiss and granodiorite gneiss (10%). Kfeldspar veins present in this section.	65.0	19.8	Vein (1 mm) of gypsum in several closely spaced fractures. 095/45 S.	
130.0	160.0	39.6	48.8		Pyroxene (5%) granodiorite gneiss with minor aplite gneiss. Well foliated, abundant quartz.	71.0	21.6	Foliation	60
160.0	196.0	48.8	59.7		Pink aplite, muscovite aplite gneiss. Some minor granodiorite gneiss (5%).	75.0	22.9	Foliation	50
						80.0	24.4	Foliation	45
						86.0	26.2	Foliation	45

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
196.0	230.0	59.7	70.1		Granodiorite gneiss with pyroxene (5%); local aplitic gneiss. Gypsum alteration common. At 228' (69.5 m) white aplite is present within granodiorite and again at 231' (70.4 m) .	86.0	26.2	Two veins (2 mm) of barren quartz. 085/40S.	
230.0	295.0	70.1	89.9		Pink aplite with local muscovite-bearing aplitic gneiss. Quartz + pyrite + fluorite veins common; locally with calcite. Large (1-1.5 cm across) pink K-feldspar "augen" present in strongly altered K-feldspar-fluorite ore. Fine grained biotite also concentrated in clots, locally.	90.0	27.4	Foliation	55
						97.0	29.6	Foliation	65
						114.0	34.7	Foliation	60
						114.0	34.7	Vein (5-6 mm) of fluorite and quartz. 075/55 NW.	
295.0	300.0	89.9	91.4		Medium grained pyroxene granodiorite (pyroxene 2%).	120.0	36.6	Foliation	75
300.0	327.0	91.4	99.7		Pink aplite and muscovite-aplite gneiss with local fluorite-bearing white aplite.	127.0	38.7	Foliation	50
327.0	348.0	99.7	106.1		Granodiorite gneiss with white aplite. Fluorite at contact between aplite and gneiss.	131.0	39.9	Foliation	45
						142.0	43.3	Foliation	65
348.0	365.0	106.1	111.3		Granodiorite gneiss with increasing aplite. Aplite: 25-30%.	150.0	45.7	Foliation	55
						152.0	46.3	Foliation	45
365.0	405.0	111.3	123.4		Granodiorite gneiss with minor aplite (less than 5%). Pyroxene locally up to 20% (fine grained masses, locally large crystals 5 mm across).	152.0	46.3	Vein of quartz with gypsum coating. 030/68 NW; slicks: 60->320°.	
405.0	427.0	123.4	130.2		Intense pyroxene alteration from 405' (123.5 m) to 412' (125.6 m). Green pyroxene with crystal up to 5 mm common; parallel to S1.	157.0	47.9	Foliation	60
						160.0	48.8	Foliation	65
						169.0	51.5	Foliation	70

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						169.0	51.5	Vein (2 mm) of K- feldspar, pyrite, galena, quartz and minor calcite. 080/32 SE.	
						172.0	52.4	Foliation	60
						174.0	53.0	Sample of aplite with hornblende (?)	
						180.0	54.9	Foliation	70
						181.0	55.2	Foliation	65
						185.0	56.4	Foliation	60
						187.0	57.0	Foliation	55
						187.0	57.0	Vein (3 mm) of K-feldspar and pyrite. 140/75 NE.	
						191.0	58.2	Foliation	60
						187.0	57.0	Foliation	50
						197.0	60.0	Vein-fracture (1 mm) of gypsum. 020/76 SE.	
						199.0	60.7	Foliation	90
						203.0	61.9	Disseminated fluorite and fine grained pyroxene parallel to foliation.	
						228.0	69.5	Foliation	75
						230.0	70.1	Vein (4 mm) of pyrite and fluorite parallel to foliation.	70
						234.0	71.3	Foliation	65
						257.0	78.3	Foliation	30
						263.0	80.2	Foliation	50
						274.0	83.5	Foliation	45
						277.0	84.4	Foliation with concentrated green pyroxene parallel to S1; 4-5 mm wide.	45
						291.0	88.7	Foliation	60
						293.0	89.3	Foliation	45
						293.0	89.3	Vein (2 mm) of pyrite and K-feldspar. 040/15 SE.	
						297.0	90.5	Foliation	40
						300.0	91.4	Foliation	40
						305.0	93.0	Foliation	30
						305.0	93.0	Vein (2 mm) of k-feldspar, calcite, pyrite and minor fluorite. 062/47 SE.	
						317.0	96.6	Foliation	45
						317.0	96.6	Vein (3 mm) of K-feldspar, pyroxene and pyrite. 038/28 SE.	
						319.0	97.2	Foliation	40
						321.0	97.8	Foliation	65

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						327.0	99.7	Foliation	30
						327.0	99.7	Vein (3 mm) of gypsum with strong slicks. Vein: 048/78 SE; slicks: down dip.	
						335.0	102.1	Foliation	15
						336.0	102.4	Foliation	40
						342.0	104.2	Foliation	45
						342.0	104.2	Vein (2 mm) of gypsum with slicks. Vein: 005/80E; slicks: down dip.	
						345.0	105.2	Foliation	40
						350.0	106.7	Foliation	45
						354.0	107.9	Foliation	45
						354.0	107.9	Two veins (1-1.5 cm) with barren quartz. 073/68 SE.	
						359.0	109.4	Foliation	40
						365.0	111.3	Foliation	45
						369.0	112.5	Foliation	40
						374.0	114.0	Foliation	45
						382.0	116.4	Vein (1.5 cm) of calcite and biotite. Parallel to foliation	45
						386.0	117.7	Foliation	45
						386.0	117.7	Vein (6-7 mm) of pyroxene, calcite and minor fluorite. 015/50 NW.	
						390.0	118.9	Foliation	40
						395.0	120.4	Foliation	40
						400.0	121.9	Foliation	40
						406.0	123.8	Vein of coarse grained green pyroxene (4-5 mm). Parallel to foliation.	45
						411.0	125.3	Sample: Biotite and calcite abundant locally and associated with pyroxenes.	
						416.0	126.8	Foliation	45
						425.0	129.5	Foliation	40
						426.0	129.8	Sample: Biotite and calcite abundant locally and associated with pyroxenes.	
	427.0		130.2		End of hole.				

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>	
				<b>Client</b>	Eaglet Mines Ltd.					
<b>Project</b>	Structural analysis								<b>Collar Elevation</b>	?
<b>Drill hole</b>	S18-81								<b>Inclination</b>	90
<b>Coordinates</b>	?								<b>Length</b>	145 m
	?								<b>Logged by:</b>	P.B.Read

Note: Assumed strike of foliation ~ 090°.

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
0.0	75.0	0.0	22.9		Overburden				
75.0	124.0	22.9	37.8		White fine grained (1 mm) gneissic aplite.	98.0	29.9	First appearance of fluorite	
						105.0	32.0	First appearance of pyroxene. Sample S18-81-105.	
						108.0	32.9	First appearance of coarse pyrite.	
						123.0	37.5	First appearance of galena and K-feldspar veins.	
124.0		37.8			Medium grained muscovite gneiss.	130.0	39.6	Foliation	65
	475.7		145.0		End of hole.				

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>	
				<b>Client</b>	Eaglet Mines Ltd.					
<b>Project</b>	Structural analysis								<b>Collar Elevation</b>	880.54 m
<b>Drill hole</b>	S19-81								<b>Inclination</b>	90
<b>Coordinates</b>	191.4 m South								<b>Length</b>	135.6 m
	255.4 m West								<b>Logged by:</b>	JAF

**Note: Assumed strike of foliation ~ 054. Core was completely split.**

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
0.0	15.0	0.0	4.6		Overburden				
15.0	325.0	4.6	99.1	QLG	Granodiorite gneiss. Medium grained pyroxene + biotite (20%) + K-feldspar and quartz.	17.0	5.2	Foliation	45
						17.0	5.2	Vein (1 cm) of fluorite and pyrite. 040/20 NW.	
						29.0	8.8	Vein (2-3 mm) of fluorite and quartz. Parallel to foliation.	40
						36.0	11.0	Foliation	35
						36.0	11.0	Vein (2 mm) of fluorite, pyrite and sphalerite. 170/70 E.	
						53.0	16.2	Foliation	35
						53.0	16.2	Vein (4 mm) of quartz and fluorite. 115/48 NE.	
						61.0	18.6	Foliation	35
						61.0	18.6	Vein (3 mm) of pyrite and lesser fluorite. 010/65E	
						68.0	20.7	Foliation	30
						68.0	20.7	Vein (1 mm) of quartz and k-feldspar. 065/65 SE.	
						73.0	22.3	Foliation	45
						73.0	22.3	Vein (0.3-10 mm) of fluorite, galena and quartz. 000/69E.	
						75.0	22.9	Foliation	40
						75.0	22.9	Vein (5 mm) of quartz with minor pyrite. 005/75E.	
						79.0	24.1	Foliation	35
						79.0	24.1	Fracture-coated with chlorite and quartz. Slicks:17->325. Fracture: 075/45 SE.	
						87.0	26.5	Foliation	45
						87.0	26.5	Vein (5 mm) of quartz, K-feldspar and fluorite. 075/45 SE.	
						94.0	28.7	Foliation	45



<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						102.0	31.1	Foliation with disseminated pyrite and fluorite parallel to S1.	50
						109.0	33.2	Foliation	45
						121.0	36.9	Foliation	60
						130.0	39.6	Foliation	50
						131.0	39.9	Foliation	45
						151.0	46.0	Foliation	40
						158.0	48.2	Foliation	40
						185.0	56.4	Foliation	40
						211.0	64.3	Foliation	25
						227.0	69.2	Foliation	30
						245.0	74.7	Foliation	25
						245.0	74.7	Vein -fracture (2-3 mm) with kaolinite. 110/70 SW.	
						257.0	78.3	Foliation	40
						257.0	78.3	Vein (2-3 mm) of kaolinite (gypsum?). 105/70 SW.	
						260.0	79.2	Foliation	30
						260.0	79.2	Vein (1 mm) of kaolinite. 040/90.	
						270.0	82.3	Foliation	30
						270.0	82.3	Vein (7-8 mm) of barren quartz. 090/32 N.	
						283.0	86.3	Disseminated fluorite parallel to foliation.	45
						283.8	86.5	Vein (1 mm) of quartz and kaolinite (?). 103/67SW.	
						288.0	87.8	Vein (1 mm) of gypsum parallel to foliation. Slickensides: 37-> 320.	40
						295.0	89.9	Foliation	45
						295.0	89.9	Vein (2 mm) of barren quartz. 055/45 SE.	
						302.0	92.1	Foliation	20
						307.0	93.6	Foliation	35
						307.0	93.6	Vein (2-3 mm) of quartz and pyrite. 045/78 SE.	
						312.0	95.1	Foliation	30
						312.0	95.1	Vein (3 -4 ) of barren quartz. 030/45 NW.	
						321.0	97.8	Foliation	35
						350.0	106.7	Foliation	50
						357.0	108.8	Foliation	35
						357.0	108.8	Vein-fracture (1 mm) of quartz. 040/30 SE.	
						360.0	109.7	Foliation	50

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						360.0	109.7	Vein- gypsum filled-fracture; Slickensides: 45->085; Vein: 155/53NE.	
						366.0	111.6	Foliation	45
						372.0	113.4	Foliation	45
						377.0	114.9	Foliation	60
						382.0	116.4	Foliation	50
						392.0	119.5	Foliation	45
						398.0	121.3	Foliation	55
						402.0	122.5	Foliation	65
						402.0	122.5	Vein (3 mm) of barren quartz. 088/46 S.	
						408.0	124.4	Foliation	50
						413.0	125.9	Foliation with 5 mm wide fluorite vein parallel to S1.	
						417.0	127.1	Foliation	50
						422.0	128.6	Foliation	55
						428.0	130.5	Foliation	50
						432.0	131.7	Vein (1.5 cm) of fluorite and pyrite parallel to foliation	45
						439.0	133.8	Foliation	45
	445.0		135.6		End of hole.	445.0	135.6	Foliation	55

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
				Eaglet Mines Ltd.					
<b>Project</b>	<b>Client</b>							<b>Collar Elevation</b>	824.03
<b>Drill hole</b>	Structural analysis							<b>Inclination</b>	90
<b>Coordinates</b>	S20-81							<b>Length</b>	66.1 m
	348.2 m South							<b>Logged by:</b>	P.B. Read
	384.7 m West								

**Note: Assumed strike of foliation ~ 054. Core was completely split.**

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
0.0	42.0	0.0	12.8		Overburden				
42.0	217.0	12.8	66.1		Medium grained (1-3 mm) biotite (3%) leucogranodiorite gneiss.	46.0	14.0	Foliation	60
						49.0	14.9	Foliation	40
						65.0	19.8	Foliation	52
						71.0	21.6	Foliation	51
						77.0	23.5	Foliation parallel to contact between overlying pink muscovite gneiss and underlying pink aplite. Disseminated fluorite.	50
						80.0	24.4	Foliation parallel to contact between overlying pink aplite and underlying muscovite gneiss. Vein (1 mm) of gypsum. 340/90.	50
						84.0	25.6	Disseminated fluorite parallel to foliation.	58
						104.0	31.7	Foliation	30
						129.0	39.3	Foliation	63
						132.0	40.2	Disseminated fluorite parallel to foliation.	65
						132.0	40.2	Vein (1 mm) of gypsum. 275/42 SW	
						140.0	42.7	Foliation	60
						147.0	44.8	Foliation	15
						147.0	44.8	Vein (2 mm) of quartz. 054/90.	
						149.0	45.4	Foliation	50
						158.0	48.2	Foliation	60
						171.0	52.1	Foliation	90
						176.0	53.6	Disseminated fluorite parallel to foliation.	50
						181.0	55.2	Foliation	52
						182.0	55.5	Foliation	55
						182.0	55.5	Vein (2 mm). 075/78SE.	
						183.0	55.8	Foliation	45

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						183.0	55.8	Vein (5 mm) of quartz and pyrite. 054/41 SE.	
						192.0	58.5	Foliation	40
						194.0	59.1	Disseminated fluorite parallel to foliation.	30
						196.0	59.7	Foliation	50
						196.0	59.7	Vein (3 mm) of quartz. 035/52 SE.	
						197.0	60.0	Disseminated fluorite parallel to foliation.	45
						205.0	62.5	Foliation	30
						213.0	64.9	Foliation	60
						213.0	64.9	Vein (3 mm) of quartz and fluorite. 054/35 NW.	
						216.0	65.8	Foliation	40
						216.0	65.8	Vein (3 mm) of quartz. 340/67 SW.	
	217.0		66.1		End of hole.				

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
					Eaglet Mines Ltd.				
<b>Project</b>	<b>Client</b> Structural analysis							<b>Collar Elevation</b>	825.31 M
<b>Drill hole</b>	S21-81							<b>Inclination</b>	90
<b>Coordinates</b>	264.3 m South							<b>Length</b>	38.7 M
	344.2 m West							<b>Logged by:</b>	P.B. Read

**Note: Assumed strike of foliation ~ 054. Core was completely split.**

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
0.0	52.0	0.0	15.8		Overburden				
52.0	127.0	15.8	38.7		Medium grained (1-3 mm) pink biotite (10%) granodiorite gneiss.	54.0	16.5	Foliation	40
						62.0	18.9	Disseminated fluorite parallel to foliation.	45
						78.0	23.8	Vein (3 mm) of quartz parallel to foliation	52
						83.0	25.3	Foliation	63
						83.0	25.3	Vein (3 mm) of gypsum. 054/42 SE.	
						86.0	26.2	Fault	
						97.0	29.6	Foliation	39
						103.5	31.5	Vein (2 mm) of quartz parallel to foliation	40
						104.0	31.7	Vein (2 mm) of quartz and fluorite. 030/33 SE.	
						107.0	32.6	Foliation	30
						107.0	32.6	Vein (3 mm) of K-feldspar. 065/63 SE.	
						109.0	33.2	Foliation	25
						109.0	33.2	Vein (2 mm) of quartz and fluorite. 054/71 SE.	
						112.0	34.1	Vein (8 cm) of quartz and K-feldspar parallel to foliation.	30
						117.0	35.7	Vein (2 cm) of K-feldspar, fluorite and molybdenum.	45
						122.5	37.3	Disseminated fluorite parallel to foliation. 5 mm thick.	32
						125.0	38.1	Foliation	52
						125.0	38.1	Vein (1 mm) of quartz. 054/61 SE.	
	127.0		38.7		End of hole.				

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>	
				<b>Client</b>	Eaglet Mines Ltd.					
<b>Project</b>	Structural analysis								<b>Collar Elevation</b>	818.43 m
<b>Drill hole</b>	S22-81								<b>Inclination</b>	90
<b>Coordinates</b>	268.2 m South								<b>Length</b>	41.8 m
	376.7 m West								<b>Logged by:</b>	P.B. Read

**Note: Assumed strike of foliation ~ 054. Core was completely split.**

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
0.0	35.0	0.0	10.7		Overburden				
35.0	137.0	10.7	41.8		White fluorite-bearing aplite with pink muscovite gneiss layers.	49.0	14.9	Foliation	70
						62.0	18.9	Foliation	66
						65.0	19.8	Foliation	58
						65.0	19.8	Vein (1 mm) of quartz. 042/52 SE.	
						70.0	21.3	Disseminated fluorite parallel to foliation.	70
						95.0	29.0	Foliation	60
						102.0	31.1	Foliation	55
						102.0	31.1	Vein (3 mm) of gypsum. 010/75 SE.	
						105.0	32.0	Foliation	30
						109.0	33.2	Disseminated pyroxene and fluorite parallel to foliation.	55
						113.0	34.4	Vein (1 cm) of quartz and pyroxene parallel to foliation.	35
						118.0	36.0	Foliation	40
						118.0	36.0	Vein (3 mm) of calcite and pyrite. 054/48 SE.	
						123.0	37.5	Disseminated fluorite and pyroxene parallel to foliation.	30
						128.0	39.0	Foliation	30
						132.0	40.2	Disseminated fluorite and pyroxene parallel to foliation.	30
						135.0	41.1	Vein (1 cm) of quartz and fluorite parallel to foliation.	45
	137.0		41.8		End of hole.				

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
<b>Project</b>	<b>Client</b>				Eaglet Mines Ltd.				
	Structural analysis							<b>Collar Elevation</b>	806.6 m
<b>Drill hole</b>	S23-81							<b>Inclination</b>	90
<b>Coordinates</b>	276.1 m South							<b>Length</b>	32.6 m
	387.2 m West							<b>Logged by:</b>	JAF

**Note: Assumed strike of foliation ~ 054. Core was completely split.**

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
0.0	50.0	0.0	15.2		Overburden				
50.0	107.0	15.2	32.6	QLG	Pink aplitic gneiss with K-feldspar augen, disseminated fluorite. Fine grained (1-3 mm) leucocratic.	57.0	17.4	Disseminated fluorite parallel to foliation. 5 mm thick.	50
						71.0	21.6	Vein (4 mm) of fluorite parallel to foliation	45
						76.0	23.2	Foliation	50
						78.0	23.8	Foliation	65
						78.0	23.8	Vein (3 mm) of barren quartz. 055/65 SE	
						90.0	27.4	Foliation	50
						96.0	29.3	Foliation	55
						96.0	29.3	Vein (3-4 mm) of barren quartz. 080/62 SE	
						105.0	32.0	Foliation	70
						105.0	32.0	Vein (3-4 mm) of quartz and fluorite. 095/25 N.	
	107.0		32.6		End of hole.	107.0	32.6	Foliation	45

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
					Eaglet Mines Ltd.				
<b>Project</b>	<b>Client</b>							<b>Collar Elevation</b>	778.39 m
<b>Drill hole</b>	Structural analysis							<b>Inclination</b>	90
<b>Coordinates</b>	S24-81							<b>Length</b>	170.0 m
	247 m South							<b>Logged by:</b>	JAF
	692 m West								

**Note: Assumed strike of foliation ~ 054. Core was completely split.**

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
0.0	26.0	0.0	7.9		Overburden				
26.0	42.0	7.9	12.8		Cream coloured, medium grained (2-3 mm) muscovite gneiss.	28.0	8.5	Foliation	50
42.0	557.7	12.8	170.0		Medium grained (2-4 mm) pyroxene (5%) leucogranodiorite gneiss.	49.0	14.9	Foliation	55
						54.0	16.5	Foliation	50
						70.0	21.3	Foliation	50
						80.0	24.4	Foliation	50
						80.0	24.4	Vein (4 mm) of quartz and fluorite. 138/37 NE.	
						92.0	28.0	Vein (5 mm) of fluorite parallel to foliation.	60
						97.0	29.6	Foliation with fluorite vein (1 cm) parallel to S1	60
						97.0	29.6	Fracture filled with kaolinite. 040/70 SE.	
						112.0	34.1	Foliation	60
						112.0	34.1	Vein (5mm) of quartz and pyrite. 055/30 SE.	
						118.0	36.0	Foliation	50
						118.0	36.0	Vein (8 cm) of quartz (90%), pyrite, sphalerite, galena and fluorite. 085/27 N.	
						121.0	36.9	Foliation	65
						121.0	36.9	Vein (5 mm) of fluorite and K-feldspar. 165/54 SW.	
						127.0	38.7	Vein of pyrite and fluorite parallel to foliation	55
						150.0	45.7	Vein (1 cm) of disseminated fluorite and pyrite parallel to foliation.	55
						150.0	45.7	Vein (1-2 mm) of fluorite and pyrite. 055/35 SE.	



<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						160.0	48.8	Foliation	60
						160.0	48.8	Vein (1.5-2 cm) of sphalerite, pyrite, galena and fluorite. 075/42 SE.	
						169.0	51.5	Foliation	50
						187.0	57.0	Foliation	60
						190.0	57.9	Vein (4 mm) of fluorite and quartz parallel to foliation.	50
						200.0	61.0	Foliation	70
						211.0	64.3	Foliation	60
						217.0	66.1	Foliation	65
						217.0	66.1	Fracture filled with kaolinite. 040/63 SE.	
						232.0	70.7	Foliation	65
						240.0	73.2	Foliation	60
						242.0	73.8	Foliation	60
						242.0	73.8	Vein (3 mm) of fluorite, quartz and pyrite. 055/20 SE.	
						251.0	76.5	Vein (1-2 mm) of pyrite and fluorite parallel to foliation.	50
						258.0	78.6	Foliation	50
						277.0	84.4	Foliation	60
						277.0	84.4	Vein-fracture (2 mm) filled with kaolinite. 055/50 SE.	
						280.0	85.3	Foliation	60
						286.0	87.2	Foliation	60
						286.0	87.2	Vein-fracture filled with kaolinite. 140/78 SW.	
						295.0	89.9	Foliation	60
						307.0	93.6	Foliation	50
						307.0	93.6	Vein (2 mm) of fluorite and pyrite. 055/20 SE.	
						319.0	97.2	Foliation	50
						331.0	100.9	Foliation	55
						343.0	104.5	Vein (1 cm ) of pyroxene, fluorite and pyrite parallel to foliation.	60
						359.0	109.4	Foliation	55
						359.0	109.4	Vein-fracture (1 mm) of fluorite. 010/27 E.	
						369.0	112.5	Disseminated pyrite and fluorite parallel to foliation.	45
						373.0	113.7	Foliation	45
						373.0	113.7	Vein-fracture filled with kaolinite. 060/50 NW.	
						410.0	125.0	Foliation	60

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						418.0	127.4	Foliation	55
						439.0	133.8	Foliation	85
						442.0	134.7	Foliation	90
						460.0	140.2	Vein (1 cm) of fluorite, pyrite and K-feldspar parallel to foliation.	40
						471.0	143.6	Foliation	70
						478.0	145.7	Foliation	65
						478.0	145.7	Vein of kaolinite. 000/65E	
						484.0	147.5	Zone ( 5 cm wide) of disseminated fluorite and pyrite parallel to foliation.	50
						491.0	149.7	Foliation	45
						491.0	149.7	Vein -fracture (1-2 mm) of kaolinite. 158/55NE.	
						507.0	154.5	Foliation	35
						514.0	156.7	Foliation	45
						514.0	156.7	Vein -fracture of kaolinite. 085/66 SE.	
						529.0	161.2	Foliation	30
						541.0	164.9	Foliation	65
						548.0	167.0	Foliation	70
	557.7		170.0		End of hole	557.0	169.8	Foliation	65

**Note: Lower 100' (30.5 m) of core is nearly devoid of fluorite.**

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
<b>Project</b>	<b>Client</b>				Eaglet Mines Ltd.				
	Structural analysis							<b>Collar Elevation</b>	856.8 m
<b>Drill hole</b>	S25-81							<b>Inclination</b>	90
<b>Coordinates</b>	196 m South							<b>Length</b>	80.1 m
	329 m West							<b>Logged by:</b>	JAF

**Note: Assumed strike of entire drill hole ~ 054. Core was completely split.**

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
0.0	63.5	0.0	19.4						
63.5	263.0	19.4	80.2	QLG	Fine grained pink aplite gneiss with local altered (muscovite) gneiss.	68.0	20.7	Foliation	50
						69.0	21.0	Vein (1.5 cm) of fluorite parallel to foliation	45
						78.0	23.8	Vein (5 mm) of fluorite parallel to foliation	30
						81.0	24.7	Foliation	40
						81.0	24.7	Vein (1.5 cm) of fluorite, quartz and pyroxene. 035/60 SE.	
						81.0	24.7	Foliation	30
						87.0	26.5	Foliation	20
						87.0	26.5	Vein (2 mm) of barren quartz. 155/87 SW.	
						94.0	28.7	Foliation with disseminated fluorite parallel to S1	15
						108.0	32.9	Foliation	40
						112.0	34.1	Foliation	35
						124.0	37.8	Foliation	40
						124.0	37.8	Vein (5mm) of quartz and pyrite. 085/90.	
						130.0	39.6	Foliation	60
						134.0	40.8	Foliation	50
						138.0	42.1	Foliation	50
						150.0	45.7	Vein (1-2 mm) of kaolinite. 060/80 SE.	
						155.0	47.2	Foliation	60
						157.0	47.9	Foliation	60
						157.0	47.9	Vein (2 mm) of gypsum. 045/38 SE.	
						161.0	49.1	Foliation	60
						173.0	52.7	Foliation	45
						180.0	54.9	Foliation	50
						203.0	61.9	Foliation	60

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						208.0	63.4	Foliation	45
						208.0	63.4	Vein (2-3 mm) of gypsum. 170/55 W.	
						213.0	64.9	Foliation	40
						229.0	69.8	Foliation	45
						244.0	74.4	Foliation	35
						248.0	75.6	Vein (5 mm) of fluorite parallel to foliation	40
						250.0	76.2	Foliation	40
						250.0	76.2	Vein-fracture (1 mm) of quartz and chlorite. 150/45 NE.	
						259.0	78.9	Foliation	40
						259.0	78.9	Vein ( $\leq$ mm) of quartz and gypsum. 175/86 SW.	
						263.0	80.2	Foliation	60
	263.0		80.2		End of hole	263.0	80.2	Vein (1 cm) of fluorite and quartz. Diffuse margins. 090/70 N.	

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
					Eaglet Mines Ltd.				
<b>Project</b>	<b>Client</b> Structural analysis							<b>Collar Elevation</b>	840.35 m
<b>Drill hole</b>	S26-81							<b>Inclination</b>	90
<b>Coordinates</b>	200 m South							<b>Length</b>	92.7 m
	394 m West							<b>Logged by:</b>	P.B Read

**Note: Assumed strike of foliation ~ 054. Core was completely split.**

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
0.0	40.0	0.0	12.2		Overburden				
4.0	304.0	1.2	92.7		Fine grained (1 mm) cream biotite (2%) aplite gneiss.	72.0	21.9	Foliation	90
						78.0	23.8	Foliation	70
						83.0	25.3	Foliation	70
						96.0	29.3	Vein (3 mm) of fluorite and quartz parallel to foliation.	45
						102.0	31.1	Disseminated fluorite parallel to foliation.	55
						102.0	31.1	Vein (1 mm) of gypsum. 010/62 SE.	
						114.0	34.7	Foliation	55
						123.0	37.5	Foliation	65
						132.0	40.2	Foliation	65
						132.0	40.2	Vein (1 mm) of gypsum. 285/56 SW.	
						152.0	46.3	Foliation	50
						158.0	48.2	Foliation	58
						158.0	48.2	Vein (4 mm) of fluorite 045/55 SE.	
						163.0	49.7	Contact between overlying muscovite gneiss and underlying fluorite-bearing aplite parallel to foliation.	30
						168.0	51.2	Foliation	53
						168.0	51.2	Vein (2 mm) of pyrite, fluorite and quartz. 054/63SE.	
						171.0	52.1	Foliation	45
						171.0	52.1	Vein (1 mm) of quartz. 055/75 SE.	
						182.0	55.5	Disseminated fluorite parallel to foliation.	50
						222.0	67.7	Disseminated fluorite and K-feldspar parallel to foliation.	30
						237.0	72.2	Foliation	40

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						237.0	72.2	Vein (1 mm) of quartz. 035/61 SE.	
						253.0	77.1	Foliation	50
						254.0	77.4	Disseminated fluorite and K-feldspar parallel to foliation.	40
						266.0	81.1	Foliation	50
						272.0	82.9	Foliation	65
						274.0	83.5	Disseminated fluorite parallel to foliation.	52
						282.0	86.0	Foliation	36
						291.0	88.7	Foliation	55
	304.0		92.7		End of hole	304.0	92.7	Foliation	52

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
				<b>Client</b>	Eaglet Mines Ltd.				
<b>Project</b>	Structural analysis							<b>Collar Elevation</b>	927.80 m
<b>Drill hole</b>	S27-81							<b>Inclination</b>	90
<b>Coordinates</b>	61 m South 260 m West							<b>Length</b>	255.1 m
								<b>Logged by:</b>	JAF

**Note: Assumed strike of foliation ~ 090. Hole begins in Snowshoe Group Schist. Unsplit core.**

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
0.0	27.5	0.0	8.4		Overburden				
27.5	226.0	8.4	68.9	Snow-shoe	Feldspar, quartz, biotite, chlorite schist with garnet locally .	31.0	9.4	Foliation	50
						33.0	10.1	Foliation	35
						33.0	10.1	Vein (2 cm) of barren quartz. 180/90.	
						42.0	12.8	Foliation with quartz vein (1 cm) parallel to S1.	30
						52.0	15.8	Foliation	30
						52.0	15.8	Vein (< 5 mm) of barren quartz. 178/66 W.	
						60.0	18.3	Foliation	25
						60.0	18.3	Minor fold- F2 (?); AXP1: 060/35NW; FAX: 10-> 295.	
						69.0	21.0	Minor fold- F2; AXP1: 105/55S; FAX: 10/270.	
					Biotite schist with abundant gamet	80.0	24.4	Foliation	25
						87.0	26.5	Foliation	10
						87.0	26.5	Vein (2 mm) of barren quartz. 180/56 W.	
						92.0	28.0	Foliation	15
						102.0	31.1	Foliation	20
						114.0	34.7	Foliation	
						114.0	34.7	Vein-fracture (1 mm) of kaolinite. 177/55 W.	
						129.0	39.3	Foliation	25
						129.0	39.3	Vein (2-3 mm) of barren quartz. 095/72 S.	
						136.0	41.5	Stringers of pyrite a few millimetres wide parallel to foliation.	30
						157.0	47.9	Foliation	40
						157.0	47.9	Minor fold- F2 (?); AXP1: 100/48N; FAX: 3-> 280.	

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						164.0	50.0	Foliation	45
						164.0	50.0	Vein (2-3 mm) of barren quartz. 118/40 SW.	
						170.0	51.8	Foliation	40
						170.0	51.8	Vein of massive quartz with minor bits of altered chloritized wallrock. Parallel to S1.	45
						178.0	54.3	Foliation	40
					Quartz, feldspar, biotite schist with garnet locally.	178.0	54.3	Minor fold- "Z" symmetry (F3?); AXP1: 103/42SW; FAX: 15-> 280.	
						195.0	59.4	Foliation	30
						195.0	59.4	Crenulations- AXP1: 045/52NW; FAX: 23 ->350.	
						200.0	61.0	Foliation	40
						217.0	66.1	Two quartz veins (1- 1.5 cm) parallel to foliation.	20
						217.0	66.1	Vein (4 mm) of quartz and pyrite. 080/50 SE.	
						221.0	67.4	Foliation	35
						221.0	67.4	Quartz vein (2 mm) cross cut S1 at moderate angle. 155/56 SW.	
226.0	229.0	68.9	69.8	QLG	Fine grained pink aplite with local clusters of pyroxene (5% or less). The contact is sharp and at low angle to S1. More Snowshoe is encountered in the next 10' (3 m) of core. Intrusive contact mostly irregular cross cut compositional layering.	226.0	68.9	Contact- 030/35 NW. Surface has slickensides: 23-> 245.	
						226.0	68.9	Vein (3-4 mm) of pyroxene and fluorite parallel to foliation.	30
						227.0	69.2	Core is split from ~ 227' (69.2 m) and beyond.	
229.0	232.0	69.8	70.7	Snow- shoe - QLG	Small (90 cm) length of biotite schist, then more pink aplite. Mixed contact.	229.0	69.8	Layers up to 1 cm) of K-feldspar rich material parallel to foliation.	45
232.0	675.0	70.7	205.7		Pink aplite gneiss with muscovite altered aplite.	237.0	72.2	Zones of irregular fluorite clusters (2- 3 mm) parallel to foliation.	20
						247.0	75.3	Foliation	35
						257.0	78.3	Foliation with abundant disseminated fluorite parallel to S1.	55
						262.0	79.9	Foliation with localized clusters of fluorite parallel to S1.	45
						277.0	84.4	Foliation	35



<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						277.0	84.4	Four veins (1-2 mm) of fluorite, pyrite and quartz. 045/30 NW.	
						285.0	86.9	Foliation	45
						285.0	86.9	Vein (2 mm) of fluorite and pyrite. 090/37 N.	
						296.0	90.2	Foliation	45
						296.0	90.2	Three veins-fractures (1 mm) with gypsum. 090/42 S.	
						298.0	90.8	Foliation	60
						298.0	90.8	Three veins with quartz and minor fluorite. 090/35 S.	
						302.0	92.1	Zones of disseminated pyrite and fluorite parallel to foliation.	60
						302.0	92.1	Five veins (1 mm) with quartz, pyrite and fluorite. 090/30 S.	
						321.0	97.8	Vein (5 mm) of quartz and fluorite parallel to foliation.	45
						326.0	99.4	Foliation	60
						328.0	100.0	Foliation	50
						328.0	100.0	Vein ( 3 mm) of K-feldspar, quartz and minor fluorite. 055/47 SE.	
						331.0	100.9	Foliation	40
						331.0	100.9	Vein (2 mm) of quartz, fluorite, galena and pyrite. 090/45 S.	
						340.0	103.6	Foliation	55
						340.0	103.6	Vein (4 mm) of K-feldspar, fluorite, galena pyrite and quartz. 115/32 SW.	
						347.0	105.8	Foliation	45
						347.0	105.8	Vein-fracture (2 mm) with gypsum. 127/90.	
						352.0	107.3	Vein (3 mm) of fluorite and pyrite parallel to foliation.	40
						357.0	108.8	Foliation	60
						357.0	108.8	Vein of quartz, K-feldspar, pyrite, galena and sphalerite. 150/52 SW.	
						363.0	110.6	Foliation	50
						363.0	110.6	Vein of quartz, K-feldspar and pyrite. 048/50 SE.	
						379.0	115.5	Foliation	40
						363.0	110.6	Foliation	55
						387.0	118.0	Vein (4 cm) of fluorite and minor	25
						394.0	120.1	Foliation	35

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						394.0	120.1	Vein (2-3 mm) of pure fluorite. 105/54 SW.	
						411.0	125.3	Foliation	45
						417.0	127.1	Foliation	40
						428.0	130.5	Zones of disseminated pyrite and fluorite parallel to foliation.	45
						441.0	134.4	Vein (1 cm) of fluorite, K-feldspar and pyroxene parallel to foliation.	55
						447.0	136.2	Foliation	50
						447.0	136.2	Three veins of quartz with minor pyrite. 090/45 S.	
						447.5	136.4	Vein (4 mm) of fluorite and pyroxene. 155/57 NE.	
						452.0	137.8	Foliation	40
						452.0	137.8	Vein (1 cm) of fluorite and calcite. ~ 000/90.	
						460.0	140.2	Foliation	50
						460.0	140.2	Vein (5 mm) of barren quartz. 080/56 SE.	
						466.0	142.0	Foliation with disseminated fluorite parallel to S1.	40
						478.0	145.7	Foliation	45
						491.0	149.7	Foliation	45
						491.0	149.7	Vein (2 mm) of quartz, pyrite and galena. 105/55 SW.	
						502.0	153.0	Foliation	50
						502.0	153.0	Vein (1-2 mm) of quartz and minor fluorite. 030/50 SE.	
						513.0	156.4	Disseminated fluorite and pyrite in 4- 5 mm zones. Parallel to foliation.	40
						527.0	160.6	Foliation	45
						534.0	162.8	Vein (2-3 mm) of fluorite, pyrite and pyroxene parallel to foliation.	45
						543.0	165.5	Vein (1 cm) of fluorite and calcite parallel to foliation.	25
						548.0	167.0	Clots of pyroxenes parallel to foliation. (5 mm wide).	30
						561.0	171.0	Foliation with vein (2 cm) of fluorite and calcite parallel to S1.	40
						573.0	174.7	Foliation	40
						573.0	174.7	Vein (5 mm) of fluorite, calcite and muscovite. 015/90.	

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						591.0	180.1	Discontinuous and irregular veins of fluorite and calcite parallel to foliation.	40
						607.0	185.0	Vein (3 mm) of pyrite and fluorite parallel to foliation.	40
						623.0	189.9	Vein of fluorite parallel to foliation.	40
						623.0	189.9	Vein-fracture (2 mm) with gypsum. 095/90.	
						636.0	193.9	Foliation	35
						644.0	196.3	Foliation	40
						659.0	200.9	Foliation	45
						659.0	200.9	Vein (2 mm) with quartz. Slickensides: 44-> 215; 080/45 S.	
						665.0	202.7	Foliation	35
675.0	680.0	205.7	207.3		Fine grained quartz-feldspar porphyry with fluorite.	680.0	207.3	Foliation	30
680.0	837.0	207.3	255.1	QLG	Pink to cream aplite	691.0	210.6	Foliation	40
						700.0	213.4	Foliation with fluorite vein (2 mm) parallel to S1	45
						703.0	214.3	Foliation	45
						703.0	214.3	Discontinuous vein (1 cm ) of pure fluorite. 05/35 W.	
						708.0	215.8	Foliation	50
						715.0	217.9	Foliation	40
						725.0	221.0	Foliation	45
						725.0	221.0	Vein (3 mm) of barren quartz. 112/32 NE.	
						734.0	223.7	Foliation	40
						748.0	228.0	Vein (1 cm) of fluorite and K-feldspar parallel to foliation.	45
						767.0	233.8	Foliation	65
						775.0	236.2	Foliation	50
						793.0	241.7	Foliation	90
						803.0	244.8	Foliation	30
						803.0	244.8	Three thin veins (1 mm) or zones of alteration (sheer?). 105/52 NE.	
						812.0	247.5	Foliation	60
						816.0	248.7	Foliation	60
						822.0	250.5	Foliation	30
						822.0	250.5	Vein-fracture (1 mm) with gypsum. 110/58 SW.	
						835.0	254.5	Foliation	55
	837.0		255.1		End of hole				

*From (feet) To (feet) From (metre) To (metre) Unit Lithology Feet Metre Primary tectonic Structure Angle*

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>	
				<b>Client</b>	Eaglet Mines Ltd.					
<b>Project</b>	Structural analysis								<b>Collar Elevation</b>	914.36 m
<b>Drill hole</b>	S28-81								<b>Inclination</b>	90
<b>Coordinates</b>	14 m North								<b>Length</b>	153.6 m
	200.1 m West								<b>Logged by:</b>	P.B Read

**Note: Assumed strike of foliation ~ 090. Unsplit core.**

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
0.0	54.0	0.0	16.5		Overburden				
54.0	72.0	16.5	21.9	Snow-shoe	Medium grained (1-2 mm) biotite-muscovite schist.	67.0	20.4	Foliation	30
						69.0	21.0	Vein (4 mm) of pyrite and K-feldspar. 355/80 NE.	
						69.0	21.0	Quartz vein parallel to foliation.	26
72.0	77.0	21.9	23.5		Medium grained (1 mm) biotite-muscovite (12%) non layered gneiss.				
77.0	78.5	23.5	23.9	Snow-shoe	Medium grained (1-2 mm) biotite-muscovite schist.	78.0	23.8	Quartz sweats parallel to foliation.	35
78.5	83.0	23.9	25.3		Medium grained (1 mm) biotite-muscovite (3%) non layered gneiss.	78.5	23.9	Intrusive contact.	
83.0	88.0	25.3	26.8	Snow-shoe	Medium grained (1 mm) biotite-muscovite schist.	85.0	25.9	Foliation	40
88.0	101.0	26.8	30.8		Medium grained (1 mm) gneissic aplite with disseminated fluorite.	92.0	28.0	Foliation	25
						92.0	28.0	Vein (2 mm) of gypsum. 085/90.	
						92.0	28.0	Vein (2 mm) of quartz and fluorite parallel to foliation. 085/90	25
						93.0	28.3	Foliation	25
						93.0	28.3	Vein (4 mm) of quartz and pyrite and fluorite. 080/90.	
101.0	104.0	30.8	31.7	Snow-shoe	Light green chloritized fine grained (1 mm) biotite-muscovite schist.	101.0	30.8	Foliation	31
104.0	118.0	31.7	36.0		Fine grained (1 mm) cream aplitic gneiss with disseminated fluorite.	106.0	32.3	Vein (4 mm) of gypsum. 083/50 SE.	
						106.0	32.3	Vein ( 5 mm) of quartz 085/50 SE.	
						112.0	34.1	Vein (1.5 cm) of fluorite parallel to foliation.	39
						112.0	34.1	Vein (2 mm) of quartz and pyrite. 055/75 SE.	

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
118.0	119.0	36.0	36.3		Chloritized light green medium grained (1 mm) biotite-muscovite schist.	118.0	36.0	Foliation	27
						118.0	36.0	Contact between aplite gneiss and schist parallel to foliation.	27
						119.0	36.3	Contact between schist and aplite gneiss parallel to foliation.	25
119.0	133.0	36.3	40.5		Aplitic gneiss with disseminated fluorite.	127.0	38.7	to 132' (40.2 m) fault.	
133.0	148.0	40.5	45.1		Medium grained (1 mm) biotite (35%) schist (Snowshoe) with minor pink aplitic sills (30 cm thick).	133.0	40.5	Foliation	28
						133.0	40.5	Faulted contact between overlying aplitic gneiss and underlying Snowshoe schist. 095/53 SW.	
						142.0	43.3	Foliation	23
						142.0	43.3	Vein of K-feldspar, fluorite and calcite. 080/90.	
148.0	151.0	45.1	46.0		Aplitic gneiss with disseminated fluorite.	149.0	45.4	to 151' (46 m) fault.	
151.0	156.0	46.0	47.5		Medium grained (1 mm) biotite schist.	154.0	46.9	Foliation	30
156.0	158.0	47.5	48.2		Cream aplitic gneiss.	156.0	47.5	Contact between overlying biotite schist and underlying aplitic gneiss. Contact irregular and unfaulted.	
158.0	187.0	48.2	57.0		Biotite (20 to 50%) schist	161.0	49.1	Foliation	23

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						163.5	49.8	to 165.0 (50.3 m) fault.	
						167.0	50.9	Foliation	45
						167.0	50.9	Vein (1 cm). 350/60 SW.	
						168.0	51.2	Disseminated fluorite zone (5 cm thick) parallel to foliation.	40
						169.0	51.5	170.0 (51.8 m) fault.	
						173.0	52.7	Foliation	20
						173.0	52.7	Vein (2 mm) of gypsum. 015/41 NW.	
						177.0	54.0	Vein (1 cm) of quartz and pyrite. 010/73 SE.	
						182.0	55.5	Disseminated fluorite zone (3 mm thick) parallel to foliation.	25
187.0	196.0	57.0	59.7		Cream to pink aplite with disseminated fluorite.	187.0	57.0	Unsalted contact between overlying Snowshoe biotite schist and underlying aplite.	
						187.0	57.0	Split core start.	
						195.0	59.4	Disseminated fluorite parallel to foliation.	33
196.0	253.0	59.7	77.1		Medium grained (2-4 mm) pyroxene (2%) leucogranodiorite gneiss.	229.0	69.8	Foliation	30
						241.0	73.5	Foliation	30
						252.0	76.8	Foliation	30
253.0		77.1			Light pinkish grey very fined quartz feldspar porphyry with disseminated fluorite	254.0	77.4	Foliation	50
						295.0	89.9	Foliation	50
						295.0	89.9	Vein (1 mm) of quartz. 063/60 SE.	
						312.0	95.1	Foliation	45
						312.0	95.1	Vein (1 mm) of quartz. 090/60 SE.	
						319.0	97.2	Foliation	52
						329.0	100.3	Foliation	38
						338.0	103.0	Foliation	40
332.0	333.0	101.2	101.5		Very fine grained quartz feldspar porphyry. Rich in purple fluorite.	343.0	104.5	Foliation	33
						343.0	104.5	Vein (2 mm) of quartz and K-feldspar. 065/62 SE.	
						387.0	118.0	Foliation	42
						387.0	118.0	Vein (1 mm) of quartz. 090/81 S.	
						400.0	121.9	Foliation	45
						411.0	125.3	Foliation	45
						423.0	128.9	Foliation	55

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						429.0	130.8	Foliation	50
						436.0	132.9	Foliation	50
						442.0	134.7	Foliation	90
						442.0	134.7	Vein (3 mm) of gypsum. 000/60 E.	
						484.0	147.5	Foliation	42
						484.0	147.5	Vein (3 mm) of quartz and pyrite. 080/50 SE.	
						487.0	148.4	Unsplit core start.	
						490.0	149.4	Foliation	50
						496.0	151.2	Foliation	80
						496.0	151.2	Vein (15 mm) of quartz and pyrite. 280/28 SW.	
						500.0	152.4	Foliation	40
						500.0	152.4	Vein (2 mm) of gypsum. 068/43 NW.	
						503.0	153.3	Foliation	45
	504.0		153.6		End of hole.				



<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
				<b>Client</b>	Eaglet Mines Ltd.				
<b>Project</b>	Structural analysis						<b>Collar Elevation</b>	933.75 m	
<b>Drill hole</b>	S29-81						<b>Inclination</b>	90	
<b>Coordinates</b>	70 m North						<b>Length</b>	221.6 m	
	186 m West						<b>Logged by:</b>	P.B Read	

**Note: Assumed strike of foliation ~ 090. Unsplit core.**

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
0.0	34.0	0.0	10.4		Overburden				
34.0	63.0	10.4	19.2	Snow-shoe	Medium grained (1 mm) medium grey layered biotite-muscovite schist	37.0	11.3	Foliation	45
						53.0	16.2	Foliation	40
						59.0	18.0	Foliation	34
						59.0	18.0	Vein (2 mm) of quartz. 085/77 SE.	
63.0	71.0	19.2	21.6		Biotite-muscovite porphyroblastic garnet schist.	68.0	20.7	Foliation	33
						68.0	20.7	Vein (2 mm) of quartz. 040/vertical.	
71.0	247.0	21.6	75.3		Biotite-muscovite-chlorite schist with rare garnet porphyroblasts.	84.0	25.6	Foliation	27
						84.0	25.6	Third phase fold; AP: 300/42NE; FA: 245/15SE. Southwest vergence.	
						96.0	29.3	Foliation	30
						96.0	29.3	Vein (2 mm) of quartz. 050/vertical.	
						110.0	33.5	Third phase fold; AP: 295/30NE; FA: 305/09NW. Southwest vergence.	
						110.0	33.5	Foliation	0
						127.0	38.7	Foliation	25
						137.0	41.8	Foliation	38
						149.0	45.4	Foliation	29
						158.0	48.2	Foliation	25
						160.0	48.8	Vein (15 cm) of quartz parallel to foliation.	29
						168.0	51.2	Foliation	27
						168.0	51.2	Vein (4 mm) of K-feldspar, quartz and muscovite. 347/90. Vein cuts foliation (and quartz vein).	
						170.0	51.8	Foliation	27

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						170.5	52.0	Vein (1 cm) of dolomite and pyrite. 005/90.	
						180.5	55.0	Foliation	31
						180.5	55.0	Vein (2 mm) of K-feldspar and dolomite. 340/90.	
						194.0	59.1	Foliation	40
						199.0	60.7	Foliation	31
						205-238	62.5 - 72.5	Anastomosing veinlets of vuggy dolomite with minor gypsum in a muscovite-rich schist.	
						210.0	64.0	Sample S28-81-210.0: vuggy dolomite vein in a muscovite schis.	
						214.0	65.2	Foliation	39
						214.0	65.2	Vein (3 cm) of vuggy dolomite and muscovite in a muscovite schist. 285/90.	
						218.0	66.4	Anastomosing veins (up to 1 cm) with dolomite rims and calcite core in muscovite schist.	
						221.0	67.4	Foliation	52
						221.0	67.4	Vein (1 cm) of vuggy dolomite in muscovite schist. 020/90.	
						235.0	71.6	Foliation	30
						241.0	73.5	Foliation	0
						241.0	73.5	Vein (1 cm) with dolomite rim and calcite core.	
						245.0	74.7	Foliation	20
						245.0	74.7	Vein (1-3 mm) of gypsum. 300/58 SW.	
						250.0	76.2	Vein (5 mm) with dolomite rim and calcite core. Parallel to foliation.	30
						250.0	76.2	Vein (1 cm) with dolomite rim and calcite core. 040/90.	
247.0	249.0	75.3	75.9		Fine grained (1 mm) biotite (3%) gneissic aplite sill	266.0	81.1	Foliation	13
249.0	273.5	75.9	83.4	Snow-shoe	Fine grained (1 mm) biotite-muscovite schist with anastomosing calcite-dolomite veins locally vuggy.	269.0-272.0	82 - 83	Vein (3 mm?) of quartz and K-feldspar. K-feldspar alteration in schist on margin of vein.	
						273.0	83.2	Vein (2 cm) or pyrite, quartz and K-feldspar parallel to foliation.	10
273.5	275.0	83.4	83.8		Pink K-feldspar aplitic gneiss sill.	274.0	83.5	Foliation	10

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						274.0	83.5	Vein (1 mm) of quartz and pyrite. 090/70 N.	
275.0	278.0	83.8	84.7		Biotite ± muscovite schist.	276.0	84.1	Vein (2 cm ) of K-feldspar, pyrite, quartz and biotite parallel to foliation.	20
278.0	291.0	84.7	88.7		Pink gneissic aplite sill with sliver of schist.	278.0	84.7	Foliation	10
						278.0	84.7	Vein (2 mm) of quartz, calcite and pyrite. 070/80 NW.	
291.0	293.0	88.7	89.3	Snow-shoe	Fine grained (1 mm) biotite (10-25%) schist	292.0	89.0	Foliation	25
293.0	298.0	89.3	90.8		Pink gneissic aplite sill.	294.0	89.6	Foliation	25
						294.0	89.6	Vein (2 mm) of quartz, pyrite and calcite. 075/80 NW.	
						296.0	90.2	Foliation	15
						296.0	90.2	Vein (5 mm) of quartz and fluorite. 020/90.	
298.0	302.0	90.8	92.1		Fine grained (1 mm) biotite (20%) schist.	296.0	90.2	Vein (1 cm) of quartz. Flat.	
302.0	318.0	92.1	96.9		Cream highly altered schist cut by gypsum veinlets. The rock consists of sericite-quartz-pyrite.	302.0	92.1	Vein and brecciated zone 10 cm thick with dolomite.	
318.0	326.0	96.9	99.4		Medium grained leucogranodiorite altered to muscovite-quartz and sparse, disseminated fluorite.				
326.0	334.0	99.4	101.8		Unidentifiable, highly altered sericite-quartz± fluorite rocks brecciated by fault.				
334.0	336.0	101.8	102.4	Snow-shoe	Biotite schist highly altered to sericite and quartz.	335.0	102.1	Foliation	25
336.0		102.4			Pink K-feldspar aplite with disseminated fluorite and veins of fluorite, pyrite and calcite.	336.0	102.4	Start split core.	
						391.0	119.2	Disseminated fluorite parallel to foliation.	29
						400.0	121.9	Disseminated fluorite parallel to foliation.	40
						413.5	126.0	Disseminated fluorite, biotite, pyroxene and calcite in zones 2 cm thick parallel to foliation.	25
						419.0	127.7	Disseminated fluorite parallel to foliation.	52

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						421.0	128.3	Vein (2 cm) or calcite, pyroxene and fluorite parallel to foliation.	45
						438.0	133.5	Foliation	45
						441.0	134.4	Disseminated fluorite in 8 cm thick zone parallel to foliation.	45
						445.0	135.6	Foliation	70
						445.0	135.6	Vein (5 mm) of quartz and fluorite.	
						447.0	136.2	Foliation	50
						447.0	136.2	Vein (0.5 metre) of quartz, K-feldspar, galena, pyrite and sphalerite.335/20 SW. Has a K-feldspar rich alteration halo.	
						452.0	137.8	Disseminated fluorite parallel to foliation.	55
						467.0	142.3	Foliation	40
						496.0	151.2	Foliation	40
						501.0	152.7	Foliation	20
						505.0	153.9	Foliation	40
						527.0	160.6	Foliation	40
						533.0	162.5	Foliation	43
						547.0	166.7	Foliation	42
						547.0	166.7	Vein (2 mm) of quartz, pyrite and K-feldspar. 292/60 SW.	
						591.0	180.1	Foliation	55
						597.5	182.1	Sparse disseminated fluorite parallel to foliation.	45
						597.5	182.1	Vein (1 mm) of calcite and pyrite. 062/66 SE.	
						632.0	192.6	Foliation	52
						649.0	197.8	Foliation	49
						649.0	197.8	Vein (1 mm) of quartz and fluorite. 055/90.	
						660.0	201.2	Unsplit core.	
						669.0	203.9	Foliation	47
						669.0	203.9	Vein (3 mm) of quartz. 090/42 S.	
						671.0	204.5	Foliation	61
						671.0	204.5	Vein (1 cm) of pyrite, fluorite, graphite, quartz and K-feldspar. 075/63 NW. Parallel late gypsum veins.	
						679.0	207.0	Foliation	63
						679.0	207.0	Vein (2 mm) of gypsum. 090/36 S.	
						679.0	207.0	Vein (3 mm) of quartz. 050/48 SE.	
						681.0	207.6	Foliation	59

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						681.0	207.6	Vein (10 mm) of quartz and dolomite. 058/44 NW.	
						681.0	207.6	Vein (2 mm) of quartz. 290/57 SW.	
						681.0	207.6	Vein (2 mm) of gypsum. 030/52 SE.	
						690.0	210.3	Split core starts.	
						690.0	210.3	Contact between overlying aplitic gneiss and fluorite-bearing aplite.	60
						697.0	212.4	Unsplit core starts.	
						702.0	214.0	Foliation	57
						702.0	214.0	Vein (4 mm) of quartz. 280/58 SW.	
						702.0	214.0	Vein (2 mm) of quartz. 057/75 NW.	
						712.0	217.0	Foliation	40
						712.0	217.0	Vein (2 mm) of quartz. 090/42 S.	
						725.0	221.0	Foliation	60
	727.0		221.6		End of hole.				

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>	
				<b>Client</b>	Eaglet Mines Ltd.					
<b>Project</b>	Structural analysis								<b>Collar Elevation</b>	940.07 m
<b>Drill hole</b>	S30-81								<b>Inclination</b>	90
<b>Coordinates</b>	117 m North								<b>Length</b>	209.3 m
	128 m West								<b>Logged by:</b>	JAF

**Note: Assumed strike of foliation ~ 090. Core is unsplit except from 435' onward.**

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
0.0	24.0	0.0	7.3		Overburden				
24.0	111.0	7.3	33.8	Snow-shoe	Quartz-feldspar-biotite± garnet schist with thin (0.5 mm) layers of hornblende schist with local garnet.	26.0	7.9	Foliation	15
						37.0	11.3	Foliation	30
						45.0	13.7	Foliation	20
						45.0	13.7	Vein (5 mm) of barren quartz. 087/37 N.	
						50.0	15.2	Foliation	20
						62.0	18.9	Foliation	25
						71.0	21.6	Foliation	30
						81.0	24.7	Foliation	25
						81.0	24.7	Vein (2 mm) of barren quartz and minor carbonate. 042/58 SE.	
						84.0	25.6	Foliation	35
						84.0	25.6	Vein of quartz and pyrite. 000/33 W.	
						91.0	27.7	Foliation	20
						91.0	27.7	F2 minor fold- (symmetrical), antiformal overturned to S; AxPI: 090/22N; FAX: 5 -> 85.	
						96.0	29.3	Foliation	30
						96.0	29.3	Vein (4 mm) of chlorite and quartz with slickensides: 85 -> east; vein: 065/90.	
						103.0	31.4	Foliation	35
						103.0	31.4	F2 minor folds (3) - symmetrical, AxPI: 080/15NW; FAX: 10 -> 62.	
						109.0	33.2	Foliation	40
						109.0	33.2	Minor folds (F3?) - west vergent "S fold". AxPI: 335/64NE; FAX: 23 -> 340.	

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
111.0	126.0	33.8	38.4		Hornblende schist with layers (0.5-1.0 m) of biotite schist. Local garnet in both.	114.0	34.7	Foliation	40
						114.0	34.7	Vein (2-3 mm) of barren quartz. 110/52 SW.	
						115.0	35.1	Foliation	30
						115.0	35.1	Minor fault with displacement of 1-2 cm. Some quartz in fracture. 095/62S.	
						121.0	36.9	Foliation	35
						121.0	36.9	Three veins (1 cm) of fibrous quartz. 037/53 NW.	
						121.0	36.9	Vein (1-2 mm) of carbonate. Crosscuts foliation parallel to quartz veins 160/55 NE.	
126.0	135.0	38.4	41.1		Biotite schist with local garnet.	126.0	38.4	Vein (2 mm) of carbonate. 125/82 NE.	
						133.0	40.5	Foliation	30
						133.0	40.5	Vein (3-4 mm) of carbonate. 010/66 W; slickensides: 63 -> E.	
135.0	151.0	41.1	46.0		Biotite schist with layers (up to 0.5 m) of hornblende schist. Both with local garnet.	135.0	41.1	Foliation	20
						135.0	41.1	Vein (1 cm) of carbonate with thin offshoot (2 mm). 062/76 SE.	
						141.0	43.0	Vein (3 cm) of coarse grained calcite parallel to foliation.	30
						143.0	43.6	Foliation	50
						143.0	43.6	Vein (2 mm) of calcite and pyrite. 055/70 NW.	
						149.5	45.6	Vein (5 mm) of quartz and coarse grained pyrite parallel to foliation.	40
						153.0	46.6	Foliation	35
151.0	225.0	46.0	68.6		Biotite schist with local garnet. Abundant quartz.	153.0	46.6	Vein (3-4 mm) of calcite and pyrite cubes. 030/72 NW.	
						157.0	47.9	Foliation	30
						157.0	47.9	Minor fold (F2)- transposed, symmetric. AxPI: 105/40NE; FAX: 15 ->295.	
						161.0	49.1	Foliation	45
						161.0	49.1	Vein (1-2 mm) of calcite and quartz. 170/90.	
						175.0	53.3	Foliation	25

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						175.0	53.3	Vein (5-6 mm) of calcite. Crosscuts quartz veins parallel to S1. 010/75W (calcite vein?)	
						184.0	56.1	Foliation	30
						184.0	56.1	Vein (4 mm) of calcite. 018/60 E.	
						186.0	56.7	Foliation	25
						186.0	56.7	Vein (3 mm) of calcite with slickensides: 30 -> 355; vein: 357/78W.	
						191.0	58.2	Foliation	20
						191.0	58.2	Vein (3 mm) of calcite, quartz and pyrite. 165/83 NE.	
						202.0	61.6	Foliation	20
						202.0	61.6	Vein (5 mm) of calcite. 000/75 W.	
						209.0	63.7	Foliation	20
						209.0	63.7	Vein of carbonate. 120/75 SW.	
						219.0	66.8		50
						219.0	66.8	Vein (5 mm) of calcite with slickensides (30 -> W). Vein: 100/82 S.	
225.0	254.0	68.6	77.4		Biotite-quartz schist with vuggy calcite and pyrite veins and gypsum coated fractures.	225.0	68.6	Foliation	20
						225.0	68.6	Vuggy calcite vein. 165/90.	
						225.0	68.6	Vein-fracture (2 mm) with gypsum. 065/72 NW.	
						226.0	68.9	Foliation	30
						226.0	68.9	Vein of carbonate, clay mineral (montmorillonite?) and chlorite. 065/82 SE.	
						235.0	71.6	Vein (2-4 mm) of quartz and calcite parallel to foliation.	20
						235.0	71.6	Vein (4 mm) of calcite, pyrite and quartz. 028/75 SE.	
						249.0	75.9	Vuggy calcite vein 3 to 4 cm wide, irregular, massive. Foliation	30
						250.0	76.2	Vuggy calcite vein 1 cm wide (no pyrite seen). 025/82 NW.	
254.0	273.0	77.4	83.2		Homblende schist, locally gametiferous with layers of biotite schist (< 0.5 m).	262.0	79.9	Foliation	35
						262.0	79.9	Vein (3 mm) of pure calcite. 014/57 W.	



<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
273.0	332.0	83.2	101.2		Biotite schist with thin layers of hornblende schist (< <0.5 m). Both locally garnetiferous.	274.0	83.5	Minor fold- F2, S vergent- AxPI: 087/25N; FAX: 10 -> 065.	
						274.0	83.5	Foliation	20
						287.0	87.5	Foliation	40
						299.0	91.1	Foliation	35
						303.0	92.4	Foliation	30
						302.0	92.1	Vein -calcite in fracture showing several centimetre offset. 00/50W.	
						303.0	92.4	Vein-fracture (2 mm) of calcite, clay mineral (montmorillonite?) and minor pyrite. 048/73 NW.	
						303.0	92.4	Vein (1 mm) of quartz and calcite; slickensides: 79 -> 310; vein: 032/82 SE.	
						315.0	96.0	Vein (7 cm) of quartz parallel to S1. Foliation	30
						317.0	96.6	Foliation	25
						317.0	96.6	Vein (1 mm) of calcite. 018/63 W.	
						331.0	100.9	Foliation	40
						331.0	100.9	Vein (1.5 cm) of quartz. 135/45 NE.	
332.0	384.0	101.2	117.0		Muscovite-rich schist (+biotite and quartz). Locally very silvery, probably phyllic alteration in part.	337.0	102.7	Foliation	30
						337.0	102.7	Vein (1.5 cm) of quartz and chlorite. 055/70 SE.	
						340.0	103.6	Vein of quartz and pyrite rich zone 3 cm down from quartz vein (20 cm) in schist. Foliation	30
					Muscovite-biotite schist with local thick quartz veins (20-25 cm).	347.0	105.8	Vein (20-25 cm) of quartz with minor calcite, galena and sphalerite at margins. 065/75 NW.	
						347.0	105.8	Vein (3 mm) of calcite. 168/76 NE.	
						370.0	112.8	Foliation	30
						370.0	112.8	Vein- gypsum filled fracture (3 mm). 105/40 SW.	
						379.0	115.5	S vergent minor fold F2- AxPI: 074/37NW; FAX: 10 -> 070.	

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
384.0	387.0	117.0	118.0		Contact-intrusive, irregular, but both rock layers have S1. Pink aplite gneiss with minor muscovite.	384.0	117.0	Foliation	65
389.0	392.0	118.6	119.5	Snow-shoe	Well foliated biotite schist.				
392.0	410.0	119.5	125.0	QLG	Pink aplite and local aplite gneiss with thin (< 15 cm) screens of biotite schist. Disseminated fluorite parallel to S1 common.	397.0	121.0	Irregular vein (1.5-2 cm) of fluorite.	
						409.0	124.7	Foliation	30
						409.0	124.7	Vein (2-3 mm) of quartz and pyrite. 015/62 W.	
410.0	455.0	125.0	138.7	QLG	Pink aplite gneiss with more abundant screens of biotite schist (0.3- 0.5 m thick), tapering again to 10-20 cm near final.	417.0	127.1	Foliation	40
						430.0	131.1	Foliation	40
						435.0	132.6	Vein (1-2 cm) of K-feldspar (QLG?) within biotite schist zone. Parallel to foliation.	45
						435.0	132.6	Core is split from 435' (132.6 m) onward.	
455.0	471.0	138.7	143.6		Pink aplite and aplite gneiss. Muscovite locally abundant.	455.0	138.7	Foliation	45
						455.0	138.7	Vein (2-3 mm) of K-feldspar, pyrite and minor fluorite. 010/70 W.	
						457.0	139.3	Foliation	40
						457.0	139.3	Vein (6 mm) of quartz, fluorite and pyrite. 000/74 W.	
471.0	494.0	143.6	150.6	QLG	Cream coloured aplitic gneiss with local medium grained leucocratic quartz rich zones.	474.0	144.5	Foliation	45
						482.0	146.9	Foliation	25
						482.0	146.9	Vein (3-4 mm) of gypsum. 100/70 SW.	
						489.0	149.0	Fault-breccia in green coloured gouge of clay + gypsum (?) 1 m wide.	
494.0	687.0	150.6	209.4		Fluorite-rich altered aplite with K-feldspar augen (1 cm+) in muscovite-rich matrix.	502.0	153.0	Foliation	40

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						508.0	154.8	Foliation and disseminated fluorite parallel to S1.	35
						515.0	157.0	Foliation	30
						515.0	157.0	Vein (2 mm) of pyrite, K-feldspar, quartz and minor fluorite. 105/78 SW.	
						523.0	159.4	Foliation	30
						523.0	159.4	Vein (1 mm) of gypsum. 070/42 NW.	
						544.0	165.8	Foliation	35
						544.0	165.8	Vein (2 mm) of gypsum and some fluorite. 095/68 S.	
						554.0	168.9	Foliation	60
						565.0	172.2	Vein (1 cm) of fluorite parallel to foliation.	35
						566.0	172.5	Foliation	30
						566.0	172.5	Vein (1 mm) of gypsum. 000/70 W.	
						581.0	177.1	Foliation	40
						586.0	178.6	Foliation	50
						593.0	180.7	Foliation	45
						607.0	185.0	Zone (1 cm) of disseminated fluorite parallel to foliation.	40
						611.0	186.2	Discontinuous vein (5 mm) of fluorite parallel to foliation.	30
						620.0	189.0	Foliation	45
						630.0	192.0	Foliation	35
						644.3	196.4	Foliation	40
						652.0	198.7	Foliation	35
						652.0	198.7	Vein-fracture (2 mm) coated with calcite. 100/65 SW.	
						660.0	201.2	Foliation	30
						662.0	201.8	Vein (5 mm) of fluorite parallel to foliation.	45
						674.0	205.4	Foliation	20
						678.0	206.7	Veins and disseminated fluorite parallel to foliation.	40
						681.0	207.6	Foliation	40
						681.0	207.6	Vein (1 mm) of barren quartz and pyrite. 047/75 SE.	
						684.0	208.5	Foliation	35
	687.0		209.4		End of hole				

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>	
				<b>Client</b>	Eaglet Mines Ltd.					
<b>Project</b>	Structural analysis								<b>Collar Elevation</b>	942.56 m
<b>Drill hole</b>	S31-81								<b>Inclination</b>	90
<b>Coordinates</b>	136.0 m North								<b>Length</b>	236.5 m
	202.0 m West								<b>Logged by:</b>	P.B.Read

**Note: Assumed strike of foliation ~ 090. Unsplit core.**

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
0.0	27.0	0.0	8.2		overburden				
27.0	77.5	8.2	23.6		Fine grained (1 mm) medium grey biotite-muscovite schist with 5 to 10% garnet porphyroblasts 3 to 8 mm in diameter.				
						28.0	8.5	Foliation	10
						42.0	12.8	Foliation	15
						44.0	13.4	Vein (2 mm) of calcite. 015/67 NW.	
						42.0	12.8	Vein (2 mm) of calcite. 280/90.	
						44.0	13.4	Foliation	15
						48.0	14.6	Foliation	15
						50.0	15.2	Foliation	14
						57.0	17.4	Foliation	14
						69.0	21.0	Foliation	10
						69.0	21.0	Vein (3 mm) of calcite. 015/90.	
77.5	98.0	23.6	29.9		Dark green hornblende schist with rare garnet porphyroblasts and minor thin (0.5 m) light grey biotite-quartz-feldspar schist.	83.0	25.3	Foliation	0
98.0	157.0	29.9	47.9		Fine grained (< 1 mm) biotite-muscovite schist with thin (0.5 m) hornblende schist layers.	111.0	33.8	Foliation	24
						111.0	33.8	Vein (3 mm) of calcite.045/90.	
						122.0	37.2	Foliation	14
						128.0	39.0	Fold, 2nd phase. AP: 090/12N; FA: 015/11NE.	
						128.0	39.0	Foliation	12
						137.0	41.8	Foliation	23
						152.0	46.3	Foliation	15
						152.0	46.3	First phase (?) fold. AP: 090/15N; FA: 040/10NE.	

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
157.0	257.0	47.9	78.3		Light to medium grey, fine grained ( 1 mm) biotite-muscovite schist with local garnet porphyroblasts (10mm). Thin muscovite- quartz-biotite layers with garnet (1%) up to 0.4 m thick between 162' (49.4 m) and 173' (52.7 m), 187 to 196' (57.0-59.8 m) and 218 to 226' (66.5-68.9 m).	162.0	49.4	Foliation	16
						162.0	49.4	Fold, 2nd phase. AP: 090/12N; FA: 085/15NE.	
						162.0	49.4	Vein (3 mm) of calcite and pyrite. 325/54 NE.	
						171.0	52.1	Foliation	18
						171.0	52.1	Vein (2 mm) of calcite and pyrite. 317/80 SW.	
						176.0	53.6	Foliation	16
						176.0	53.6	Fold, 3rd phase. AP: 090/38N; FA: 085/05NE. S vergent.	
						178.0	54.3	Vein (3 mm) of pyrite and calcite. 355/90.	
						182.0	55.5	Foliation	27
						182.0	55.5	Fold, 3rd phase. AP: 275/49NE; FA: 275/flat. S vergent.	
						200.0	61.0	Foliation	18
						200.0	61.0	Fold, 2nd or 3rd phase. AP: 280/34NE; FA: 280/flat. Vergence, southwest.	
						210.0	64.0	Foliation	19
						230.0	70.1	Foliation	35
						240.0	73.2	Foliation	15
						240.0	73.2	First phase (?) fold. AP: 090/15N; FA: 003/15NE.	
257.0	275.0	78.3	83.8		Dark green hornblende ± chlorite ± biotite, sparsely garnetiferous schist with thin biotite schist layers.	264.0	80.5	Foliation	15
						264.0	80.5	First phase (?) fold. AP: 090/15N; FA: 090/flat. Vergence south.	
275.0	301.0	83.8	91.7		Medium grained (1 mm) light to medium grey biotite, muscovite and garnet porphyroblasts (5%) schist within (0.4 m) hornblende-plagioclase schist layers.	277.0	84.4	Foliation	38

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						284.0	86.6	Vein with calcite and gypsum.	
						292.0	89.0	Foliation	35
						292.0	89.0	Vein (2 mm) of calcite. 340/76NE.	
						297.0	90.5	Foliation	39
						297.0	90.5	Vein (2 mm) of calcite. 280/48SW..	
301.0	350.0	91.7	106.7		Medium grained (1-2mm) biotite-muscovite schist with local garnet porphyroblasts. Gypsum veinlets common.	301.0	91.7	Vein unknown strike dip vertical. Vuggy calcite, pyrite and magnetite with epidote wall rock alteration.	10
						318.0	96.9	Foliation	23
						325.0	99.1	Vein (2 cm) of quartz and pyrite parallel to foliation.	
						335.0	102.1	Foliation	12
						340.0	103.6	Foliation	5
350.0	355.0	106.7	108.2		Pink aplite.	350.0	106.7	Fault- 30 cm thick gouge and fault breccia.	
355.0	368.0	108.2	112.2		Medium grained biotite-muscovite schist.	358.0	109.1	Foliation	12
368.0	374.0	112.2	114.0		Cream to pink highly altered aplite cut by gypsum veinlets.	374.0	114.0	Foliation	31
374.0		114.0			Medium grained (1-3 mm) leucogranodiorite gneiss with 3% pyroxene.	384.0	117.0	Disseminated fluorite and quartz parallel to foliation.	46
						384.0	117.0	Vein (3 mm) of gypsum. 070/58SE.	
						390.0	118.9	Foliation	35
						390.0	118.9	Vein (4 mm) of quartz. 275/70SW.	
						400.0	121.9	Foliation	45
						400.0	121.9	Vein (4 mm) of quartz and pyrite. 010/52SE.	
						429.0	130.8	Foliation	40
						429.0	130.8	Vein (1 cm) of calcite, quartz and fluorite. 035/90.	
						430.0	131.1	Split core starts.	20
						441.0	134.4	Sparse disseminated fluorite parallel to foliation.	
						447.0	136.2	SampleS31-81-447. Gypsum ?	
						450.0	137.2	First appearance of pyroxene.	38
						455.0	138.7	Foliation	30
						488.0	148.7	Foliation	40
						508.0	154.8	Foliation	45
						512.0	156.1	Disseminated fluorite parallel to foliation.	30

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						519.5	158.3	Disseminated fluorite parallel to foliation.	30
						531.0	161.9	Disseminated fluorite parallel to foliation.	
						532.0	162.2	Foliation	27
						532.0	162.2	Vein (1 mm) of pyrite, quartz and galena. 080/90.	30
						539.0	164.3	Disseminated fluorite parallel to foliation.	30
						560.0	170.7	Disseminated fluorite parallel to foliation.	
						560.0	170.7	Vein (10 mm) of fluorite and calcite. 325/63SW.	30
						563.0	171.6	Foliation	35
						566.0	172.5	Disseminated fluorite and K-feldspar parallel to foliation.	15
						575.0	175.3	Disseminated fluorite, K-feldspar and pyroxene parallel to foliation.	35
						580.0	176.8	Disseminated fluorite parallel to foliation.	
						590.0	179.8	Fault gouge 30 cm.	45
						613.0	186.8	Disseminated fluorite and pyrite parallel to foliation.	
						613.0	186.8	Vein (1 mm) of calcite. 090/40S.	
						619.0	188.7	Foliation	38
						619.0	188.7	Vein (3 mm) of K-feldspar. 090/47S.	40
						624.0	190.2	Foliation	35
						626.0	190.8	Disseminated fluorite parallel to foliation.	
						630.0	192.0	Start of unsplit core.	
						639.0	194.8	Foliation	35
						655.0	199.6	Foliation	30
						655.0	199.6	Vein (1.5 cm) of calcite, pyrite and fluorite. 045/90.	
						665.0	202.7	Foliation	35
						678.0	206.7	Foliation	40
						684.0	208.5	Foliation	45
						684.0	208.5	Vein (1 mm) of muscovite in muscovitized rocks. 090/90.	
						687.0	209.4	Foliation	45
						687.0	209.4	Vein (20 cm) of quartz, K-feldspar, pyrite and galena. 080/67 NW.	
						696.0	212.1	Sparse disseminated fluorite and pyrite parallel to foliation.	

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						699.0	213.1	Foliation	35
						699.0	213.1	Vein (15 mm) of calcite, quartz, fluorite and pyrite. 055/65 NW.	
						705.0	214.9	Foliation	47
						705.8	215.1	Vein (2 mm) of pyrite and calcite. 325/50 SW.	
						709.0	216.1	Foliation	40
						709.0	216.1	Vein (8 mm) of pyrite, calcite and quartz. 313/54 SW.	
						714.0	217.6	Start of split core.	
						727.0	221.6	Foliation	45
					Many irregular gypsum veinlets.	727.0	221.6	Vein (10 mm) of quartz and K-feldspar. 090/50 S.	45
						731.0	222.8	Sparse disseminated fluorite parallel to foliation.	
						745.0	227.1	Foliation	40
						745.0	227.1	Vein (2 mm) of gypsum. Slickensides: 215/. 090/40S.	
						748.5	228.1	Foliation	41
						748.5	228.1	Vein (2 mm) of gypsum. 285/45 SW.	
						758.0	231.0	Foliation	38
						758.0	231.0	Vein (1-3 mm) of quartz and pyrite. 090/58 S.	
						761.5	232.1	Foliation	41
						761.5	232.1	Vein (2 cm) of fluorite, K-feldspar and muscovite. 070/41 SE.	
						761.5	232.1	Vein (2 mm) of gypsum. 090/41 S.	
						761.5	232.1	Foliation	41
						774.0	235.9	Vein (2 mm) of gypsum. 030/66 NW.	
						768.0	234.1	Foliation	35
						774.0	235.9	Foliation	30
	776.0		236.5		End of hole.				



<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
<b>Project</b>	<b>Client</b>			Eaglet Mines Ltd.					947.23 m
<b>Drill hole</b>	Structural analysis							<b>Collar Elevation</b>	90
<b>Coordinates</b>	S32-81							<b>Inclination</b>	230.7 m
	192 m North							<b>Length</b>	JAF
	100m West							<b>Logged by:</b>	

**Note: Assumed strike of foliation ~ 090. Core is unsplit within Snowshoe Group.**

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
0.0	16.0	0.0	4.9		Overburden				
16.0	56.0	4.9	17.1	Snowshoe	Quartz rich biotite schist with local garnet.	20.0	6.1	Foliation	20
						33.0	10.1	Foliation	30
						33.0	10.1	Minor fold- F2, west vergent; AxPI: 120/37NE; FAX: 17 -> 023.	
						34.0	10.4	Vein (1 cm+) of calcite. 015/90.	35
56.0	61.0	17.1	18.6		Fault gouge. Green, local fine breccia	56.0	17.1	Fault gouge and fine breccia zone 5' (1.5 m) thick.	
						79.0	24.1	Foliation	40
						92.5	28.2	Foliation	35
						92.5	28.2	Vein of quartz, calcite, pyrite and vuggy calcite. 073/80 SE.	
						99.0	30.2	Foliation	35
105.0	108.0	32.0	32.9		Thin layer (0.3 m) of light green quartzite with garnet.				
						112.0	34.1	Foliation	30
						112.0	34.1	Vein of calcite, pyrite and quartz. 065/74 NW.	
						128.0	39.0	Foliation	15
						128.0	39.0	Two veins (2-3 mm) of calcite. 075/67 SE.	
						136.0	41.5	Foliation	25
						147.0	44.8	Foliation	30
						147.0	44.8	Minor fold - "crenulation", west vergent, F3 (?). AxPI: 003/57E; FAX: 20 -> 010.	
						150.0	45.7	Foliation	
	162.0		49.4			150.0	45.7	Thick vein (4 cm) of mildly vuggy calcite. 155/65vNE.	
162.0	226.0	49.4	68.9		Biotite schist with local garnet layers. Layers of hornblendes schist locally (10-15 cm).	160.0	48.8	Foliation	35
						162.0	49.4	Foliation	40
						174.0	53.0	Foliation	45

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						176.0	53.6	Foliation	45
						184.0	56.1	Foliation	20
						184.0	56.1	Vein (3 mm) of calcite with slickensides: 25 ->305; vein: 107/47NE.	
						186.0	56.7	Foliation	30
						186.0	56.7	Vein (2-3 mm) of vuggy calcite (locally) and calcite + pyrite. 092/73S, show offset of ~ 5 mm.	
						191.0	58.2	Foliation	35
						191.0	58.2	Vein (4 mm) of calcite. 025/72 SE.	
						201.0	61.3	Foliation	45
						201.0	61.3	Vein (2 mm) of calcite. 043/72 SE.	
						206.0	62.8	Foliation	35
						209.0	63.7	Foliation	30
						209.0	63.7	Vein (5 mm) of vuggy calcite and local gypsum. 010/82 W.	
						212.0	64.6	Foliation	35
						212.0	64.6	Minor fold- southwest vergent, "S fold" F3. AxPI: 135/65NE; FAX: 27 - > 315.	
226.0	235.0	68.9	71.6		Biotite schist with local garnet. Hornblende schist layers increasing frequency.	226.0	68.9	Foliation	25
						231.0	70.4	Foliation	60
						231.0	70.4	Irregular vein (2-4 mm) of calcite. 070/65 NW.	
235.0	273.0	71.6	83.2		Hornblende schist with layers of biotite schist (15-20 cm) and biotite quartzite (local, 15 cm or less).	242.0	73.8	Foliation	25
						255.0	77.7	Foliation	25
						258.0	78.6	Foliation	20
						258.0	78.6	Minor fold- F2-south vergent, "S fold" F3. AxPI: 040/13NW; FAX: 8-> 042.	
273.0	319.0	83.2	97.2		Biotite schist and biotite quartzite; hornblende schist is absent.	264.0	80.5	Foliation	20
						274.0	83.5	Foliation	15
						274.0	83.5	Vein (2-3 mm) of calcite and minor pyrite. 165/50 NE.	
						280.0	85.3	Foliation	10

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						280.0	85.3	Vein (3 mm) of calcite with slickensides: 68-> S; vein: 160/74SW.	
						292.0	89.0	Foliation	20
						305.0	93.0	Foliation	15
						305.0	93.0	Vein (1 mm) of calcite. 010/84 W.	
						317.0	96.6	Foliation	10
319.0	340.0	97.2	103.6		Biotite schist with abundant hornblende, grey quartzite.	320.0	97.5	Foliation	20
						332.0	101.2	Minor fold- F2- east vergent. AxPI: 025/50NW; FAX: 25> 010.	
						332.0	101.2	Foliation	15
						332.0	101.2	Foliation	25
						332.0	101.2	Vein (2 mm) of calcite; offset S1. Slicks: 48° -> W; vein: 100/83N.	
						334.0	101.8	Minor fold- symmetric F2; AxPI: 060/25NW; FAX: 20> 325.	
340.0	407.0	103.6	124.1		Biotite schist, local garnet, with minor hornblende schist (15-20 cm thick).	341.0	103.9	Foliation	25
						348.0	106.1	Foliation	20
						348.0	106.1	Vein (1 cm). 160/75SW.	
						355.0	108.2	Foliation	25
						355.0	108.2	Vein (2 mm) of calcite and chlorite. 165/80 NE.	
						364.0	110.9	Foliation	20
						364.0	110.9	Vein-fracture (1 mm) of calcite with 3 to 4 mm normal offset. 367/72SE.	
						369.0	112.5	Foliation	
						369.0	112.5	Vein-fracture (2 mm) of calcite and quartz with offset. 093/56 NW.	
						384.0	117.0	Foliation	25
						384.0	117.0	Minor fold- F2 (?) southeast vergent. AxPI: 088/43NW; FAX: 14> 085.	
						402.0	122.5	Foliation	40
						402.0	122.5	Minor fold- F2 south vergent. AxPI: 097/52N; FAX: 3> 084.	
						404.0	123.1	Foliation	40
						404.0	123.1	Vein (2-3 mm) of calcite. 170/37W.	

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
407.0	499.0	124.1	152.1		Biotite schist, local garnet, with minor biotite quartzite. Hornblende schist is absent. Muscovite is very abundant in schist (medium grained). Biotite is fine grained.	409.0	124.7	Foliation	15
						415.0	126.5	Foliation	25
						424.0	129.2	Foliation	25
						424.0	129.2	Vein-fracture (2 mm) of gypsum. 032/45 NW.	
						437.0	133.2	Foliation	30
						451.0	137.5	Foliation	25
						451.0	137.5	Vein (1 mm) of calcite. 060/90.	
						461.0	140.5	Foliation	35
						461.0	140.5	Vein (2 mm) of calcite. 152/67 NE.	
						467.0	142.3	Foliation	45
						468.0	142.6	Foliation	35
						468.0	142.6	Vein (3 mm) of calcite. 162/57 SW.	
						491.0	149.7	Foliation	25
						491.0	149.7	Vein (2-3 mm) of calcite. 102/67 SW.	
						494.0	150.6	Vein (< 1mm) calcite coated fracture. 175/90.	
499.0	526.0	152.1	160.3	QLG + Snow-shoe	Intercalations of pink aplite sills (0.3-0.5 m) within sericite-biotite schist. Minor carbonate veins. Contact are irregular (intrusive) but both rock types bear the S1 foliation.	497.0	151.5	Foliation	35
						512.0	156.1	Foliation	35
						512.0	156.1	Gypsum veins (1-2 mm) closely spaced with many veins in succession. 075/52 SE.	
526.0	539.0	160.3	164.3		Pink aplite, locally foliated. Fluorite in veins and disseminated.	526.0	160.3	Foliation	25
						526.0	160.3	Vein (2 mm) of quartz. 140/72SW.	
539.0	570.0	164.3	173.7		Fluorite abundant in some rock. Foliation lacking.	539.0	164.3	Foliation	30
						539.0	164.3	Core is split from 539' (164.3 m) onward.	
						539.0	164.3	Vein (4 mm) of quartz. 012/46 W.	

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						548.0	167.0	Foliation	30
						553.0	168.6	Fault - 7' (2.1 m) of gouge, clay-altered chips	
570.0	757.0	173.7	230.7		Pyroxene alteration present. Irregular layers of pyroxene associated with fluorite and pyrite.	569.0	173.4	Foliation	45
						576.0	175.6	Foliation	35
						578.0	176.2	Vein (15 cm) of gypsum. Unoriented.	
						593.0	180.7	Foliation	45
						600.0	182.9	Disseminated fluorite parallel to S1; also gypsum coating on S1. Foliation	45
						602.5	183.6	Disseminated fluorite parallel to S1. Foliation	50
						602.5	183.6	Vein-fracture (2 mm) with gypsum. 130/60NE.	
						607.0	185.0	Foliation	40
						616.0	187.8	Foliation	40
						616.0	187.8	Vein (1 mm) of barren quartz. 160/70 SW.	
						624.0	190.2	Foliation	15
						624.0	190.2	Vein (2-3 mm) with gypsum. 018/90.	
						632.0	192.6	Foliation	20
						632.0	192.6	Vein (3 mm)- calcite coating in fracture. 097/73 SE.	
						643.0	196.0	Zones of fluorite and minor pyroxene parallel to foliation.	20
						643.0	196.0	Vein of quartz and calcite. 059/82 SE.	
						647.0	197.2	Foliation	15
						647.0	197.2	Vein (2-3 mm) of barren quartz. 031/82 SE.	
						651.0	198.4	Foliation	0
						671.0	204.5	Foliation	20
						676.0	206.0	Foliation	15
						676.0	206.0	Vein. 175/72W.	
						692.0	210.9	Foliation	35
						696.0	212.1	Vein (3-4 mm) of fluorite parallel to foliation.	30
						696.0	212.1	Foliation	35
						698.0	212.8	Vein (2-3 mm) of gypsum. 005/80 E.	

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						707.0	215.5	Foliation	25
						707.0	215.5	Vein (1.5 cm) of quartz and pyrite. 092/62 N.	
						712.0	217.0	Vein of fluorite parallel to foliation	15
						716.0	218.2	Foliation	20
						716.0	218.2	Vein- gypsum coated fracture (1 mm). 082/75 NW.	
						720.0	219.5	Coating of gypsum on S1. Foliation.	35
						722.0	220.1	Vein- gypsum coated fracture. 110/57 NE.	
						732.0	223.1	Disseminated fluorite parallel to S1. Foliation	25
						738.0	224.9	Foliation	15
						738.0	224.9	Vein- gypsum coated fracture. 017/90.	
						741.0	225.9	2.75 m fault breccia. 0.5 to 3 cm fragments of aplite in clay in gouge matrix.	
						757.0	230.7	Foliation	25
757.0			230.7		End of hole.	757.0	230.7	Vein- gypsum filled fracture (2 mm). 105/65 NE.	

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
				<b>Client</b>	Eaglet Mines Ltd.				
<b>Project</b>	Structural analysis							<b>Collar Elevation</b>	947.23 m
<b>Drill hole</b>	S33-81							<b>Inclination</b>	90
<b>Coordinates</b>	57.6 m North							<b>Length</b>	220.3 m
	320.9 m West							<b>Logged by:</b>	JAF

**Note: Assumed strike of foliation ~ 090.**

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
0.0	67.0	0.0	20.4		Overburden.				
67.0	76.5	20.4	23.3		Fine grained (1 mm) medium grey biotite-muscovite schist with 15 to 20% garnet porphyroblasts.	67.0	20.4	Foliation	22
						75.0	22.9	Foliation	10
						75.0	22.9	Fold 1st phase. AxPI: 090/10S; FAX: 240/05SW.	
76.5	112.0	23.3	34.1		Fine grained medium grey biotite-muscovite schist with 5 to 20% garnet porphyroblasts and thin (0.5 m thick) dark green garnetiferous hornblende chlorite schist.	88.0	26.8	Foliation	5
						91.0	27.7	Foliation	0
						91.0	27.7	Fold 3rd phase. AxPI: 330/65NE; FAX: 330/05NW. Southwest vergence- crenulation in 330/05 NW.	
						92.0	28.0	Foliation	20
						92.0	28.0	Fold 3rd phase. AxPI: 330/59NE; FAX: 330/flat. Vergence southwest.	
112.0	128.0	34.1	39.0		Medium grained (2 mm) muscovite-rich schist.				
128.0	130.0	39.0	39.6		Cream aplite sill.				
130.0	133.0	39.6	40.5		Medium grained (1 mm) muscovite-rich biotite schist.				
133.0		40.5			Cream aplite with minor disseminated fluorite and pyrite.	137.0	41.8	Foliation	35
						140.0	42.7	Start of split core.	
						152.0	46.3	First appearance of dark green pyroxene. Numerous gypsum veinlets.	
						166.0	50.6	Foliation	50

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						191.0	58.2	Disseminated K-feldspar, fluorite and pyrite parallel to foliation.	85
						191.0	58.2	Vein (1 mm) of pyrite and quartz. 355/31 SW.	
						193.0	58.8	Disseminated fluorite parallel to foliation.	50
						197.0	60.0	Foliation	
						197.0	60.0	Vein (4 mm) of quartz and gypsum. 090/36 S.	
						209.0	63.7	Disseminated fluorite parallel to foliation.	55
						211.0	64.3	Foliation	40
						217.0	66.1	Foliation	64
						217.0	66.1	Vein (1 mm) of gypsum. 275/45 SW.	
						231.0	70.4	Disseminated fluorite and pyroxene parallel to foliation.	58
						260.0	79.2	Foliation	55
						269.0	82.0	Disseminated fluorite and pyroxene parallel to foliation.	42
						280.0	85.3	Foliation	59
						280.0	85.3	Vein (1 mm) of gypsum and calcite. 035/46 SE.	
						286.0	87.2	Foliation	52
						286.0	87.2	Vein (4 mm) of K-feldspar and fluorite. 090/44 S.	
						291.0	88.7	Foliation	55
						294.0	89.6	Foliation	55
						296.5	90.4	Disseminated fluorite and serpentized pyroxene parallel to foliation.	30
						301.0	91.7	Disseminated fluorite parallel to foliation.	35
						301.0	91.7	Vein (2 mm) of K-feldspar. 275/52 SW.	
						304.5	92.8	Vein (15 mm) of calcite, fluorite and pyroxene parallel to foliation.	40
						316.0	96.3	Vein (2 mm) of K-feldspar, fluorite and pyrite. 090/62 S.	
						316.0	96.3	Foliation	52
						316.0	96.3	Vein (10 mm) of calcite and pyroxene.	0
						324.0	98.8	Disseminated fluorite and K-feldspar parallel to foliation.	44



<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						324.0	98.8	Vein (2 mm) of K-feldspar. 090/50 S.	
						334.0	101.8	Foliation	30
						339.0	103.3	Foliation	35
						342.0	104.2	Foliation	40
						349.0	106.4	Foliation	41
						349.0	106.4	Vein (1 mm) of fluorite and pyrite. 010/55 SE.	
						367.0	111.9	Disseminated fluorite and muscovite parallel to foliation.	32
						370.0	112.8	Unsplit core starts at 370' (112.8 m).	
						371.0	113.1	Disseminated fluorite parallel to foliation.	41
						371.0	113.1	Vein (3 mm) of gypsum. 090/32 S.	
						378.0	115.2	Disseminated fluorite and pyroxene parallel to foliation.	43
						378.0	115.2	Vein (2 mm) of quartz and gypsum. 300/32 SW.	
						385.0	117.3	Foliation	35
						385.0	117.3	Vein (1 mm) of quartz, pyrite and gypsum. 090/45 S.	
						394.0	120.1	Disseminated fluorite, pyroxene and K-feldspar parallel to foliation.	48
						394.0	120.1	Vein (2mm) of fluorite, pyrite and quartz. 090/42 S.	
						400.0	121.9	Disseminated fluorite and pyroxene parallel to foliation.	34
						400.0	121.9	Vein (3 mm) of calcite and gypsum. 090/72 S.	
						405.0	123.4	Disseminated fluorite and K-feldspar parallel to foliation.	57
						405.0	123.4	Vein (1 mm) of gypsum. 090/53S.	
						421.0	128.3	Disseminated fluorite and pyroxene parallel to foliation.	38
						421.0	128.3	Vein (2 mm) of gypsum. 000/38 W.	
						432.0	131.7	Disseminated fluorite, K-feldspar and pyroxene parallel to foliation.	52
						437.0	133.2	Vein (18 cm) of quartz pyrite and fluorite.	
						443.0	135.0	Foliation	45
						455.0	138.7	Foliation	35
						456.5	139.1	Foliation	41

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						456.5	139.1	Vein (2 mm) of gypsum. 065/78 SE.	51
						463.5	141.3	Foliation	21
						463.5	141.3	Vein (6 cm) of quartz, pyrite ± fluorite. 275/72 SW.	
						473.0	144.2	Foliation	45
						473.0	144.2	Vein of pyrite, fluorite and quartz. 045/53 SE.	
						478.5	145.8	Minor disseminated fluorite parallel to foliation.	40
						478.5	145.8	Vein (1 mm) of gypsum. 277/85 NE.	
						478.5	145.8	Vein (2 mm) of pyrite and K-feldspar. 290/45 SW.	
						484.0	147.5	Sparsely disseminated fluorite parallel to foliation.	
						484.0	147.5	Vein set (8), 2 mm thick of pyrite galena and K-feldspar. 028/43SE.	
						496.0	151.2	Foliation	40
						496.0	151.2	Vein (3 mm) of calcite and quartz. 062/49 SE.	43
						511.0	155.8	Foliation	42
						511.0	155.8	Vein (2 mm) of quartz and sphalerite. 028/47 SE.	
						527.0	160.6	Foliation	65
						527.0	160.6	Vein (2 mm) of quartz and sphalerite. 090/55 S.	
						532.0	162.2	Disseminated fluorite and pyroxene parallel to foliation.	
						532.0	162.2	Vein (2 mm) of pyrite and quartz. 310/51 SW.	60
						545.0	166.1	Foliation	48
						550.0	167.6	Unsplit core end at 550' (167.7 m).	35
						560.0	170.7	Foliation	41
						565.0	172.2	Foliation	52
						575.0	175.3	Foliation	49
						575.0	175.3	Vein (1 cm ) of quartz. 300/35 SW.	
						592.0	180.4	Foliation	50
						603.0	183.8	Foliation	62
						608.5	185.5	Foliation	38
						622.0	189.6	Disseminated fluorite and first appearance of pyroxene. Parallel to foliation.	

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						641.0	195.4	Foliation	38
						653.0	199.0	Foliation	42
						661.0	201.5	Foliation	45
						666.0	203.0	Sparsely disseminated fluorite and biotite parallel to foliation.	62
						676.0	206.0	Foliation	40
						685.0	208.8	Foliation	38
						680.0	207.3	Split core end at 680' (207.3 m)	
						685.0	208.8	Vein (2 mm) of gypsum. 290/33SW.	
						694.0	211.5	Coarse pyroxene parallel to foliation.	
						701.0	213.7	Foliation	49
						701.0	213.7	Vein (2 mm) of gypsum. 037/54 SE.	
						717.0	218.5	Foliation	38
	722.8		220.3		End of hole.				905.89 m
									90

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
					Eaglet Mines Ltd.				178.9 m
<b>Project</b>	<b>Client</b>							<b>Collar Elevation</b>	JAF
<b>Drill hole</b>	Structural analysis							<b>Inclination</b>	
	S34-81							<b>Length</b>	
<b>Coordinates</b>	0.6 m North							<b>Logged by:</b>	
	321.9 m West								<i>Angle</i>

**Note: Assumed strike of foliation ~ 090. Core is unsplit for the most part.**

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
0.0	99.0	0.0	30.2		Overburden				45
99.0	120.0	30.2	36.6	Snow-shoe	Biotite schist- (garnet not present in this section) with thin sills of pink aplite (QLG) less than 20 cm thick (contains local fluorite).	102.0	31.1	Foliation	35
						107.0	32.6	Foliation	35
						107.0	32.6	Vein-fracture (1 mm) with carbonate (calcite + dolomite?). 005/85 SE.	
124.0	234.0	37.8	71.3		Pink aplite and muscovite aplite gneiss. Zones of disseminated fluorite common (1 m+)	120.0	36.6	Coating of gypsum (1 mm) on S1 surface. Foliation	
						134.0	40.8	Foliation	45
						134.0	40.8	Vein- irregular gypsum coated fracture. 085/57 S.	
						141.0	43.0	Foliation	50
						141.0	43.0	Vein (2 mm) of gypsum. 075/45 SE.	
						144.5	44.0	Foliation	50
						146.0	44.5	Foliation	55
						152.0	46.3	Foliation	45
						160.0	48.8	Foliation	60
						165.0	50.3	Foliation	60
						174.0	53.0	Foliation	60
						174.0	53.0	Vein (3 mm) of gypsum. 022/56 SE.	40
						177.0	54.0	Foliation	55
						177.0	54.0	Vein with vuggy calcite. 095/84 N.	
						182.0	55.5	Foliation	55
						182.0	55.5	Vein (1-2 mm) of gypsum. 088/70 N.	
						184.0	56.1	Vein and zones of pyroxene and fluorite 1- 1.5 cm wide	

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						187.0	57.0	Foliation	60
						187.0	57.0	Vein (5 mm) of quartz with slickensides; vein: 117/53 SW; slicks: 32°->272.	
						190.0	57.9	Foliation	65
						190.0	57.9	Vein (2 mm) of gypsum. 065/63 SE.	
						199.0	60.7	Foliation	55
						199.0	60.7	Vein (1-2 mm) of quartz with slickensides. Slicks: 35°-> W; vein: 177/69 W.	
						204.0	62.2	Foliation	55
						204.0	62.2	Vein (3-4 mm) of fluorite, pyrite and quartz. 060/62 SE.	
						210.0	64.0	Foliation	65
						210.0	64.0	Vein (1-2 mm) of fluorite, quartz and minor pyrite. 105/45 SW.	
						211.0	64.3	Vein (2 mm) of fluorite, molybdenite, pyrite, sphalerite and quartz. 085/55 S.	
						219.0	66.8	Foliation	60
						219.0	66.8	Vein (2-4 mm) of fluorite, quartz and pyrite. 104/55 SW.	
						230.0	70.1	Foliation	35
						230.0	70.1	Vein (1 cm) of calcite and fluorite. 010/60 E.	
						233.0	71.0	Vein (1.5 cm) of pyroxene, fluorite and K-feldspar parallel to foliation.	60
234.0	250.0	71.3	76.2		Muscovite aplite. Potassic alteration common over 2 metres.	237.0	72.2	Foliation	55
						237.0	72.2	Vein (2 mm) of quartz and ms. 042/46 SE.	
						238.0	72.5	Vein (1 cm) of fluorite, K-feldspar and quartz parallel to S1. Ms altered zone.	60
						242.0	73.8	Foliation	70
						242.0	73.8	Vein (1 cm) of fluorite, calcite, pyroxene and pyrite. 060/68 SE.	
						249.0	75.9	Foliation	50
						249.0	75.9	Vein (2 mm) of quartz and fluorite. 095/68 S.	
250.0	257.0	76.2	78.3		Pyroxene alteration zone reached. Pyroxene and fluorite abundant (10%).	252.0	76.8	Vein of fluorite, pyroxene and calcite parallel to foliation.	

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						253.0	77.1	Foliation	
						253.0	77.1	Vein (2 mm) of quartz and fluorite. 065/60 SE.	
257.0	587.0	78.3	178.9		Carbonate alteration widespread for 3 m. Locally present vuggy calcite.	258.0	78.6	Calcite veins and widespread alteration (vuggy) parallel to foliation	45
						263.0	80.2	Foliation	75
						263.0	80.2	Vein: 049/44 SE; slicks: 28°-<212°.	
						275.0	83.8	Foliation	50
						275.0	83.8	Vein- gypsum filled fracture (2 mm). 040/82 SE.	
						278.0	84.7	Foliation	60
						278.0	84.7	Vein of quartz and minor carbonate. 118/90	
						282.0	86.0	Disseminated fluorite very abundant sub-parallel to S1. Foliation	45
						295.0	89.9	Veins and dissemination of fluorite and minor calcite parallel to foliation.	70
						305.0	93.0	Foliation	50
						312.0	95.1	Foliation	50
						317.0	96.6	Vein (4 mm) of muscovite and fluorite parallel to foliation.	70
						321.0	97.8	Foliation	55
						327.0	99.7	Foliation	50
						337.0	102.7	Foliation	45
						337.0	102.7	Vein of gypsum. 357/90.	
						347.0	105.8	Foliation	40
						351.0	107.0	Foliation	40
						351.0	107.0	Vein (3 mm) of gypsum. 040/68 NW.	
						357.0	108.8	Foliation	70
						357.0	108.8	Vein (4 mm) of gypsum. 127/62 NE.	
						360.0	109.7	Zones of pyroxene and fluorite parallel to foliation.	45
						360.0	109.7	Vein (3-4 mm) of gypsum. 142/53.	
						371.0	113.1	Foliation	65
						371.0	113.1	Vein (4-5 mm) of gypsum. 080/30 S.	
						377.0	114.9	Vein of quartz and fluorite. 130/38 SW.	

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						386.0	117.7	Vein (4 mm) of fluorite, pyrite and quartz. 094/48 S.	
						386.0	117.7	Foliation	50
						392.0	119.5	Closely spaced veins (2-4 mm thick) of quartz and pyrite. 025/44 SE.	
						392.0	119.5	Foliation	55
						395.0	120.4	Foliation	70
						395.0	120.4	Vein of quartz, sphalerite and minor fluorite. 140/38 SW.	
						403.0	122.8	Vein (2 mm) of quartz. Slicks: 34->000; vein: 155/88 SW.	
						403.0	122.8	Foliation	50
						415.0	126.5	Foliation	40
						420.0	128.0	Foliation	50
						420.0	128.0	Vein (2 mm) of gypsum. 160/58 NE.	
						431.0	131.4	Vein (1 mm) of gypsum. 080850N.	
						436.0	132.9	Foliation	50
						436.0	132.9	Vein (2-3 mm) of gypsum. 100/46 SW.	
						440.0	134.1	Vein (2 mm) of gypsum. 055/63 NW.	
						442.0	134.7	From 442' to 500' (134.7-152.4 m) core is completely split	
						453.0	138.1	Foliation	50
						453.0	138.1	Vein- gypsum coated fracture. 000/90.	
						481.0	146.6	Foliation	55
						488.0	148.7	Foliation	60
						488.0	148.7	Vein (2 mm) of gypsum. 005/65 E.	
						497.0	151.5	Foliation	30
						500.0	152.4	From 500' (152.4 m) to end, core is mostly intact.	
						506.0	154.2	Foliation	25
						506.0	154.2	Vein of fluorite, 2 cm and 8 mm thick. 043.52 SE.	
						510.0	155.4	Foliation	25
						510.0	155.4	Vein- gypsum coating in fracture. 016/45 W; Slicks: 14->215.	
						516.0	157.3	Foliation	30
						522.0	159.1	Foliation	35
						522.0	159.1	Vein (1-2 mm) of quartz and pyrite. 015/63 E.	

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						527.0	160.6	Foliation	45
						530.0	161.5	Foliation	50
						530.0	161.5	Vein (2-3 mm) of quartz and minor calcite. 045/68 NW.	
						535.0	163.1	Veins of large crystals of pyroxene and fluorite parallel to foliation. Also gypsum veins.	45
						543.0	165.5	Foliation	30
						543.0	165.5	Vein (3-4 mm) of quartz and minor fluorite. 160/50 SW.	
						552.0	168.3	Foliation	50
						552.0	168.3	Vein (2 mm) of barren quartz. 147/48 SW.	
						552.0	168.3	Vein (3-4 mm) of gypsum. 106/75 NE.	
						557.0	169.8	Foliation	55
						557.0	169.8	Vein (2 mm) of quartz and pyrite. 012/66 E.	
						571.0	174.0	Foliation	50
						577.0	175.9	Foliation	40
						577.0	175.9	Vein (15 cm) of massive quartz (barren). 150/43 NE.	
						585.0	178.3	Vein (5 cm) of vuggy calcite. 062/68 NW.	
						585.0	178.3	Foliation	45
	587		178.9		End of hole.				



<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>	
				<b>Client</b>	Eaglet Mines Ltd.					
<b>Project</b>	Structural analysis								<b>Collar Elevation</b>	874.43 m
<b>Drill hole</b>	S35-81								<b>Inclination</b>	90
<b>Coordinates</b>	82.1 m south 540.0 m West								<b>Length</b>	155.4 m
								<b>Logged by:</b>	JAF	

**Note: Assumed strike of foliation ~ 090. Core is unsplit for the most part.**

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
0.0	29.0	0.0	8.8		Overburden				
29.0		8.8		QLG	Pink aplite and ms aplite gneiss with local granodiorite gneiss (medium grained, pyroxene 5-10%).	37.0	11.3	Foliation	45
						37.0	11.3	Vein (1.5 cm) of fluorite parallel to S1. 060/65 NW.	
						50.0	15.2	Foliation	60
						54.0	16.5	Zones of disseminated fluorite a few cm wide parallel to foliation.	45
						74.0	22.6	Foliation	45
						82.0	25.0	Zones of disseminated fluorite 3-4 cm wide parallel to foliation.	55
						91.0	27.7	Foliation	50
						91.0	27.7	Vein ( 1.5 cm) of fluorite. 130/73 SW.	
						101.0	30.8	Foliation	50
						113.0	34.4	Foliation	55
						113.0	34.4	Veins (1-2 mm) of pyrite and minor fluorite. 040/35 SE.	
						126.0	38.4	Foliation	40
						137.0	41.8	Foliation	35
						137.0	41.8	Two veins (1 cm) of fluorite and quartz. 175/70 SW.	
						152.0	46.3	Foliation	40
						161.0	49.1	Foliation	55
						161.0	49.1	Discontinuous vein (1-2 mm) of pyrite and minor fluorite. 085/54 SW.	
						179.0	54.6	Zones of disseminated fluorite parallel to foliation.	50
						189.0	57.6	Foliation	60
						202.0	61.6	Foliation	50
						207.0	63.1	Foliation	45

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						207.0	63.1	Vein (4-5 mm) of pyrite, quartz and minor fluorite. 030/38 SE.	
						217.0	66.1	Foliation	60
						217.0	66.1	Veins (2) of quartz and pyrite 3-4 mm wide. 060/60 SE.	
						222.0	67.7	Vein (1 cm) of fluorite parallel to foliation.	50
						236.0	71.9	Foliation	45
						243.0	74.1	Foliation	50
						243.0	74.1	Vein (2 mm) of barren quartz. 083/55 N.	
						248.0	75.6	Vein (3 mm) of kaolinite sub-parallel to foliation. 055/50 NW.	
						267.0	81.4	Zones of disseminated fluorite and pyroxene parallel to foliation.	60
						272.0	82.9	Foliation	50
						272.0	82.9	Vein (4-5 mm) of fluorite, galena, sphalerite and quartz. 055/40 SE.	
						280.0	85.3	Veins (2) of fluorite and quartz 3 mm wide parallel to foliation.	50
						288.0	87.8	Foliation	55
						294.0	89.6	Thin disseminated fluorite zones parallel to foliation.	60
						302.0	92.1	Foliation	45
						319.0	97.2	Zones of disseminated fluorite parallel to foliation.	55
						328.0	100.0	Foliation	60
						328.0	100.0	Vein (1-2 mm) of fluorite and quartz. 070/35 SE.	
						345.0	105.2	Foliation	70
						345.0	105.2	Vein (2 mm) of fluorite and quartz. 040/30 SE.	
						351.0	107.0	Vein (3 mm) of K-feldspar, fluorite and quartz. 010/64 E.	
						355.0	108.2	Vein (3 mm) of quartz, pyrite and fluorite. 027/55 SE.	
						357.0	108.8	Vein (2 cm) of quartz with thin (< 1 mm) fluorite at margin. Parallel to foliation.	65
						368.0	112.2	Vein (6-7 mm) of fluorite and pyroxene parallel to foliation.	65
						378.0	115.2	Foliation	60
						378.0	115.2	Vein (2-3 mm) of barren quartz. 065/57 SE.	
						380.0	115.8	Foliation	55

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						380.0	115.8	Two quartz veins (2-3 mm), barren. 035/55 SE.	
						396.0	120.7	Vein and disseminated fluorite and pyroxene (4-5 mm wide) parallel to foliation.	65
						400.0	121.9	Foliation	40
						407.0	124.1	Vein (3-4 cm) of fluorite and minor quartz parallel to foliation.	50
						421.0	128.3	Veins (2-3 cm) of fluorite and quartz parallel to S1. Foliation.	70
						425.0	129.5	Foliation	55
						425.0	129.5	Three veins (1-2 mm) with quartz and minor galena and fluorite. 055/30 SE.	
						442.0	134.7	Foliation	55
						452.0	137.8	Zones of fluorite 1 cm thick parallel to foliation.	45
						452.0	137.8	Vein-fracture with kaolinite. 000/67 E.	
						457.0	139.3	Veins (0.5-1.0 cm) of fluorite and quartz parallel to foliation.	50
						466.0	142.0	Vein (2 cm) of fluorite sub-parallel to foliation.	50
						481.0	146.6	Foliation	40
						490.0	149.4	Foliation	50
						496.0	151.2	Foliation	60
	510		155.4		End of hole.				

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>	
				<b>Client</b>	Eaglet Mines Ltd.					
<b>Project</b>	Structural analysis								<b>Collar Elevation</b>	819.90 m
<b>Drill hole</b>	S36-81								<b>Inclination</b>	90
<b>Coordinates</b>	191.2 m south 555.2 m West								<b>Length</b>	169.7 m
								<b>Logged by:</b>	JAF	

**Note: Assumed strike of foliation ~ 054. Core is intact, unsplit.**

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
0.0	42.0	0.0	12.8		overburden				
42.0	62.0	12.8	18.9		Pink aplite and muscovite-bearing aplite gneiss.	43.0	13.1	Foliation	50
						43.0	13.1	Vein (2 mm) of quartz. 055/25 SE.	
						47.0	14.3	Foliation	50
						47.0	14.3	Vein (1-2 mm) of quartz and fluorite. 055/30SE.	
62.0	88.0	18.9	26.8		Medium grained (1-3 mm) granodiorite gneiss. Pyroxene (2%), quartz (10%), K-feldspar.	64.0	19.5	Foliation	55
						66.0	20.1	Foliation	50
						77.0	23.5	Foliation	55
						87.0	26.5	Foliation	55
						92.0	28.0	Vein (4-5 mm) of pyroxene and fluorite parallel to foliation.	60
						103.0	31.4	Foliation	55
						103.0	31.4	Vein (2 cm) of fluorite. 032/55 NW.	
						112.0	34.1	Foliation	55
						114.0	34.7	Foliation	55
						114.0	34.7	Vein-fracture with kaolinite. 055/85 NW.	
						123.0	37.5	Foliation	65
						123.0	37.5	Fractures filled with pyrite and ± fluorite. 1mm thick. 055/30 NW.	
						135.0	41.1	Foliation	50
						135.0	41.1	Vein (1 cm) of fluorite and quartz. 025/58 SE.	
						142.0	43.3	Foliation	
						142.0	43.3	Vein (6 mm) of quartz. 085/53 S.	
						152.0	46.3	Vein (4 mm) of fluorite and quartz parallel to foliation.	50
						158.0	48.2	Foliation	55

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						158.0	48.2	Vein (5 mm) of fluorite, quartz and pyroxene. 175/54 NE.	
						177.0	54.0	Foliation	45
						177.0	54.0	Vein (3-4 mm) of pyrite, quartz and fluorite. 055/50 SE.	
						179.0	54.6	Foliation	45
						179.0	54.6	Vein (5 cm) of pyroxene, fluorite and pyrite. 085/52 N.	
						184.0	56.1	Foliation	45
						184.0	56.1	Four quartz veins ( 3 mm wide) barren. 090/40 S.	
						192.0	58.5	Foliation	50
						192.0	58.5	Vein (1 cm) of barren quartz. 140/64 SW.	
						202.0	61.6	Foliation	45
						202.0	61.6	Fracture (2 mm) kaolinite coated. 170/75 SE.	45
						209.0	63.7	Foliation	45
						209.0	63.7	Vein (1.5 cm) of fluorite and minor pyroxene. 065/42 NW.	
						219.0	66.8	Foliation	40
						219.0	66.8	Vein (4-5 mm) of pyroxene and fluorite. 170/57 SW.	
						227.0	69.2	Foliation	40
						232.0	70.7	Foliation	40
						232.0	70.7	Vein (8-10 mm) of pure fluorite. 170/86 SW.	
						250.0	76.2	Foliation	55
						250.0	76.2	Vein (2 cm) of fluorite and quartz. 095/17 N.	
						261.0	79.6	Disseminated fluorite parallel to S1. Zones 1 cm wide. Foliation.	50
						268.0	81.7	Foliation	50
						268.0	81.7	Four quartz veins (4 mm wide). 055/35 NW.	
						277.0	84.4	Foliation	55
						277.0	84.4	Vein (3 mm) of fluorite, quartz and sphalerite (?). 050/44 SE.	
						289.0	88.1	Vein (2 cm) of fluorite conformable to foliation	55
						289.0	88.1	Vein (1 cm) of barren quartz. 100/60 SW.	
						298.0	90.8	Foliation	50
						298.0	90.8	Vein ( 2 cm) of fluorite, quartz and pyroxene). 135/65 SW.	

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						302.0	92.1	Foliation	65
						312.0	95.1	Foliation	45
						312.0	95.1	Vein (3 mm) of barren quartz. 055/?	40
						321.0	97.8	Vein (5 mm) of fluorite parallel to S1. Foliation	55
						339.0	103.3	Foliation	60
						339.0	103.3	Vein-fracture (1-2 mm) with kaolinite. 175/80 SW.	
						346.0	105.5	Foliation	45
						351.0	107.0	Foliation	60
						351.0	107.0	Vein of barren quartz. 055/55 SE.	
						362.0	110.3	Foliation	40
						376.0	114.6	Foliation	50
						376.0	114.6	Vein (4 mm) of barren quartz. 055/35 SE.	
						383.0	116.7	Foliation	50
						402.0	122.5	Foliation	50
						402.0	122.5	Irregular vein (5-6 mm) of pure fluorite. 075/55 NW.	
						411.0	125.3	Foliation	60
						426.0	129.8	Zones of disseminated fluorite parallel to foliation. 2-3 cm wide.	55
						434.0	132.3	Disseminated pyrite and calcite parallel to foliation. 2 mm wide.	50
						448.0	136.6	Foliation	35
						448.0	136.6	Vein-fracture (1 mm) with kaolinite. 135/75 NE.	
						461.0	140.5	Foliation	50
						473.0	144.2	Foliation	45
						473.0	144.2	Vein (2-3 mm) of fluorite. Horizontal.	
						482.0	146.9	Zones of disseminated fluorite, pyroxene and calcite parallel to foliation.	60
						492.0	150.0	Foliation	45
						505.0	153.9	Zones of disseminated fluorite parallel to foliation. ~ cm wide.	60
						522.0	159.1	Foliation	50
						529.0	161.2	Foliation	60
						529.0	161.2	Vein (3-4 mm) of barren quartz, 030/40 SE.	
						534.0	162.8	Foliation	60
						534.0	162.8	Discontinuous vein (1-2 cm) of fluorite. 115/45 NE.	

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						549.0	167.3	Foliation	50
						549.0	167.3	Vein (1.5 cm) of fluorite and quartz sub-parallel to S1. 055/58 NW.	
	557		169.8		End of hole.	554.0	168.9	Foliation	45

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
				<b>Client</b> Eaglet Mines Ltd.					
<b>Project</b>	Structural analysis						<b>Collar Elevation</b>	820.02 m	
<b>Drill hole</b>	S37-81						<b>Inclination</b>	90	
<b>Coordinates</b>	129.7 m south						<b>Length</b>	175.8 m	
	770.5 m West						<b>Logged by:</b>	PBR/JAF	

**Note: Assumed strike of foliation ~ 054° (section of 045°). Core is completely split.**

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
0.0	7.0	0.0	2.1		Overburden				
7.0	28.0	2.1	8.5		Leucocratic muscovite aplite gneiss.	21.0	6.4	Foliation	45
28.0	30.0	8.5	9.1		Medium grained pyroxene (?) (10%) granodiorite gneiss (sd).	29.0	8.8	Foliation	45
30.0	32.0	9.1	9.8		Fluorite-bearing aplitic gneiss	29.0	8.8	Sample S37-81-28.5' (8.7 m)	
32.0	35.0	9.8	10.7		Medium grained pyroxene (?) granodiorite gneiss (sd).	33.0	10.1	Foliation	50
35.0	39.0	10.7	11.9		Muscovite-fluorite bearing aplitic gneiss.				
39.0	51.0	11.9	15.5		Medium grained pyroxene (?) granodiorite gneiss (sd).	47.5	14.5	Sample S37-81-47.5 feet (sd)	
51.0	53.0	15.5	16.2		Muscovite- aplitic gneiss.				
53.0	56.0	16.2	17.1		Medium grained pyroxene (?) granodiorite gneiss (sd).				
56.0	68.5	17.1	20.9		Muscovite- aplitic gneiss.	61.0	18.6	Foliation	45
68.5	71.5	20.9	21.8		Medium grained pyroxene (?) granodiorite gneiss (sd).				
71.5	94.5	21.8	28.8		Muscovite- aplitic gneiss with K-feldspar, fluorite and pyrite.				
94.5	103.0	28.8	31.4		Medium grained pyroxene (?) granodiorite gneiss (sd).	97.5	29.7	Foliation	30
103.0	107.0	31.4	32.6		Muscovite- aplitic gneiss.				
107.0	111.0	32.6	33.8		Medium grained pyroxene (?) granodiorite gneiss (sd).	107.5	32.8	Foliation	45
111.0	116.0	33.8	35.4		Aplitic gneiss.				
116.0	119.5	35.4	36.4		Medium grained pyroxene (?), up to 3 mm, 15%, granodiorite gneiss (sd).				
119.5	124.0	36.4	37.8		K-feldspar, pyroxene ± fluorite pegmatite (?)				
124.0	144.0	37.8	43.9		Medium grained pyroxene (5-20%) syenodiorite gneiss (sd).	137.0	41.8	Foliation	55



<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
144.0	147.0	43.9	44.8		K-feldspar- muscovite aplitic gneiss.				
147.0	151.0	44.8	46.0		Fine grained pyroxene (?) 10% syenodiorite gneiss (sd).	148.0	45.1	Foliation	45
151.0	173.5	46.0	52.9		K-feldspar-fluorite-pyroxene (2%) muscovite (<5%) aplite gneiss.				
173.5	175.5	52.9	53.5		Medium grained (1-4 mm) pyroxene (30%) syenodiorite gneiss (sd).				
175.5	179.0	53.5	54.6		K-feldspar-fluorite-pyroxene aplite.				
179.0	185.0	54.6	56.4		Medium grained (1-2 mm) pyroxene (10%) syenodiorite gneiss (sd).	185.0	56.4	Foliation	30
185.0	392.0	56.4	119.5		K-feldspar ± fluorite ± pyroxene aplite, local muscovite zones passing downward into muscovite-bearing aplite gneiss.	294.0	89.6	Foliation	40
						312.0	95.1	Foliation	55
						319.0	97.2	Foliation	55
						344.0	104.9	Foliation	60
						348.0	106.1	Foliation	55
						351.0	107.0	Foliation	55
						351.0	107.0	muscovite abundant in aplite.	
						387.0	118.0	Foliation	45
392.0	393.0	119.5	119.8		Medium grained (2 mm) pyroxene (25%) syenodiorite gneiss (sd).				
393.0	397.0	119.8	121.0		K-feldspar ± fluorite aplite.				
397.0	402.0	121.0	122.5		Fine grained pyroxene (5-40%) syenodiorite gneiss (sd).				
402.0	413.0	122.5	125.9		K-feldspar aplitic gneiss				
413.0	423.0	125.9	128.9		Pyroxene (10%) syenodiorite gneiss with disseminated fluorite parallel to foliation.	414.0	126.2	Foliation	45
423.0	480.0	128.9	146.3		K-feldspar ± fluorite ± pyroxene aplite, becoming muscovite-bearing at 442'.	423.0	128.9	Foliation	45
						447.0	136.2	Foliation	60
480.0	492.0	146.3	150.0		muscovite-bearing aplitic gneiss.	483.0	147.2	Foliation	50
492.0	495.0	150.0	150.9		Medium grained leucocratic pyroxene (?) granodiorite gneiss.	492.0	150.0	Foliation	55
495.0	529.0	150.9	161.2		Pink muscovite-K-feldspar aplitic gneiss.	495.0	150.9	Foliation	55

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						499.0	152.1	Foliation	50
						512.0	156.1	Foliation	50
						512.0	156.1	Vein (1 mm) of fluorite and pyrite. 043/67 SE.	
						515.0	157.0	Foliation	35
529.0	536.0	161.2	163.4		Leucocratic biotite (?) (~ 3%) granodiorite gneiss.	531.0	161.9	Foliation	30
536.0	562.0	163.4	171.3		Pink K-feldspar aplite- grading into pink muscovite aplitic gneiss.	550.0	167.6	Foliation	40
562.0	572.0	171.3	174.3		Medium grained (1-2 mm) pyroxene (?) granodiorite gneiss.	562.0	171.3	Foliation	50
572.0	574.0	174.3	175.0		Pink muscovite-bearing aplitic gneiss.	572.0	174.3	Foliation	40
574.0	577.0	175.0	175.9		Pyroxene (?) (10%) granodiorite gneiss.	574.0	175.0	Foliation	40
	577.0		175.9		End of hole.				

**Note: sd - syenodiorite**

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
<b>Project</b>	<b>Client</b>	Eaglet Mines Ltd.							
<b>Drill hole</b>	S38-81							<b>Collar Elevation</b>	823.82 m
<b>Coordinates</b>	77.3 m south 860.9 m West							<b>Inclination</b>	90
								<b>Length</b>	172.8 m
								<b>Logged by:</b>	PBR/JAF

**Note: Assumed strike of foliation ~ 054° (section of 045°).**  
Core has been completely split and prevents the measuring of vein or orientations.

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
0.0	12.0	0.0	3.7		Overburden				
12.0	26.0	3.7	7.9		Medium grained (1-2 mm) leucocratic biotite (?) ( 5%) granodiorite gneiss.	14.0	4.3	Foliation	55
26.0	40.0	7.9	12.2		Aplitic gneiss with disseminated fluorite and local muscovite zones.	34.0	10.4	Foliation	40
40.0	62.0	12.2	18.9		Cream, medium grained (1-2 mm) leucocratic pyroxene (?) (5- 10%) granodiorite gneiss.	52.0	15.8	Foliation	50
62.0	84.5	18.9	25.8		Pink aplitic gneiss with disseminated fluorite and local muscovite aplitic gneiss.	54.0	16.5	Foliation	40
84.5	88.0	25.8	26.8		Medium grained (1-3 mm) leucocratic pyroxene (?) ( 5%) granodiorite gneiss.				
88.0	95.0	26.8	29.0		Cream coloured aplitic gneiss with minor fluorite.				
95.0	101.0	29.0	30.8		Medium grained (1-3 mm) pink granodiorite gneiss.				
101.0	108.0	30.8	32.9		Pink aplitic gneiss.	105.0	32.0	Foliation	45
108.0	110.0	32.9	33.5		Pink medium grained (2-4 mm) leucocratic pyroxene (?) granodiorite gneiss.	108.0	32.9	Foliation	60
110.0	115.0	33.5	35.1		Pink aplite locally gneissic,				
115.0	122.0	35.1	37.2		Pink leucocratic pyroxene (?) granodiorite gneiss.				
122.0	179.0	37.2	54.6		Pink aplite with disseminated fluorite. Local muscovite gneiss zones.	136.0	41.5	Foliation	45
						172.0	52.4	Foliation	45
179.0	180.0	54.6	54.9		Aplitic gneiss with 1 feet pyroxene rich zone.	176.0	53.6	Foliation	45

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
180.0	213.0	54.9	64.9		Pink aplite with local muscovite aplitic gneiss ( $\pm$ fluorite).	195.0	59.4	Foliation	60
						206.0	62.8	Foliation	50
213.0	214.0	64.9	65.2		1' (30 cm) section of pyroxene granodiorite gneiss.	213.0	64.9	Foliation	55
214.0	227.0	65.2	69.2		Medium grained pyroxene (10%) granodiorite gneiss with disseminated fluorite.	215.0	65.5	Foliation	50
227.0	280.0	69.2	85.3		Pink aplite with fluorite rich zones and local zones of muscovite rich aplitic gneiss.	229.0	69.8	Foliation	45
280.0	287.0	85.3	87.5		Medium grained (1-2 mm) leuco granodiorite gneiss. Pyroxene (2%).				
287.0	337.0	87.5	102.7		Pink aplite with fluorite rich zones and local zones.	334.0	101.8	Foliation	55
337.0	341.0	102.7	103.9		Medium grained (1-3 mm) pyroxene (10%) granodiorite gneiss with disseminated fluorite.	337.0	102.7	Foliation	55
341.0	366.0	103.9	111.6		Pink K-feldspar rich aplite grading downhole into cream aplite.	362.0	110.3	Foliation	60
366.0	376.0	111.6	114.6		Medium grained (1-4 mm) leucocratic pyroxene (2-10%) granodiorite gneiss.	369.0	112.5	Foliation	60
376.0	383.0	114.6	116.7		muscovite-fluorite bearing pink aplite.				
383.0	410.0	116.7	125.0		Medium grained cream to pink leuco granodiorite gneiss (pyroxene 5%). Disseminated fluorite.	385.0	117.3	Foliation	40
						408.0	124.4	Foliation	60
410.0	422.0	125.0	128.6		Pink K-feldspar aplite with disseminated fluorite and local muscovite zones.				
422.0	426.0	128.6	129.8		Pink pyroxene-fluorite syenodiorite gneiss.	426.0	129.8	Foliation	65
426.0	437.0	129.8	133.2		Pink aplite with disseminated fluorite. Fluorite locally foliated.	437.0	133.2	Foliation	40
437.0	441.0	133.2	134.4		Pink muscovite (5-20%)-aplitic gneiss.	441.0	134.4	Foliation	40
441.0	452.0	134.4	137.8		Aplite with disseminated fluorite.				
452.0	455.0	137.8	138.7		muscovite-bearing aplitic gneiss.				

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
455.0	460.0	138.7	140.2		Pink aplite with disseminated fluorite.				
460.0	468.0	140.2	142.6		Pink ms-bearing aplitic gneiss.				
468.0	496.0	142.6	151.2		Pink fluorite bearing pegmatitic aplite.				
496.0	500.0	151.2	152.4		Pyroxene-fluorite granodiorite gneiss.				
500.0	504.0	152.4	153.6		Pink aplite with disseminated fluorite.				
504.0	512.0	153.6	156.1		Pink muscovite (5-25%) aplitic gneiss and pink aplite zones with disseminated fluorite.	506.0	154.2	Foliation	45
512.0	519.0	156.1	158.2		Pink aplite with disseminated fluorite.				
519.0	522.0	158.2	159.1		Pink muscovite-bearing aplitic gneiss.	522.0	159.1	Foliation	50
522.0		159.1			Cream to pink aplitic gneiss with disseminated fluorite-biotite-pyroxene. Locally pyroxene syenodiorite gneiss less than 2' (60 cm) thick.	549.0	167.3	Foliation	60
	567.0		172.8		End of hole.	563.0	171.6	Foliation	50

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
<b>Project</b>	<b>Client</b>			Eaglet Mines Ltd.				<b>Collar Elevation</b>	860.1 m
<b>Drill hole</b>	Structural analysis							<b>Inclination</b>	90
<b>Coordinates</b>	S39-81							<b>Length</b>	163.6 m
	36.5 m south							<b>Logged by:</b>	JAF/PBR
	769.2m West								

**Note: Assumed strike of foliation ~ 054° (section of 045°). Much of the core has been split, leaving about 10% (or less) intact.**

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
0.0	9.0	0.0	2.7		Overburden				
9.0	25.0	2.7	7.6		Cream to pink aplite with disseminated fluorite and pyrite	15.0	4.6	Foliation	50
25.0	37.0	7.6	11.3		Pink, medium grained (1- 4 mm) leucocratic biotite (?) pyroxene granodiorite gneiss.	32.0	9.8	Foliation	55
37.0	70.0	11.3	21.3		Pink aplite with disseminated fluorite and pyrite	46.0	14.0	Foliation	45
						55.0	16.8	Foliation	60
						62.0	18.9	Foliation	65
70.0	72.0	21.3	21.9		Cream aplite.				
72.0	74.0	21.9	22.6		Cream to pink medium grained (1- 4 mm) leucocratic pyroxene (5%) granodiorite gneiss.	74.0	22.6	Foliation	50
74.0	76.0	22.6	23.2		Pink aplite with massive fluorite and pyrite, pegmatite.				
76.0	98.5	23.2	30.0		Pink, medium grained (1- 3 mm) leucocratic pyroxene (3%) granodiorite zone and pink aplite zone between 81 and 82'.	80.0	24.4	Foliation	45
						97.0	29.6	Foliation	50
98.5	121.0	30.0	36.9		Cream aplite with local gneissic zones . Disseminated fluorite and minor pyroxene (?)	113.0	34.4	Foliation	50
121.0	132.0	36.9	40.2		Cream coloured, medium grained (1-3 mm) leucocratic pyroxene (5%) granodiorite gneiss.	123.0	37.5	Foliation	55
132.0	135.0	40.2	41.1		Pink aplite with disseminated fluorite.				
135.0	139.0	41.1	42.4		Muscovite bearing (20%) aplitic gneiss.	136.5	41.6	Foliation	65
139.0	165.0	42.4	50.3		Pink aplite with disseminated fluorite.				

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
165.0	201.0	50.3	61.3		Pink, medium grained (1-3 mm) pyroxene (3- 25%) granodiorite gneiss.	171.0	52.1	Foliation	60
						181.0	55.2	Foliation	60
						189.0	57.6	Foliation	50
						198.0	60.4	Foliation	50
201.0	235.0	61.3	71.6		Pink aplite with disseminated fluorite and pyrite. Local pyroxene granodiorite gneiss zone.	221.5	67.5	Foliation	45
235.0	236.0	71.6	71.9		Medium grey fine grained (1 mm) biotite-rich (40%) pyroxene (?) gneiss with fluorite.	235.5	71.8	Vein (5 mm) of fluorite parallel to foliation.	50
						235.5	71.8	Foliation	50
236.0	239.0	71.9	72.8		Medium grained (1-3 mm) K-feldspar, pyroxene (5%) granodiorite.				
239.0	250.0	72.8	76.2		Pink aplite with disseminated fluorite and zones of muscovite bearing pink aplitic gneiss.	243.0	74.1	Foliation	40
250.0	254.0	76.2	77.4		White medium grained (1-3 mm) pyroxene (3%) granodiorite gneiss.	251.0	76.5	Foliation	35
254.0	270.0	77.4	82.3		Pink aplite and aplite gneiss with disseminated fluorite. Local muscovite-rich aplitic gneiss.				
270.0	278.0	82.3	84.7		Pink muscovite-rich (20%) aplite gneiss.	275.0	83.8	Foliation	45
278.0	341.0	84.7	103.9		Cream muscovite ± biotite aplite gneiss with disseminated fluorite.	282.0	86.0	Foliation	45
						297.0	90.5	Foliation	50
						306.0	93.3	Vein (3 mm) of fluorite. 355/73 SW.	
						311.0	94.8	Foliation	40
						321.0	97.8	Foliation	50
						334.0	101.8	Foliation	45
341.0	350.0	103.9	106.7		Pink aplite with fluorite clots.				
350.0	352.0	106.7	107.3		Cream leucocratic pyroxene (3%) aplite gneiss.	352.0	107.3	Foliation	60
352.0	368.0	107.3	112.2		Pink aplite with disseminated fluorite and minor muscovite.				
368.0	378.0	112.2	115.2		Cream to pink aplite gneiss with fluorite parallel to foliation.	371.0	113.1	Vein of fluorite parallel to foliation.	50

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						371.0	113.1	Foliation	50
378.0	384.0	115.2	117.0		Cream pyroxene (5 to 15%) aplite gneiss.	378.0	115.2	Foliation	55
						383.0	116.7	Foliation	60
384.0	406.0	117.0	123.8		Pink aplite with disseminated fluorite and muscovite (5%).				
406.0	410.0	123.8	125.0		Cream leucocratic pyroxene (1%) aplite gneiss.				
410.0	420.0	125.0	128.0		Pink muscovite-bearing aplite.	411.0	125.3	Foliation	60
420.0	427.0	128.0	130.2		Pyroxene (3 to 10%) aplitic gneiss.	421.0	128.3	Foliation	55
427.0	480.0	130.2	146.3		Muscovite (5%) aplite with local pyroxene aplite layers.	430.0	131.1	Foliation	50
						447.0	136.2	Foliation	40
						467.0	142.3	Foliation	65
						476.0	145.1	Foliation	55
480.0	493.0	146.3	150.3		Cream medium grained (1 to 3 mm) pyroxene (10%) granodiorite gneiss.	486.0	148.1	Foliation	50
493.0	500.0	150.3	152.4		Pink aplite.	500.0	152.4	Foliation	60
500.0	502.0	152.4	153.0		Cream medium grained (1 to 3 mm) leucocratic pyroxene (5%) granodiorite gneiss.				
502.0	508.0	153.0	154.8		Pink aplite.				
508.0	513.0	154.8	156.4		Pyroxene (5 to 35%) gneiss.	509.0	155.1	Vein (5 mm) of fluorite parallel to foliation.	50
						509.0	155.1	Foliation	50
513.0		156.4			Pink aplite with disseminated fluorite and local pyroxene aplite gneiss zones.	517.0	157.6	Foliation	45
	536.7		163.6		End of hole.	527.0	160.6	Foliation	50



<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>	
				<b>Client</b>	Eaglet Mines Ltd.					
<b>Project</b>	Structural analysis								<b>Collar Elevation</b>	873.74 m
<b>Drill hole</b>	S40-81								<b>Inclination</b>	90
<b>Coordinates</b>	30.8 m south 679.4 m West								<b>Length</b>	91.7 m
								<b>Logged by:</b>	PBR/JAF	

**Note: Assumed strike of foliation ~ 054° (section of 045°). Core has been completely split.**

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
0.0	52.0	0.0	15.8		Overburden.				
52.0	102.0	15.8	31.1		Quartz and fluorite vein				
102.0	106.0	31.1	32.3		Pink aplite with disseminated fluorite.				
106.0	132.0	32.3	40.2		Cream to pink medium grained (1-3 mm) leucocratic granodiorite gneiss.				
132.0	196.0	40.2	59.7		Pink aplite with disseminated fluorite and local pyroxene aplitic gneiss layers.				
196.0	238.0	59.7	72.5		Muscovite aplite gneiss.				
238.0	244.0	72.5	74.4		Medium grey, fine grained (1 mm) biotite (35%) + muscovite gneiss and disseminated fluorite.				
244.0	275.0	74.4	83.8		Pink muscovite-bearing (10 to 20%) gneiss and local aplite layers and disseminated fluorite and calcite clots.				
275.0	301.0	83.8	91.7		Cream aplite gneiss.				
	301.0		91.7		End of hole.				

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>	
				<b>Client</b>	Eaglet Mines Ltd.					
<b>Project</b>	Structural analysis								<b>Collar Elevation</b>	876 m
<b>Drill hole</b>	S41-81								<b>Inclination</b>	90
<b>Coordinates</b>	60 m north								<b>Length</b>	160.6 m
	780 m west								<b>Logged by:</b>	PBR

**Note: Assumed strike of foliation ~ 054° (section of 045°). Core has been completely split**

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
0.0	17.0	0.0	5.2		Overburden				
17.0	70.0	5.2	21.3		Fine grained (1 mm) cream to medium grey biotite (5 to 48%) layered aplite and biotite gneiss.				
70.0	91.0	21.3	27.7		Pink aplite with fluorite.	71.0	21.6	Vein (5 mm) of fluorite parallel of foliation.	60
91.0		27.7			Pink and cream medium grained (1-3 mm) leucocratic granodiorite gneiss. End of lithology logging.	174.0	53.0	Vein (3 mm) of galena, pyrite, calcite and biotite. 070/50SE.	
					<b>End of lithology logging.</b>				
						195.0	59.4	Vein (5 mm) of fluorite. 045/40NW.	
						212.0	64.6	Late vein (3 mm) with kaolinite. 027/68SE.	
						234.0	71.3	Contact between overlying fine grained biotite (30%) gneiss and underlying aplite parallel to foliation.	60
						311.5	94.9	Fluorite-muscovite vein. 067/74NW.	
						355.0	108.2	Vein of fluorite, calcite and pyrite. 056/62NW.	
						362.0	110.3	Zone (5 mm thick) of disseminated fluorite parallel to foliation.	50
						387.0	118.0	Vein (2 mm) of fluorite. 012/62SE.	
						397.5	121.2	Vein (3mm) of fluorite and muscovite parallel to foliation.	75
						508.5	155.0	Vein (1 mm) of kaolinite. 042/85NW.	
	527.0		160.6		End of hole.				



<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
					Eaglet Mines Ltd.				
<b>Project</b>	<b>Client</b>							<b>Collar Elevation</b>	888 m
	Structural analysis							<b>Inclination</b>	90
<b>Drill hole</b>	S42-81							<b>Length</b>	175.9 m
<b>Coordinates</b>	40 m south							<b>Logged by:</b>	PBR
	600 m west								

**Note: Core has been completely split.**

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
0.0	17.0	0.0	5.2		Overburden				
						32.0	9.8	Vein (2-4 mm) of fluorite and muscovite parallel to foliation.	50
						47.0	14.3	Vein (1 cm) of fluorite and pyrite parallel to foliation.	50
						93.0	28.3	Vein (6 mm) of biotite, fluorite and pyrite parallel to foliation.	30
						121.0	36.9	Vein (2 mm) of fluorite. 050/68 SE.	
						125.0	38.1	Vein (5 mm) of fluorite, biotite and pyrite parallel to foliation.	50
						145.0	44.2	Foliation parallel to fine grained biotite paragneiss overlying aplite boundary.	55
						150.0	45.7	Vein (5 mm) of fluorite and K-feldspar.	0
						306.0	93.3	Vein (3 mm) of fluorite and pyrite. 088/50SE.	
						425.0	129.5	Vein (1-2 cm) of fluorite and pyroxene. 010/53NW.	
						446.5	136.1	Vein (1 mm) of kaolinite. 045/69NW.	
						448.0	136.6	Vein (1 mm) of kaolinite. 083/73SE.	
						449.0	136.9	Vein (1-2 mm) of fluorite and pyrite parallel foliation.	55
						455.0	138.7	Vein (1 mm) of fluorite and pyrite parallel foliation.	40
						527.0	160.6	Vein (1 cm) of fluorite and pyrite parallel foliation.	
						529.0	161.2	Vein (3 mm) of fluorite and pyrite. 320/78SW. (May be vertical).	
						529.5	161.4	Vein (5 mm) of fluorite and quartz. 020/63NW.	

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						533.0	162.5	Vein (2 mm) of fluorite and pyrite. 340/68NE.	
						546.0	166.4	Vein (3 mm) of fluorite parallel foliation.	60
						551.0	167.9	Vein (1.5 cm) of fluorite parallel foliation.	60
						576.0	175.6	Disseminated fluorite and pyroxene parallel to foliation.	80
	577.0		175.9		End of hole.				

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
<b>Project</b>	<b>Client</b>				Eaglet Mines Ltd.				
<b>Drill hole</b>	Structural analysis							<b>Collar Elevation</b>	907.5 m
<b>Coordinates</b>	S43-81							<b>Inclination</b>	90
	1.0 m north							<b>Length</b>	185.0 m
	293.0 m West							<b>Logged by:</b>	JAF

**Note: Assumed strike of foliation of 90 °. Core is completely split.**

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
0.0	52.0	0.0	15.8		Overburden				
52.0	129.0	15.8	39.3	Snow-shoe	Biotite schist. Muscovite abundant.	106.0	32.3	First appearance of gypsum veins.	
						113.0	34.4	First appearance of K-feldspar in a quartz-K-feldspar-pyrite vein.	
						120.0	36.6	First appearance of fluorite in a K-feldspar vein.	
129.0	137.5	39.3	41.9		Sill of pink aplite (25-35 cm) and gneissic aplite with biotite schist slivers.	134.0	40.8	Vein (2 mm) of calcite. 038/80NW.	
137.5	149.0	41.9	45.4		Medium grained (2 mm) medium grey biotite schist.				
149.0	310.0	45.4	94.5		Pink to cream coloured aplite with fluorite and local up to 3 feet long fine grained (1 mm) muscovite aplitic gneiss zones. At 212' 64.6 m the aplite becomes white to 254' (77.4 m). Sample S43-81-254' (77.4 m): coarse muscovite for K/Ar dating.	150.0	45.7	First appearance of pyroxene.	
						182.0	55.5	Vein (1 cm) of barren quartz. 080/53SE.	
						191.0	58.2	Vein (2-10 mm) of fluorite, calcite and pyrite. 010/85W.	
						200.0	61.0	Vein (1-1.5 cm) of fluorite, calcite and pyrite. 105/53N.	
						214.0	65.2	Vein of fluorite, quartz and pyrite. 150/55	
						225.0	68.6	Vein (2 mm) of greenish gypsum. 035/73NW.	
						227.0	69.2	Vein (1 cm) of fluorite and calcite parallel to foliation.	35
						237.0	72.2	Vein (5 mm) of fluorite. 150/3SW	

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						244.0	74.4	Vein (2 mm) of gypsum. 110/85SW.	
						250.0	76.2	Vein (1-2 mm) of quartz and pyrite. 047/53 NW.	
						267.0	81.4	Vein (5-6 mm) of fluorite and pyrite. 105/44SW.	
						276.0	84.1	Vein of quartz and fluorite. 060/42SE.	
						278.0	84.7	Vein (1 cm) of fluorite, calcite and pyrite. 010/40W.	
						306.0	93.3	Vein (2 mm) of fluorite, calcite and pyrite. 010/88E.	
						306.0	93.3	Vein-fracture (1 mm) with gypsum. 097/75N	
310.0	324.0	94.5	98.8		Pink greenish fine grained grano to syeno diorite; pyroxene (5%), muscovite (2%).	315.0	96.0	Vein-fracture (1-2 mm) with gypsum. 090/83S.	
						319.0	97.2	Vein (1 mm) of pyroxene and pyrite parallel to S1. Foliation	55
324.0	366.0	98.8	111.6		Pink medium grained (1-3 mm) muscovite aplite gneiss with thin (25 cm) pink aplite sills and disseminated fluorite.	325.0	99.1	Vein (1 cm) of fluorite, calcite and pyroxene. 175/72E.	
						343.0	104.5	Vein (5 mm+) of fluorite, calcite and pyrite. 085/53SE.	
						353.0	107.6	Vein of K-feldspar, fluorite and pyroxene. 033/71NW.	
366.0	385.0	111.6	117.3		Pink aplite with fluorite.	368.0	112.2	Fault- 12 feet of gouge and local breccia. Fluorite seen.	
385.0	433.0	117.3	132.0		Pink medium grained (1-3 mm) aplitic muscovite gneiss and thin pink aplite zones 3 feet long.	387.0	118.0	Vein (2 mm) of pyrite, quartz and fluorite. 107/49SE.	
						389.0	118.6	Vein (2-4 mm) of fluorite, quartz and pyrite. 093/42S.	
						412.0	125.6	Vein (3-4 mm) of fluorite and calcite. 000/55W.	
						421.0	128.3	Disseminated fluorite parallel to S1. Foliation	50
						428.0	130.5	Vein (1 cm+) of fluorite and calcite. 012/63 NW.	
433.0	528.0	132.0	160.9		White aplite with disseminated fluorite. Thin white aplitic gneiss zones.	442.0	134.7	Vein (1-2 mm) of calcite and very minor fluorite.	088/90

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						462.0	140.8	Vein (2 cm) of fluorite and K-feldspar. 077/64SE.	
						462.0	140.8	Vein-fracture (2 mm) with gypsum. 002/90	
						482.0	146.9	Vein (2 mm) of gypsum. 108/43NE.	
						502.0	153.0	Vein (3 mm) of fluorite, pyrite and quartz. 025/53NW.	
						506.0	154.2	Vein (3-4 cm) of massive fluorite. 055/58SW.	
						516.0	157.3	Vein (1 cm) of fluorite parallel to foliation.	35
528.0	531.0	160.9	161.9		Granodiorite with pyroxene gneiss margins.	529.0	161.2	Vein of fluorite and disseminated fluorite parallel to foliation.	45
						530.0	161.5	Vein (1-1.5 cm) of pyroxene and fluorite. 031/58NW.	
531.0	532.0	161.9	162.2		White aplite.				
532.0	544.0	162.2	165.8		White to pink aplitic gneiss.	541.0	164.9	Vein (1-1.5 cm) of fluorite parallel to foliation.	60
	540.0		164.6		Medium grained pyroxene (5%) granodiorite gneiss strongly foliated. Local white aplite.	544.0	165.8	Vein (1-1.5 cm) of fluorite and calcite. 085/57NW.	
544.0	557.0	165.8	169.8		Medium grained (2-4 mm) pyroxene (10%) granodiorite gneiss.	551.0	167.9	Vein (1 mm) of quartz and pyrite. 125/47SW.	
557.0	561.0	169.8	171.0		White aplite.				
561.0	609.0	171.0	185.6		Medium grained (2-4 mm) pyroxene (10%) granodiorite gneiss.	563.0	171.6	Vein (2 mm) of quartz and pyrite. 125/57SW.	
						568.0	173.1	Vein-fracture (2-3 mm) with gypsum. 022/48SE.	
						575.0	175.3	Vein-fracture (1 mm) with gypsum. 033/57SE.	
						594.0	181.1	Vein (2-3 mm) of gypsum with slickensides. Vein: 046/78NW; slicks: 60 ->240.	
						604.0	184.1	Vein (3 mm) of barren quartz. 170/32E.	
	609.0		185.6		End of hole.				



<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
<b>Project</b>	<b>Client</b>				Eaglet Mines Ltd.			<b>Collar Elevation</b>	904 m
<b>Drill hole</b>	S44-81							<b>Inclination</b>	90
<b>Coordinates</b>	33 m south 259.0 m West							<b>Length</b>	145.4 m
								<b>Logged by:</b>	JAF

Note: core is at 374' (114 m) below surface at level of adit #2.

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
0.0	40.0	0.0	12.2		Overburden				
40.0	92.0	12.2	28.0	Snow-shoe	Biotite schist with minor pink aplite and muscovite aplite gneiss in thin sills (1 m or less). Muscovite aplite is very thin (< 10 cm) and minor. Gypsum alteration is present from top of hole. Also biotite+calcite+pyrite.	45.0	13.7	Vein-fracture (1 mm) with gypsum. 032/85SE.	
						71.0	21.6	Vein. 020/90.	
						80.0	24.4	Vein (2-3 mm) of quartz and K-feldspar. 036/68SE.	
92.0	137.0	28.0	41.8	QLG	Cream to pink aplite and aplite gneiss with thin zones of leuco granodiorite gneiss 25 cm thick with locally abundant biotite.	100.0	30.5	Vein-fracture with gypsum. 038/69SE.	
						112.0	34.1	Vein of pyroxene, pyrite and minor molybdenite. 015/45NW.	
						120.0	36.6	Vein-fracture (1 mm) with gypsum. 115/84SW.	
137.0	185.0	41.8	56.4		At 137' (41.8 m): White aplite with minor pink aplite containing small veins and seams of fluorite+pyrite.				
					At 144' (43.9 m): cream to pink aplite with zones of muscovite aplite (up to 0.5m), coarse grained K-feldspar+muscovite and occasional veins of fluorite.	157.0	47.9	Vein (4-5 mm) of fluorite and minor quartz and K-feldspar. 080/63NW.	
						164.0	50.0	Vein (3 cm) of fluorite and K-feldspar.	
						167.0	50.9	Vein (2 mm) of calcite with slickensides; slicks: 38->315; vein: 045/45NW.	

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						181.0	55.2	Vein-fracture (1 mm) with coated gypsum. 070/35NW.	
185.0	201.0	56.4	61.3		Pink aplite (muscovite absent) with locally strong pyroxene+fluorite alteration; non foliated.	192.0	58.5	Vein of quartz and pyrite. 002/72E.	
201.0	203.0	61.3	61.9		Mafic sill- biotite+K-feldspar+local muscovite (alteration). Small fluorite veins also.				
215.0	259.0	65.5	78.9		Pink aplite, granodiorite gneiss layers (5-10 cm thick) and minor coarse grained muscovite gneiss (locally 10 cm thick).	205.0	62.5	Vein (2-3 mm) of quartz and minor fluorite. 045/47SE.	
						228.0	69.5	Disseminated fluorite and calcite parallel to S1. Foliation	45
						233.0	71.0	Vein of fluorite, pyroxene and pyrite. 140/60NE.	
						253.0	77.1	Vein (6-7 mm) of calcite and minor quartz. 115/47SW.	
259.0	290.0	78.9	88.4		Cream to pink aplite with thin zones of coarse muscovite-K-feldspar gneiss. No granodiorite gneiss. Muscovite gets finer downward in halo, where biotite becomes locally abundant within fluorite-bearing pink aplite and aplite gneiss.	270.0	82.3	Vein (2-3 mm) of gypsum. 087/65S.	
						282.0	86.0	Vein (1.5 cm) of fluorite. 158/53NE.	
						284.0	86.6	Vein (1 cm+) of fluorite, calcite and pyroxene. 057/46NW.	
290.0	330.0	88.4	100.6	Ore zone	White aplite with local pink aplite and concentrated fluorite (ore zone) passing downward into muscovite aplite as thin (< 10 cm) layers in white aplite.	292.0	89.0	Very abundant disseminated fluorite parallel to S1. Foliation.	45
						296.0	90.2	Vein of gypsum several mm wide. 088/63N.	
						303.0	92.4	Vein-fracture (1 mm) with gypsum. 078/90.	
						311.0	94.8	Fluorite parallel to foliation.	45
						319.0	97.2	Vein (2 mm) of gypsum. 095/56N.	

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
330.0	336.0	100.6	102.4	Ore zone	Granodiorite gneiss with medium grained biotite abundant.				
	336.0	102.4		Ore zone	White aplite with fluorite and large pyroxene crystals at contact with granodiorite below.	337.0	102.7	Vein of quartz, pyroxene and minor pyrite. 018/72W.	
347.0	357.5	105.8	109.0		Medium grained pyroxene (10%) granodiorite gneiss. Fluorite common- disseminated parallel to S1.	357.0	108.8	Pyroxene rich layers parallel to S1 (alteration?). Foliation.	50
357.0	377.0	108.8	114.9	Ore zone	White aplite with fluorite clusters.	374.0	114.0	Foliation is "banded" fluorite ore (~ parallel to S1 in gneiss)	60
377.0	384.0	114.9	117.0		Coarse muscovite-Kfeldspar aplite gneiss and thin (15-20 cm) layer of pink aplite.	379.0	115.5	Vein (2-4 mm) of gypsum. 165/74SW.	
384.0	407.0	117.0	124.1	Ore zone	White aplite + fluorite with several (5-10 cm) minor muscovite-Kfeldspar gneiss layers.	397.0	121.0	Veins and broad zone of fluorite mineralization parallel to foliation.	55
						397.0	121.0	Vein (2 mm) of gypsum with slickensides; vein: 005/83W; slicks: 18 -> S.	
407.0	449.0	124.1	136.9		Medium grained granodiorite with thin layers (10 cm) of cream coloured aplite with fluorite and pyrite mineralization. (possible shoots from main ore).	411.0	125.3	Vein -parallel to S1- gypsum lined fractures (2 mm) . 015/67E.	
						421.0	128.3	disseminated fluorite and pyroxene parallel to foliation.	40
						428.0	130.5	Vein(2 mm) of gypsum. 062/85NW.	
						441.0	134.4	Vein (1 cm) of barren quartz. 065/60NW.	
449.0	477.0	136.9	145.4		Medium grained pyroxene granodiorite gneiss. No aplite present. Strong gypsum alteration locally.	457.0	139.3	Vein-fracture (2 mm) with gypsum. 032/76SE.	
						462.0	140.8	Vein (2-3 mm) containing quartz and molybdenite parallel to foliation.	50
						468.0	142.6	Vein (4 mm) of gypsum with slickensides. Vein: 045/74SE; slicks: 42 ->235.	

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
	477.0		145.4		End of hole.	475.0	144.8	Vein (5 mm) of pyroxene and quartz.	

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
<b>Project</b>	<b>Client</b>				Eaglet Mines Ltd.			<b>Collar Elevation</b>	911.6
<b>Drill hole</b>	S45-81							<b>Inclination</b>	90
<b>Coordinates</b>	901 m south 223 m west				incorrect: 0 to2 m south			<b>Length</b>	175.6 m
								<b>Logged by:</b>	JAF

Note: Assumed strike of foliation ~090. Core has been completely split. Depth in core at level of adit #2 is 121.6 m.

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
0.0	54.0	0.0	16.5		Overburden				
54.0	96.0	16.5	29.3	Snow-shoe	Biotite schist, locally feldspathized, bleached whitish in gradational (intrusive) contact with pink aplite gneiss with locally abundant muscovite gneiss. At 96' (29.3 m) zone of intermixing (sills) ~ 1 m.	69.5	21.2	Vein-fracture (2 mm) with gypsum. 142/67SW.	
						89.0	27.1	Vein (4-5 mm) of quartz and K-feldspar parallel to foliation.	35
96.0	124.0	29.3	37.8	QLG.	Pink aplite gneiss and muscovite gneiss. Disseminated fluorite common. Muscovite gneiss has local zones (10cm) of coarse grained muscovite+Kfeldspar alteration. Gypsum alteration. Pyrite + fluorite mineralization.	104.0	31.7	Vein (< 1 mm) of quartz and minor gypsum. 165/88NE.	
						116.0	35.4	Vein of gypsum with slicks. Vein: 035/90; slicks: vertical.	
124.0	157.0	37.8	47.9		Fine grained muscovite+ biotite aplite gneiss - finely foliated pale pink to cream coloured with very local muscovite and K-feldspar alteration (zones 10 cm or less).	135.0	41.1	Vein-fracture (4 mm) with gypsum. 075/78SE.	
157.0	176.0	47.9	53.6	Snow-shoe	Biotite schist with several zones (10 cm) of pink aplite; upper contact is sharp with aplite gneiss; lower contact is mixed with granodiorite gneiss.	163.0	49.7	Vein (6 mm) of quartz, fluorite and pyrite. 005/63W.	
176.0	185.0	53.6	56.4		Granodiorite gneiss extremely altered, medium grained and leucocratic with pink aplite becoming abundant at 185'.				

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
185.0	206.0	56.4	62.8		Cream coloured aplite, fine grained, no muscovite, in gradational contact with granodiorite gneiss.	189.0	57.6	Foliation + gypsum coated fracture (1 mm). 000/65E.	
						199.0	60.7	Surface coated with gypsum (1 mm) parallel to foliation.	35
206.0	214.0	62.8	65.2		Medium grained granodiorite gneiss. No obvious pyroxene.	209.0	63.7	Vein (3 mm) of quartz, pyrite and minor fluorite. 130/48SW.	
214.0	240.0	65.2	73.2		Pink aplite, non-foliated, medium grained locally with thin zones of medium grained granodiorite gneiss (20-30 cm thick).	215.0	65.5	Vein (2-3 mm) of quartz and pyrite. 140/68SW.	
240.0	264.0	73.2	80.5		Pink to cream aplite with fine grained biotite, locally cutting weak foliation. Disseminated fluorite abundant.	245.0	74.7	Vein (2 mm) of barren quartz. 085/75S.	
264.0	332.0	80.5	101.2		Muscovite aplite gneiss, fine grained. Muscovite in pink K-feldspathized gneiss.	272.0	82.9	Irregular fluorite and Kfeldspar vein (4 mm) parallel to foliation.	50
						287.0	87.5	Vein (3-4 mm) of fluorite and Kfeldspar. 093/64S.	
						298.0	90.8	Vein (5 mm) of fluorite, muscovite and pyrite. 132/43SE.	
						321.0	97.8	Vein (1 cm) of fluorite parallel to foliation.	40
						327.0	99.7	Vein (1 cm) of fluorite, calcite and pyrite. 162/33NE.	
332.0	371.0	101.2	113.1		Coarse muscovite-Kfeldspar aplite, pink, locally foliated with zones (15-20 cm thick) of non-muscovite pink aplite (unfoliated).	332.0	101.2	Vein (3 mm) of barren quartz. 143/55SW.	
						357.0	108.8	Vein (2-3 mm) of gypsum. Vein: 097/77N; slicks: 32 -> E.	
						365.0	111.3	Zones of disseminated fluorite throughout foliation	40
						367.0	111.9	Vein (1 cm) of fluorite and calcite. Horizontal.	
371.0	398.0	113.1	121.3		Pink aplite; very local muscovite-bearing zones in otherwise fine to medium grained Kfeldspar-rich aplite.	377.0	114.9	Vein (1-1.5 cm) of fluorite and calcite. 025/59SE.	

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
398.0	490.0	121.3	149.4	Ore	White aplite with very local pink zones, Calcite+ fluorite + pyrite comprise the alteration phases in this "ore zone". The aplite is very homogeneous and exactly as seen in adit #2.	405.0	123.4	Vein of quartz and minor pyrite. 165/39NE.	
						423.0	128.9	Vein (3 mm) of Kfeldspar, quartz and pyrite. 078/43Se.	
						428.0	130.5	Vein (3-4 mm) of quartz. 130/35NE.	
						476.0	145.1	Vein (3 cm) of fluorite and quartz. 155/80NE.	
						484.0	147.5	Vein (5 mm) of fluorite and quartz. 1161/25NE.	
490.0	512.0	149.4	156.1		White aplite + fluorite ore. (~lower contact of "ore zone") with thin granodiorite gneiss layers (20- 40 cm thick).	491.0	149.7	Vein. 070/60SE.	
512.0	550.0	156.1	167.6			Granodiorite gneiss . Local white aplite; both very altered by gypsum and clay (?). Aplite is thin, 10-20 cm. Granodiorite contains very little pyroxene, except at upper contact with ore, where green pyroxene, calcite +Kfeldspar comprise alteration suite.	514.0	156.7	Vein-fracture (1 mm) with gypsum. 002/82E.
					524.0		159.7	Vein (5 mm- 1.0cm) of quartz, fluorite and pyrite. 150/45SW.	
					525.0		160.0	Vein (4 mm) of fluorite, pyroxene and calcite parallel to foliation.	50
					531.0		161.9	Vein (2 cm) of calcite and fluorite parallel to foliation.	55
					536.0		163.4	Vein (1 cm) of barren quartz. 000/50E.	
550.0	576.0	167.6	175.6		White aplite and Kfeldspar + fluorite + plagioclase (?)- ore interval. This "mixed" zone of white aplite/ fluorite+ medium grained leuco granodiorite gneiss extents to base of hole. Lower contact not encountered.	552.0	168.3	Vein (5-6 mm) of fluorite. 110/48NE.	
						554.0	168.9	Vein (0.5 mm) of pyrite. 012/83E.	
						561.0	171.0	Vein-fracture (2 mm) with gypsum. 080/80S.	

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						570.0	173.7	Vein (3 mm) of barren quartz. 065/47SE..	
	576.0		175.6		End of hole	576.0	175.6	Vein (2 cm) of quartz parallel to S1. Foliation	40



<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
<b>Project</b>	<b>Client</b>			Eaglet Mines Ltd.				<b>Collar Elevation</b>	900.1 m
<b>Drill hole</b>	S46-81							<b>Inclination</b>	90
<b>Coordinates</b>	30 m south 330 m west							<b>Length</b>	132.9 m
								<b>Logged by:</b>	P.B.Read

Note: Assumed strike of foliation ~090. Core split.

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
0.0	101.0	0.0	30.8		Overburden				
101.0	110.0	30.8	33.5		Fine grained (1 mm) biotite (30%) schist with pink aplite sills up to 0.5 m thick.	104.0	31.7	First appearance of disseminated fluorite in a pink aplite sill.	
						106.0	32.3	First gypsum vein	
110.0	120.0	33.5	36.6		Highly altered to gypsum Snowshoe schist.	110.0	33.5	Vein (30 cm) of quartz. Orientation unknown.	
120.0	137.0	36.6	41.8		Fine grained (1 mm) biotite (25%) schist and thin pink aplite sills.	135.0	41.1	Vein (2 mm) 030/69NW.	
137.0	150.0	41.8	45.7		Pink aplite in unfaulted contact with Snowshoe. Contains a few slivers of fine grained (1 mm) muscovite schist, probably altered Snowshoe.				
150.0	170.0	45.7	51.8		Medium grained (2-4 mm) leuco-quartz monzonite highly altered by gypsum.				
170.0	201.0	51.8	61.3		Pink to cream aplite with disseminated fluorite and aplite gneiss.	192.0	58.5	Disseminated fluorite parallel to foliation.	90
						203.0	61.9	Disseminated fluorite and first pyroxene parallel to foliation.	
201.0	205.0	61.3	62.5		Fine to medium grained (1-3 mm) leuco granodiorite and granodiorite gneiss.				
205.0	207.0	62.5	63.1		Pink medium grained (2 mm) muscovite gneiss.				
207.0	220.0	63.1	67.1		White and local pink aplite with disseminated pyrite and fluorite.				
220.0	237.0	67.1	72.2		Pink medium grained (2-3) muscovite gneiss.	234.0	71.3	Disseminated fluorite parallel to foliation.	60
237.0	265.0	72.2	80.8		Pink to white aplite with disseminated fluorite and pyrite.				

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
265.0	288.0	80.8	87.8		Pink medium grained (2-3) muscovite gneiss with thin pink aplite zones	272.5	83.1	Disseminated fluorite parallel to foliation.	65
						282.0	86.0	Vein (2 cm) of pyrite and Kfeldspar. 293/34SW.	
						283.0	86.3	Vein (3 cm) of quartz, pyrite, galena and Kfeldspar. 293/35SW.	
288.0	291.0	87.8	88.7		Medium grained (2-4) leuco granodiorite gneiss.				
291.0	293.0	88.7	89.3		Pink aplite.				
293.0	314.0	89.3	95.7		Pink medium grained (1-2 mm) muscovite-bearing aplite gneiss zones in pink aplite with disseminated pyrite and fluorite.				
314.0	323.0	95.7	98.5		Pink medium grained (2-3 mm) muscovite gneiss.	322.0	98.1	Vein (2 mm) of gypsum. Slicks: 256/30SW; 305/34SW.	
323.0	330.0	98.5	100.6		Pink medium grained (2-3 mm) pegmatitic aplite with disseminated fluorite.				
330.0	341.0	100.6	103.9		Medium grained (2-3 mm) pink muscovite gneiss.	335.0	102.1	Vein (5 mm) of fluorite and muscovite. 295/85NE.	
						336.0	102.4	Vein (1 mm) of gypsum. 080/55NW.	
341.0	347.0	103.9	105.8		Medium grained (2-4 mm) pyroxene (3%) leuco granodiorite gneiss.	345.0	105.2	Vein (3 mm) of quartz. 280/54SW.	
347.0	394.0	105.8	120.1		White to locally pink aplite with disseminated fluorite	367.0	111.9	Fluorite, calcite, pyrite and Kfeldspar parallel to foliation.	70
394.0	403.0	120.1	122.8		Medium grained (1-3 mm) biotite - pyroxene (5%) leuco- granodiorite gneiss.	395.0	120.4	Vein (2 mm) of gypsum. 070/55NW.	
403.0	412.0	122.8	125.6		Pink medium grained (1 mm) muscovite aplite gneiss.				
412.0	425.0	125.6	129.5		Medium grained (1-3 mm) biotite - pyroxene (5%) leuco granodiorite gneiss.	424.0	129.2	Zone (2 cm wide) of fluorite and pyroxene parallel to foliation.	60
425.0	434.0	129.5	132.3		Pink biotite -bearing (3%) aplitic gneiss.				
	436.0		132.9		End of hole.				

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
<b>Project</b>	<b>Client</b>				Eaglet Mines Ltd.				
<b>Drill hole</b>	S47-81							<b>Collar Elevation</b>	895.6 m
<b>Coordinates</b>	65.5 m south 269.5 m west							<b>Inclination</b>	90
								<b>Length</b>	142.0 m
								<b>Logged by:</b>	P.B.Read

Note: Assumed strike of foliation ~090. Core split.

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
0.0	16.0	0.0	4.9		Overburden				
16.0	39.0	4.9	11.9		Pink to cream gneissic aplite. Disseminated fluorite appear at top of hole.	22.0	6.7	Foliation and first appearance of pyroxene	45
39.0	51.0	11.9	15.5		Pink aplite and pegmatitic aplite with fluorite and thin (< 0-3 mm) pink muscovite layers.	39.0	11.9	Foliation defined by fluorite.	59
						44.0	13.4	Vein (1 mm) of Kfeldspar and pyrite. 310/51SW.	
						45.0	13.7	Disseminated fluorite and Kfeldspar parallel to foliation.	50
51.0	80.0	15.5	24.4		Pink aplite gneiss with disseminated fluorite	54.0	16.5	Zone (6 cm) of Kfeldspar, fluorite and pyrite parallel to foliation.	24
						63.0	19.2	Disseminated fluorite parallel to foliation.	45
						64.0	19.5	Disseminated fluorite parallel to foliation.	45
						73.0	22.3	Disseminated fluorite parallel to foliation.	41
						77.0	23.5	Disseminated fluorite parallel to foliation.	32
80.0	115.0	24.4	35.1		Medium grained (2-4 mm) leuco granodiorite gneiss cut by irregular gypsum veinlets and calcite+fluorite veins.	103.0	31.4	Pyroxene parallel to foliation.	49
115.0	119.0	35.1	36.3		Pink medium grained (2 mm) muscovite gneiss.				
119.0	120.5	36.3	36.7		Medium grained (1 mm) dark grey biotite-muscovite (40%) schist sill. Many irregular gypsum veinlets.				
120.0	130.0	36.6	39.6		Pink muscovite-bearing gneissic aplite with disseminated fluorite.				

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
130.0	187.0	39.6	57.0		Pink aplite with disseminated pyrite and fluorite.	186.0	56.7	Disseminated fluorite parallel to foliation.	35
187.0	190.0	57.0	57.9		Medium grained (2-4 mm) muscovite gneiss				
190.0	206.0	57.9	62.8		Pink aplite with disseminated fluorite and pyrite.				
206.0	211.0	62.8	64.3		Medium grained (2-3 mm) pyroxene (3%) leuco-granodiorite gneiss.				
211.0	250.0	64.3	76.2		Pink aplite with disseminated fluorite and pyrite.	212.0	64.6	Disseminated pyroxene and fluorite parallel to foliation.	
						217.0	66.1	Disseminated fluorite parallel to foliation.	35
250.0	253.0	76.2	77.1		Medium grained (2-4 mm) leuco-granodiorite gneiss.				
253.0	267.0	77.1	81.4		Pink fine grained (1 mm) muscovite-bearing gneissic aplite.				
267.0	291.0	81.4	88.7		Pink to white aplite with disseminated fluorite.				
291.0	295.0	88.7	89.9		Medium grained (2-4 mm) leuco-granodiorite gneiss.	294.0	89.6	Vein (1 mm) of Kfeldspar. 359/83SW.	
295.0	297.0	89.9	90.5		White aplite.				
297.0	318.0	90.5	96.9		Medium grained (2 mm) pink muscovite gneiss and thin pink aplitic pegmatites.				
318.0	352.0	96.9	107.3		Medium grained (1-3 mm) pyroxene (5%) leuco-granodiorite gneiss with thin (25 cm) pink or white aplite zones.	323.0	98.5	Vein (3 mm) of quartz and pyrite. 090/50S.	
						327.0	99.7	Vein (3 mm) of quartz and pyrite. 065/53SE.	
						332.0	101.2	Disseminated fluorite parallel to foliation.	58
352.0	371.0	107.3	113.1		White aplite with disseminated fluorite.				
371.0	381.0	113.1	116.1		Medium grained (2-4 mm) pyroxene (5%) leuco-granodiorite gneiss.	377.0	114.9	Vein (1 mm) of quartz and fluorite. 345/43NE.	
381.0	398.0	116.1	121.3		White aplite with disseminated fluorite.	393.0	119.8	Disseminated fluorite parallel to foliation.	55
398.0	421.0	121.3	128.3		Pink aplitic gneiss with pink or white aplite zones up to 2' (60 cm) long.	399.0	121.6	Disseminated fluorite and pyroxene parallel to foliation.	45

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						399.0	121.6	Vein (2 mm) of gypsum.	320/60SW.
						413.0	125.9	Vein (10 mm) of fluorite.	320/35NE.
421.0	422.0	128.3	128.6		Medium grained (1-3 mm) leuco-granodiorite gneiss.				
422.0	449.0	128.6	136.9		White aplite gneiss with fluorite.				
449.0	451.0	136.9	137.5		Medium grained (2-3 mm) leuco-granodiorite gneiss.				
451.0	455.0	137.5	138.7		White pegmatitic aplite.				
455.0	466.0	138.7	142.0		Medium grained (2-3 mm) pyroxene (5%) leuco-granodiorite gneiss.				
	466.0		142.0		End of hole.				

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
<b>Project</b>	<b>Client</b>			Eaglet Mines Ltd.					
<b>Drill hole</b>	Structural analysis							<b>Collar Elevation</b>	896.9 m
<b>Coordinates</b>	S48-81							<b>Inclination</b>	90
	73.6 m south							<b>Length</b>	148.4 m
	241.4 m west							<b>Logged by:</b>	P.B.Read

Note: Assumed strike of foliation ~090. Core split.

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
0.0	12.0	0.0	3.7		Overburden				
12.0	47.0	3.7	14.3		Fine grained (1 mm) biotite (10-20%) aplitic gneiss with scarce Kfeldspar augen- looks like Snowshoe with thin fine grained (1 mm) leucocratic biotite (2%) gneissic aplite sills.	22.0	6.7	Vein (1 mm) of K-feldspar and quartz. 275/66SW.	
						27.0	8.2	First appearance of pyroxene.	
						32.0	9.8	Vein (10 cm) of vuggy quartz, calcite and K-feldspar. Orientation unknown.	
						34.0	10.4	First appearance of fluorite and pyrite. Fluorite, quartz and calcite vein of unknown orientation.	
47.0	61.0	14.3	18.6		Medium grained (2-4 mm) biotite-pyroxene granodiorite gneiss and leuco granodiorite gneiss.				
61.0	74.0	18.6	22.6		White to cream aplite and gneissic aplite.				
74.0	77.0	22.6	23.5		Medium grained (1-3 mm) leuco-granodiorite gneiss.				
77.0	83.0	23.5	25.3		Pink aplite with disseminated fluorite	82.0	25.0	Vein (6 cm) of fluorite and Kfeldspar parallel to foliation.	25
						91.0	27.7	Vein (1 mm) of gypsum. 005/47SE.	
83.0	100.0	25.3	30.5		Fine grained (1 mm) aplitic gneiss.				
100.0	102.0	30.5	31.1		Leucogranodiorite gneiss.				
102.0	111.0	31.1	33.8		Pink aplite with disseminated fluorite				
111.0	125.0	33.8	38.1		Fine grained (1 mm) pink muscovite-bearing aplitic gneiss with fluorite.				

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
125.0	127.0	38.1	38.7		Granodiorite gneiss				
127.0	139.0	38.7	42.4		Pink aplite with disseminated fluorite	137.0	41.8	Zone (2 cm thick) of fluorite and Kfeldspar parallel to foliation.	30
139.0	143.0	42.4	43.6		Pink medium grained (2-3 mm) muscovite gneiss.				
143.0	189.0	43.6	57.6		Cream to pink aplite with disseminated fluorite	150.0	45.7	to 250' core is cut by gypsum veinlets.	
						177.0	54.0	Vein (6 mm) of vuggy calcite. 080/49 SW.	
189.0	192.0	57.6	58.5		Medium grained (2-4 mm) leucogranodiorite gneiss.				
192.0	215.0	58.5	65.5		Pink aplite with disseminated fluorite	210.0	64.0	Vein (1 cm) of fluorite and Kfeldspar parallel to foliation.	56
215.0	219.0	65.5	66.8		Pink medium grained (1-3 mm) muscovite gneiss.				
219.0	227.0	66.8	69.2		Pink aplite with disseminated fluorite				
227.0	237.0	69.2	72.2		Leucogranodiorite gneiss.				
237.0	244.0	72.2	74.4		Pale pink aplite with disseminated fluorite				
244.0	247.0	74.4	75.3		Leucogranodiorite gneiss.				
247.0	255.0	75.3	77.7		Pink aplite.				
255.0	261.0	77.7	79.6		Medium grained (2-4) leucogranodiorite with 1% pyroxene.				
261.0	267.0	79.6	81.4		Pink aplite.				
267.0	289.0	81.4	88.1		Medium grained (2-4) leucogranodiorite with 1% pyroxene.				
289.0	301.0	88.1	91.7		Medium grained (2-4) leucogranodiorite with 1% pyroxene.				
301.0	354.0	91.7	107.9		Pink and white aplite with disseminated fluorite. Mainly white aplite (at 351')				
354.0	379.0	107.9	115.5		Medium grained (2-4 mm) pyroxene (1%) leuco granodiorite.	367.0	111.9	Zone (3 cm thick) of pyroxene and fluorite parallel to foliation.	39
379.0	407.0	115.5	124.1		White aplite with disseminated fluorite				
407.0	487.0	124.1	148.4		Medium grained (2-4 mm) pyroxene (5%) granodiorite gneiss and rare thin aplite zones up to 25 cm.	411.0	125.3	Vein (1 mm) of gypsum. 060/32SE.	
						413.0	125.9	Vein (4 cm) of quartz. 082/69SE.	
						415.0	126.5	Zone (1 cm thick) of fluorite parallel to foliation.	45

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						420.0	128.0	Vein (5 mm) of quartz. 010/54NW.	
						424.0	129.2	Zone (4 cm thick) of pyroxene and calcite parallel to foliation.	25
						428.0	130.5	Zone (2 cm thick) of fluorite, pyroxene, K-feldspar and biotite. parallel to foliation.	54
						445.0	135.6	Vein (4 mm) of quartz, pyrite and galena. 310/50SW.	
						458.5	139.8	Zone (8 cm) of calcite, fluorite and pyroxene parallel to foliation.	35
						461.0	140.5	Vein (2 mm) of quartz, galena and pyrite. 065/48SE.	
						467.0	142.3	Vein (1 mm) of pyrite and quartz. 285/65SW.	
						471.0	143.6	Vein (2 mm) of gypsum. 063/85NW.	
						483.0	147.2	Vein (2 mm) of quartz and pyrite. 315/47SW.	
	487.0		148.4		End of hole.				



<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
<b>Project</b>	<b>Client</b>			Eaglet Mines Ltd.				<b>Collar Elevation</b>	893.9 m
<b>Drill hole</b>	Structural analysis							<b>Inclination</b>	90
<b>Coordinates</b>	S49-81							<b>Length</b>	139.0 m
	54.5 m south							<b>Logged by:</b>	P.B.Read
	303.4 m west								

Note: Assumed strike of foliation ~090. Unsplit core to 106.5' (32.5 m).

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
0.0	82.0	0.0	25.0		Overburden				
82.0	90.0	25.0	27.4		Fine grained (1 mm) biotite (10%) aplitic gneiss.	86.0	26.2	First disseminated fluorite	
90.0	98.5	27.4	30.0		Pink fine grained muscovite-bearing aplite.				
98.5	106.5	30.0	32.5		Fine grained (1 mm) cream coloured aplite.				
106.5	108.0	32.5	32.9		Pale purple fluorite-bearing aplite.				
108.0	117.0	32.9	35.7		Pink medium grained (1-2 mm) muscovite-bearing aplitic gneiss.				
117.0	132.0	35.7	40.2		Pink aplite with disseminated fluorite.				
132.0	142.0	40.2	43.3		Medium grained (1-3) pyroxene (2%) leuco granodiorite gneiss.	139.0	42.4	First appearance of pyroxene.	
142.0	157.0	43.3	47.9		Pink aplite with disseminated fluorite.				
157.0	176.0	47.9	53.6		White aplite with disseminated fluorite.				
176.0	176.3	53.6	53.7		Light grey to purple quartz-feldspar micro porphyry. Sample S49-81-176.0' 53.6 m).				
176.3	179.0	53.7	54.6		White aplite with disseminated fluorite.	178.0	54.3	Vein (2 mm) of quartz. 292/65 SW.	
179.0	313.0	54.6	95.4		Pink medium grained (1-2 mm) muscovite-bearing aplitic gneiss.	186.0	56.7	Vein (3 mm) of quartz and pyrite. 090/52S.	
						191.0	58.2	Vein (1 mm) of Kfeldspar. 090/46S.	
						199.0	60.7	Disseminated fluorite and pyroxene parallel to foliation.	45
						222.0	67.7	Disseminated pyroxene and fluorite parallel to foliation.	23
						244.0	74.4	Disseminated fluorite and pyroxene parallel to foliation.	40

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						249.0	75.9	Disseminated pyroxene and minor fluorite parallel to foliation.	43
						249.0	75.9	Vein (4 mm) of quartz and pyrite. 090/50S.	
						256.0	78.0	Vein (1 mm) of gypsum. 090/77S.	
						261.0	79.6	Disseminated fluorite parallel to foliation.	53
						290.0	88.4	Disseminated pyroxene parallel to foliation.	65
						299.0	91.1	Disseminated fluorite, K-feldspar and carbonate parallel to foliation.	60
						308.0	93.9	Disseminated K-feldspar and fluorite parallel to foliation.	50
313.0	336.0	95.4	102.4		White aplite with disseminated fluorite.				
336.0	350.0	102.4	106.7		Medium grained (2-4 mm) pyroxene (8%) granodiorite gneiss.	341.0	103.9	Vein (2 mm) of pyrite, fluorite and quartz. 090/62S.	
						364.0	110.9	Vein (1 mm) of gypsum. 090/62S. Slicks: 123/50SE.	
						347.0	105.8	Disseminated pyroxene and fluorite parallel to foliation.	50
						380.0	115.8	Disseminated fluorite parallel to foliation.	40
						394.0	120.1	Disseminated fluorite, K-feldspar and calcite parallel to foliation.	50
						401.0	122.2	Disseminated fluorite, pyroxene and calcite parallel to foliation.	60
						405.0	123.4	Disseminated pyrite and fluorite parallel to foliation.	50
						411.0	125.3	Vein (2-3 mm) of quartz and pyrite. 081/39SE.	
						416.0	126.8	Vein (3 mm) of galena, pyrite and quartz. 305/59SW.	
						423.0	128.9	Vein (5 mm) of quartz. 090/59S.	
						428.0	130.5	Vein (2 mm) of quartz and pyrite. 085/26SE.	
						432.0	131.7	Vein (4 cm) of Kfeldspar, quartz, pyrite and fluorite parallel to foliation.	50
						442.0	134.7	Vein (6 mm) of quartz, galena, sphalerite and pyrite. 090/55S.	
						456.5	139.1	Vein (3 mm) of gypsum. 090/72S.	
	456.0		139.0		End of hole.				

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
<b>Project</b>	<b>Client</b>			Eaglet Mines Ltd.				<b>Collar Elevation</b>	918.67 m
<b>Drill hole</b>	Structural analysis							<b>Inclination</b>	90
<b>Coordinates</b>	S58-82							<b>Length</b>	178.61 m
	32.0 m north							<b>Logged by:</b>	PBR
	258.6 m West								

**Note: Assumed strike of foliation of 90 °. Unsplit core.**

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
0.0	17.0	0.0	5.2		Overburden				
17.0	37.0	5.2	11.3		Medium grey fine grained (1 mm) layered biotite-muscovite schist.	25.5	7.8	Vein (5 mm) of vuggy calcite. 308/70 SE.	
37.0	39.0	11.3	11.9		Dark green, biotite-bearing hornblende schist.				
39.0	234.5	11.9	71.5		light to medium grey, medium grained (2 mm) biotite-muscovite schist and very rare garnet porphyroblasts.	85.0	25.9	Irregular veins of dolomite of unknown orientation.	
						91.5	27.9	Vuggy veins of dolomite of unknown orientation. Sample S58-82-91.5' (27.9 m).	
						89.0-	27.1-	Heavily muscovitized schist cut by	
						92.0	28.0	vuggy dolomite vein.	
						124.5	37.9	Vein (3 mm) of calcite. 030/71 SE.	
						181.0	55.2	Vein (2 mm) of calcite and pyrite, Unknown orientation.	
					3' (91 cm) wide sericitized zone from 204' to 207' (62.2- 63.1 m).	204.5	62.3	Vein (2 mm) of gypsum. First appearance of gypsum. Orientation unknown.	
						209.0	63.7	Vein (3 mm) of gypsum. 325/90.	
					Starting at 210' (64 m) are thin (0.5 m) gneissic aplite sills.				
234.5	242.0	71.5	73.8		Cream to pink aplitic gneiss. Sparsely disseminated pyrite and fluorite.	234.5	71.5	Cream aplitic gneiss un faulted contact with Snowshoe schist.	
242.0	244.0	73.8	74.4	Snow-shoe	Fine grained ( 1mm) biotite (20%) schist	235.0	71.6	First appearance of fluorite.	

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
244.0	263.0	74.4	80.2		Fine grained (1 mm) cream to pink muscovite (25%) schist derived from Snowshoe. Sparse disseminated fluorite at 248.0' (75.6 m). Heavy sericitization.				
263.0	304.0	80.2	92.7		Cream aplite with disseminated fluorite and pyrite. Screens of fine grained (1 mm) muscovite gneiss (probably altered Snowshoe).	282.0	86.0	First appearance of pyroxene.	
304.0	345.0	92.7	105.2		Pink medium grained (1-2 mm) muscovite gneiss cut by sills of aplite up to 1 metre long with disseminated fluorite.	295.0	89.9	Disseminated fluorite parallel to foliation	68
						311.0	94.8	Vein (3 mm) of pyrite, fluorite, calcite, K-feldspar and quartz. 075/43 SE.	
						316.0	96.3	Vein (3 mm) of quartz, K-feldspar and pyroxene. 090/33 S.	
						319.0	97.2	Vein (2 mm) of pyrite and K-feldspar. 090/47S.	
						321.0	97.8	Disseminated fluorite and K-feldspar parallel to foliation.	
345.0	365.0	105.2	111.3		Cream to pink pegmatitic aplite with coarse fluorite, calcite and muscovite and slivers of medium grained (2 mm) muscovite gneiss.				
365.0	371.0	111.3	113.1		Medium grained (2 mm) cream coloured muscovite gneiss.				
371.0	382.0	113.1	116.4		Pink to cream slightly gneissic aplite and disseminated fluorite.				
382.0	391.0	116.4	119.2		Cream, medium grained (1-2 mm) muscovite gneiss with K-feldspar and fluorite knots.				
391.0	395.0	119.2	120.4		White aplite.				
395.0	397.0	120.4	121.0		Muscovite gneiss.				
397.0	406.0	121.0	123.8		Fault breccia and gouge				
406.0	430.0	123.8	131.1		Pink pegmatitic aplite with disseminated fluorite and pyrite.	423.0	128.9	Disseminated fluorite parallel to foliation	42
						428.0	130.5	to 448' (136.6 m) Split core.	
						463.0	141.1	Disseminated fluorite parallel to foliation	48
						448.0	136.6	to 539' (164.3 m) unsplit core.	

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						463.0	141.1	Vein (2 mm) of pyrite and quartz. 290/39 SW	
					Starts at 515' (157 m) many gypsum veins to end of hole.	535.0	163.1	to 543' (165.5 m) split core.	
						543.0	165.5	to 553' (168.6 m) Unsplit core.	
						544.0	165.8	Vein (1 mm) of gypsum. 300/48 SW.	
						553.0	168.6	to 586' (178.6 m) Split core.	
						556.0	169.5	Disseminated fluorite and K-feldspar parallel to foliation.	52
						561.0	171.0	First appearance of certain pyroxene.	
	586.0		178.6		Note that this hole only has pyroxene alteration and no pyroxene granodiorite gneiss present. End of hole.				

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
<b>Project</b>	<b>Client</b>			Eaglet Mines Ltd.				<b>Collar Elevation</b>	881.66 m
<b>Drill hole</b>	Structural analysis							<b>Inclination</b>	90
<b>Coordinates</b>	S59-82							<b>Length</b>	117.6 m
	103.8 m south							<b>Logged by:</b>	PBR
	284.2 m west								

**Note: Assumed strike of foliation of 90 °. Unsplit core.**

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
			0.0						
0.0	25.0	0.0	7.6		Overburden				
25.0	57.0	7.6	17.4		Cream to pink aplite with discontinuous fluorite + pyrite and slivers of fine grained altered (muscovite) aplite gneiss all cut by veins of quartz, fluorite and pyrite.	25.0	7.6	First appearance of fluorite at top of hole.	
						29.5	9.0	First gypsum vein	
						44.0	13.4	Vein (2 cm ) of fluorite.	
						54.0	16.5	Vein (10 cm) of fluorite, quartz and pyrite.	
57.0	59.5	17.4	18.1		Fine grained (1 mm) biotite (15%) aplitic gneiss.	57.0	17.4	Contact between aplite and fine grained biotite (15%) aplitic gneiss.	
59.5	62.0	18.1	18.9		Medium grained (2-4 mm) leuco-granodiorite.				
62.0	64.0	18.9	19.5		highly altered (gypsum) muscovite gneiss.				
64.0	79.0	19.5	24.1		Cream to pink aplite with discontinuous fluorite + pyrite.	64.0	19.5	to 74' (22.6 m) split core.	
						66.0	20.1	White aplite dike 2 cm thick.	
						74.0	22.6	Disseminated fluorite parallel to foliation.	32
						74.0	22.6	to 80' (24.4 m) unsplit core.	
79.0	90.0	24.1	27.4		Cream, fine grained (1 mm) biotite (2-20%) aplitic gneiss.	80.0	24.4	to 86' (26.2 m) split core	
90.0	94.0	27.4	28.7		Cream gneissic aplite.	90.5	27.6	Vein (1 mm) of gypsum. 058/70 SE.	
94.0	101.0	28.7	30.8		Medium grained (2-4 mm) leuco-granodiorite.				
101.0	116.0	30.8	35.4		Pink aplite with muscovite gneiss	106.0	32.3	First appearance of pyroxene.	
116.0	134.0	35.4	40.8		Pyroxene (20%) granodiorite gneiss and aplite sills with disseminated fluorite.	116.0	35.4	Contact between pink aplite and pyroxene leucograndiorite gneiss.	

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
134.0	153.0	40.8	46.6		Pink aplite and local fine grained (1 mm) muscovite gneiss layers.				
153.0	160.0	46.6	48.8		Pink medium grained (2-4 mm) pyroxene (2%) leuco-granodiorite gneiss.				
160.0	163.0	48.8	49.7		Pink aplite and local fine grained (1 mm) muscovite gneiss layers and disseminated fluorite.				
163.0	168.0	49.7	51.2		Quartz vein	163.0	49.7	to 289' (88.1 m) split core	
						163.0	49.7	to 168' (51.2 m). Vein of quartz, galena, pyrite, tetrahedrite (?), fluorite and sphalerite.	
					Sample S59-82-167' (50.9 m) for opaques				
182.0	212.0	55.5	64.6		Quartz vein	182.0	55.5	to 212' (64.6 m). Vein of quartz, fluorite, galena, pyrite, K-feldspar. The vein is vuggy. Locally with calcite and gypsum crystals.	
					Sample S59-82-188' (57.3 m) for opaques				
					Sample S59-82-196' (59.7 m) for opaques				
212.0	239.0	64.6	72.8		Medium grained (2-4 mm) leuco-granodiorite gneiss with less than 2% pyroxene..				
239.0	250.0	72.8	76.2		Medium grained (2 mm) pink to purple aplite with disseminated fluorite.				
250.0	251.0	76.2	76.5		Leuco-granodiorite gneiss.	250.0	76.2	First appearance of pyroxene.	
251.0	286.0	76.5	87.2		White aplite.	279.0	85.0	Vein (2 mm) of quartz and gypsum. 090/44S.	
						281.0	85.6	Disseminated fluorite parallel to foliation.	45
286.0	292.0	87.2	89.0		Fine grained (1 mm) layered pyroxene (10%) gneiss.				
						288.0	87.8	Vein (1 mm) of gypsum. 043/68 NW.	
						289.0	88.1	to 335' unsplit core.	
						292.0	89.0	Disseminated fluorite and pyroxene parallel to foliation.	40
292.0	331.0	89.0	100.9		Medium grained (1-3 mm) leuco-granodiorite gneiss.	302.0	92.1	Vein (1-3 mm) of quartz, pyrite and calcite. 090/63 S.	

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						309.0	94.2	Vein (2 mm) of quartz and pyrite. 340/51 NE.	
						320.0	97.5	Vein (2 mm) of gypsum. 075/72 SE.	
331.0	362.0	100.9	110.3		White aplite with disseminated fluorite.	335.0	102.1	to 345' (105.2 m) Split core.	
						345.0	105.2	to 356' (108.5 m) unsplit core.	
362.0	364.0	110.3	110.9		Medium grained (2-4 mm) leuco-granodiorite gneiss with no mafic minerals.				
364.0	377.0	110.9	114.9		White aplite with disseminated fluorite.				
377.0	386.0	114.9	117.7		Pink biotite (3%) leuco-granodiorite gneiss.	380.0	115.8	Vein. 080/51 SE.	
	386.0		117.7		End of hole.				



<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
<b>Project</b>	<b>Client</b>			Eaglet Mines Ltd.					
<b>Drill hole</b>	Structural analysis							<b>Collar Elevation</b>	886.19 m
<b>Coordinates</b>	S60-82							<b>Inclination</b>	90
	64.5 m south							<b>Length</b>	120.7 m
	338.8 m west							<b>Logged by:</b>	PBR

**Note: Assumed strike of foliation of 90 °. Unsplit core.**

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
0.0	100.0	0.0	30.5		overburden				
100.0	106.0	30.5	32.3		Medium grained (2-3 mm) pyroxene granodiorite and leuco-granodiorite gneiss.				
106.0	114.0	32.3	34.7		Cream coloured slightly gneissic aplite	109.0	33.2	First appearance of pyroxene.	
114.0	124.0	34.7	37.8		Fine grained (1 mm) muscovite-bearing aplitic gneiss.	111.0	33.8	First appearance of fluorite.	
124.0	149.0	37.8	45.4		Cream coloured slightly gneissic aplite				
149.0	159.0	45.4	48.5		Medium grained (2-4 mm) pyroxene (2 to 5%) leuco-granodiorite gneiss.				
159.0	163.0	48.5	49.7		White slightly gneissic aplite.				
163.0	171.0	49.7	52.1		Medium grained (1-2 mm) muscovite-bearing gneiss.				
171.0	183.0	52.1	55.8		White and pink gneissic aplite with disseminated fluorite.	171.0	52.1	Vein (5 mm) of quartz and pyrite. 090/61 S.	
						175.0	53.3	Vein (4 mm) of pyrite, fluorite and quartz. 090/51 S.	
						179.0	54.6	Disseminated fluorite parallel to foliation.	42
183.0	191.0	55.8	58.2		Medium grained (2-4 mm) leuco-granodiorite gneiss.				
191.0	193.0	58.2	58.8		Medium grained (1-2 mm) muscovite gneiss.				
193.0	195.0	58.8	59.4		Cream aplite with disseminated fluorite.				
195.0	196.0	59.4	59.7		Medium grained (1-3 mm) leuco-granodiorite gneiss.				
196.0	197.0	59.7	60.0		White aplite.				

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
197.0	205.0	60.0	62.5		Medium grained (1-3 mm) leuco-granodiorite gneiss.	197.0	60.0	Contact between aplite and granodiorite gneiss truncates foliation (in granodiorite ). 086/61 SE.	
205.0	216.0	62.5	65.8		Medium grained (1-3 mm) muscovite gneiss and aplite gneiss portion	211.0	64.3	Vein (5 mm) of fluorite, gypsum, K-feldspar and quartz. 090/33S.	
216.0	219.0	65.8	66.8		Felsic dike with no fluorite (?)	217.0	66.1	Sample S60-82-217' (66.1 m).	
219.0	224.0	66.8	68.3		Medium grained (1-3 mm) muscovite gneiss.		0.0		
224.0	281.0	68.3	85.6		Pink aplite and aplitic gneiss with local pyroxene and fluorite disseminations. Some muscovite gneiss slivers.	235.0	71.6	Vein (3 mm) of quartz and fluorite. 300/53 SW.	
						245.0	74.7	Disseminated fluorite and pyroxene parallel to foliation.	
281.0	288.0	85.6	87.8		Pink medium grained (1-3 mm) muscovite gneiss.				
289.0	312.0	88.1	95.1		Pink slightly gneissic aplite	293.0	89.3	Disseminated fluorite parallel to foliation.	
						293.0	89.3	Vein of gypsum and chalcedony. 340/76 SW.	
						293.0	89.3	Sample S60-82-293' (89.3 m); Chalcedony-quartz vein.	
						296.0	90.2	to 306' (93.3 m) split core.	
						306.0	93.3	to 336' (102.4 m) unsplit core	
312.0	315.0	95.1	96.0		Pink medium grained (1-2 mm) muscovite-bearing gneiss.				
315.0	326.0	96.0	99.4		Pink slightly gneissic aplite				
326.0	342.0	99.4	104.2		Medium grained (2-4 mm) leuco-granodiorite gneiss.	334.0	101.8	Vein (2 mm) of gypsum. 245/75 SW.	
						335.0	102.1	Vein (3 mm) of gypsum. 305/90	
						336.0	102.4	to 386' (117.7 m) split core.	
342.0	377.0	104.2	114.9		White to pink aplite. Disseminated fluorite.	354.0	107.9	Disseminated fluorite parallel to foliation.	
377.0	396.0	114.9	120.7		Medium grained (2 mm) biotite (15%?) pyroxene (5%) granodiorite gneiss. Sample S60-82-385.5' (117.5 m) typical.	386.0	117.7	to 396' (120.7 m) unsplit core.	
	396.0		120.7		End of hole.				

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
<b>Project</b>	<b>Client</b>			Eaglet Mines Ltd.				<b>Collar Elevation</b>	888.44 m
<b>Drill hole</b>	Structural analysis							<b>Inclination</b>	90
<b>Coordinates</b>	S61-82							<b>Length</b>	129.8 m
	60.4 m south							<b>Logged by:</b>	PBR
	368.4 m west								

**Note: Assumed strike of foliation of 90°. Unsplit core to 186'.**

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
0.0	99.0	0.0	30.2		Overburden				
99.0	106.0	30.2	32.3		Altered (sericite) aplite gneiss	102.0	31.1	First appearance of fluorite.	
106.0	131.0	32.3	39.9		Medium grained (2-4 mm) pyroxene (5%) leuco-granodiorite gneiss	109.0	33.2	First appearance of pyroxene.	
						112.0	34.1	First appearance of gypsum vein	
131.0	136.0	39.9	41.5		Pink pyroxene- fluorite bearing aplite				
136.0	161.0	41.5	49.1		Medium grained (1-2 mm) pink muscovite-bearing aplitic gneiss.				
161.0	166.0	49.1	50.6		Cream aplitic gneiss				
166.0	169.0	50.6	51.5		Medium grained (1-3 mm) pyroxene (3%) leuco-granodiorite gneiss				
169.0	186.0	51.5	56.7		Medium grained (1-2 mm) muscovite-bearing pink aplitic gneiss.				
186.0	197.0	56.7	60.0		Cream to pink pegmatitic aplite with fluorite.	186.0	56.7	to 196' (59.7 m) split core.	
						196.0	59.7	to 230' (70.1 m) unsplit core.	
197.0	226.0	60.0	68.9		Cream and pink aplitic gneiss with thin muscovite gneiss sections.				
226.0	239.0	68.9	72.8		Pink to white aplite with disseminated fluorite	228.0	69.5	Vein (2 mm) of K-feldspar. 090/25S.	
						230.0	70.1	to 240' (73.2 m) split core.	
239.0	249.0	72.8	75.9		Medium grained (2-4 mm) pyroxene (2%) leuco-granodiorite gneiss	240.0	73.2	Vein (2 mm) of gypsum. 090/57 S.	
						240.0	73.2	to 349' (106.4 m) unsplit core	
						247.0	75.3	Vein (1 mm) of gypsum. 072/65 NW.	
						247.0	75.3	Vein (2 mm) of quartz. 280/48 SW.	

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
249.0	264.0	75.9	80.5		Fine grained (1 mm) biotite (5-20%) aplitic gneiss locally altered to fine grained muscovite-bearing aplitic gneiss.	257.0	78.3	Disseminated pyroxene parallel to foliation.	
264.0	270.0	80.5	82.3		White aplite with disseminated fluorite, pyroxene and pyrite.				
270.0	275.0	82.3	83.8		Medium grained (1-3 mm) pyroxene (2-10%) granodiorite gneiss				
275.0	301.0	83.8	91.7		Pink aplite gneiss locally altered to pink muscovite-bearing aplitic gneiss.				
301.0	312.0	91.7	95.1		Medium grained (2-4 mm) leuco-granodiorite gneiss	302.0	92.1	Vein (3 mm) of gypsum. 090/66 S.	
312.0	334.0	95.1	101.8		Pink to white gneissic aplite.	317.0	96.6	Vein (2 mm) of quartz. 090/44 S.	
						323.0	98.5	Disseminated fluorite and pyroxene parallel to foliation.	53
334.0	341.0	101.8	103.9		Medium grained (1-3 mm) pyroxene (5%) leuco granodiorite gneiss	337.0	102.7	Sample S61-62-337' (102.7 m).	
341.0	349.0	103.9	106.4		Pink gneissic aplite.				
349.0	368.0	106.4	112.2		Pink to white aplite with fluorite	349.0	106.4	to 359' (109.4 m) split core	
		0.0				359.0	109.4	to 385' (117.3 m) unsplit core	
368.0	370.0	112.2	112.8		Medium grained (2-4 mm) leuco granodiorite gneiss				
370.0	372.0	112.8	113.4		White gneissic aplite.				
372.0	381.0	113.4	116.1		Medium grained (2-4 mm) leuco granodiorite gneiss				
381.0	400.0	116.1	121.9		White aplite with disseminated fluorite,.	385.0	117.3	to 405' (123.4 m) split core	
400.0	403.0	121.9	122.8		Medium grained (1-3 mm) pyroxene (15%) granodiorite gneiss				
403.0	406.0	122.8	123.8		White aplite with disseminated fluorite,.				
406.0	426.0	123.8	129.8		Medium grained (1-3 mm) pyroxene granodiorite gneiss				
	426.0		129.8		End of hole.				

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
<b>Project</b>	<b>Client</b>			Eaglet Mines Ltd.				<b>Collar Elevation</b>	905.62 m
<b>Drill hole</b>	Structural analysis							<b>Inclination</b>	90
<b>Coordinates</b>	S62-82							<b>Length</b>	160.3 m
	3.7 m north							<b>Logged by:</b>	JAF
	348.7 m west								

**Note: Assumed strike of foliation of 90 °. Most core is intact, but has been split where noted.**

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
0.0	52.0	0.0	15.8		Overburden				
52.0	90.0	15.8	27.4	Snow-shoe	Biotite schist; abundant muscovite and local garnet. Local carbonate veins present	70.0	21.3	Vein (1 mm) of calcite and minor pyrite. 160/72 NE.	
90.0	125.0	27.4	38.1	QLG	Pink to cream aplite and muscovite-bearing aplite gneiss with lesser pyroxene granodiorite gneiss. Pyroxene alteration obvious. Disseminated fluorite.	108.0	32.9	Large pyroxene crystals distributed parallel to S1. Foliation	60
						119.0	36.3	Vein -fracture with gypsum. Slicks: 37 -> SW; vein: 042/40 SE.	
125.0	197.0	38.1	60.0		Pink aplitic gneiss with abundant muscovite -bearing aplite gneiss. Granodiorite gneiss absent.	131.0	39.9	Vein (1 mm) of quartz with small amount of gypsum coating. Vein: 127/53 SW; slicks: 39 -> W	
						136.0	41.5	Vein (5 mm) of barren quartz. 092/38S.	
						141.0	43.0	Vein (1-1.5 cm) of fluorite, pyrite, K-feldspar and minor pyroxene. 015/76 NW.	
						144.0	43.9	Vein (2 mm) of fluorite and calcite. 115/37 SW.	
						147.0	44.8	Vein (2 mm) of K-feldspar, calcite and pyrite. 085/47 S.	
						150.0	45.7	Vein ( 1 cm) of barren quartz. 084/38 S.	
						150.0	45.7	Vein ( 2 mm) of K-feldspar, pyrite. 084/38 S.	
						156.0	47.5	Vein of quartz, K-feldspar and pyrite. 080/33 SE.	
						162.0	49.4	Vein-fracture (2 mm) of gypsum. 068/72 SE.	
						166.0	50.6	to 186° (56.7 m) Core is split.	

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						188.0	57.3	Vein of K-feldspar, fluorite and calcite. 102/48SW.	
						191.0	58.2	Two veins of K-feldspar, quartz, fluorite, pyrite and calcite. 082/56S.	
197.0	320.0	60.0	97.5		Pink aplite gneiss with pyroxene alteration and local zones of pyroxene (5%) granodiorite gneiss (5-10 cm thick).	212.0	64.6	Five veins (2 mm wide) of quartz, pyrite and minor fluorite. 078/45 SE.	
						225.0	68.6	Zones of disseminated fluorite and pyroxene parallel to foliation.	45
						225.0	68.6	Vein (3 mm) of K-feldspar, pyrite and fluorite. 066/50 SE.	
						238.0	72.5	Vein -fracture ( 1 mm) of gypsum. 060/78 SE.	
						249.0	75.9	Vein (2 mm) of gypsum with slicks. Vein: 168/79 NE; slicks: 65 ->SE.	
						267.0	81.4	Vein ( 2-3 mm) of barren quartz. 107/65 SW.	
						281.0	85.6	Vein-fracture with gypsum. 088/72 NW.	
						283.0	86.3	Vein (3 mm) of quartz, calcite and minor fluorite. 105/32 SW.	
						296.0	90.2	Vein (4 cm) of calcite and fluorite. 065/50 NW.	
						299.0	91.1	Vein-fracture (2 mm) with gypsum. 097/53 S.	
						313.0	95.4	Vein (5 mm) of quartz, pyrite and minor fluorite. 118/56 SW.	
320.0	526.0	97.5	160.3		Pyroxene granodiorite gneiss with layers of pale aplite and aplite gneiss. Pyroxene and fluorite alteration common.	321.0	97.8	Zones of disseminated pyroxene and fluorite parallel to foliation.	55
						334.0	101.8	Vein (1 mm) of gypsum. 058/85 NW.	
						341.0	103.9	Two veins (3-4 mm wide) of gypsum. 072/45 SE.	
						345.0	105.2	Three veins (3- 4 mm wide) of barren quartz. 015/53SE.	
						358.0	109.1	Vein (2 mm) of gypsum. 090/45 S.	

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						389.0	118.6	Fault- 4' (1.2 m) of core in gap-possible breccia fragments (30 cm).	
						401.0	122.2	Vein (3 mm) of gypsum and quartz. 095/58 S.	
						411.0	125.3	Vein-fracture of gypsum. 089/47S.	
						421.0	128.3	Three veins-fractures (1-2 mm wide) with gypsum. 119/45 SW.	
						436.0	132.9	to 450' (137.1 m) core has been split.	
						460.0	140.2	Vein (2 mm) of gypsum. 173/75 SE.	
						464.0	141.4	Vein of gypsum parallel to foliation. Slickensides: 45 -> N.	45
						468.0	142.6	Two veins (2-3 mm wide) with gypsum. Vein: 078/54 NW; slicks: 42 -> 056.	
						486.0	148.1	Vein (3-4 mm) of gypsum. 062/68 SE; slicks: -> down dip.	
						494.0	150.6	Vein of calcite, fluorite, fluorite and K-feldspar. 156/46 SW.	
						503.0	153.3	Vuggy calcite vein (2 cm). Open cavity with euhedral calcite. 096/46N.	
						510.0	155.4	Vein (5 mm) of fluorite, calcite, pyrite and sphalerite. 145/15 NE.	
						515.0	157.0	Zones (3 cm wide) of disseminated fluorite, calcite and pyroxene parallel to foliation.	45
						520.0	158.5	Vein -fracture (1 mm) of gypsum. 080874 N.	
	526.0		160.3		End of hole.				

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
				Eaglet Mines Ltd.					
<b>Project</b>	<b>Client</b> Structural analysis							<b>Collar Elevation</b>	878.80 m
<b>Drill hole</b>	S63-82							<b>Inclination</b>	90
<b>Coordinates</b>	90.9 m South 371.6 m West							<b>Length</b>	157.3 ,
								<b>Logged by:</b>	JAF

**Note:** Assumed strike of foliation of 90 °. Core has been only partially split, approximately 98 %. Core is at 293' (89.3 m) in adit #2.

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
0.0	109.0	0.0	33.2		Overburden.				
109.0	123.0	33.2	37.5		Pink aplite gneiss with thin muscovite zones.	114.0	34.7	Vein (4 mm) of fluorite, muscovite and K-feldspar parallel to foliation.	40
123.0	145.0	37.5	44.2		Medium grained pyroxene (10%) granodiorite gneiss with thin (< 10 cm) pink aplite, non-gneissic.	123.0	37.5	Vein (3 mm) of quartz, fluorite and sphalerite 107/47 SW.	
						131.0	39.9	Vein (2 -3 mm) of gypsum. 028/74 SE.	
						136.0	41.5	Vein (3 mm) of barren quartz. 095/41 S.	
145.0	170.0	44.2	51.8		Pink aplite and muscovite aplite gneiss with thin zones of pyroxene granodiorite gneiss (15 cm). Granodiorite is minor (5%). Muscovite aplite is fine grained (1 mm).	151.0	46.0	Vein (10 cm) of fluorite and calcite. 175/51 E.	
						169.0	51.5	Vein (2-3 mm) of pyrite, fluorite and quartz. 140/43 SW.	
170.0	191.0	51.8	58.2	pyroxene alteration zone	Medium grained pyroxene granodiorite gneiss and fine grained pink biotite-pyroxene (?) aplite (No muscovite).	180.0	54.9	Vein of quartz. 045/46 SE.	
191.0	230.0	58.2	70.1		Gneiss- some pyroxene granodiorite zones (15-20 cm wide) which are associated with cream to white aplite (with pyroxene?) 10 to 20 cm wide.	196.0	59.7	Vein (2 mm) of quartz and fluorite	
						196.0	59.7	Vein (3 mm) of quartz, fluorite and sphalerite 135/40 SW.	
						198.0	60.4	Vein (6 mm) of K-feldspar. 125/52 SW.	
						214.0	65.2	Vein-fracture (2 mm) of gypsum. 095/64 S.	



<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>					
						218.0	66.4	Two veins(1-2 cm) of quartz and fluorite. 090/35S.						
						221.0	67.4	Vein (2 mm) of gypsum. 063/78SE.						
						226.0	68.9	Vein (2 mm) of gypsum with slickensides. Vein : 107/66 NE; Slicks: 28 ->100.						
230.0	250.0	70.1	76.2		Medium grained pyroxene (10%) granodiorite gneiss with local pink aplite and minor aplite gneiss with local muscovite-bearing zones (muscovite is medium grained (1-2 mm); but appears to be coarse grained variety of muscovite aplite gneiss).	233.0	71.0	Vein (2-3 mm) of quartz, K-feldspar and gypsum alteration. 140/54 SW.						
						250.0	291.0	76.2	88.7	Cream to pink aplite.	266.0	81.1	Vein (2-3 mm) of quartz. Shows minor offset. 066/85 NW.	
											275.0	83.8	Vein (4 mm) of barren quartz. 082/43 S.	
						291.0	295.0	88.7	89.9	Muscovite aplite gneiss. Coarse muscovite in pink aplite, locally well foliated with some non-gneissic zones (5-10 cm).	291.0	88.7	Vein (2-4 mm) of gypsum and minor quartz. 085/50 S.	
						295.0	334.0	89.9	101.8	Medium grained pyroxene (1%) granodiorite gneiss with very thin pink aplite zones (5-10 cm).	295.0	89.9	Vein (4 mm) of pyroxene, fluorite and pyrite. 125/54 SW.	
											304.0	92.7	Fracture-coated with gypsum less than 1 mm. 090/40S.	
											306.0	93.3	Vein-fracture (1 mm) with gypsum. 050/63 SE.	
											318.0	96.9	Vein (1 cm) of barren quartz. 095/40 S.	
											324.0	98.8	Two veins (5 and 10 mm) of quartz, pyrite and gypsum. 035/48 SE.	
											328.0	100.0	to 368' (112.2 m) Core is split.	
334.0	373.0	101.8	113.7		White aplite with local thin pink zones. Local foliation weakly developed.. Fluorite disseminated throughout ("ore zone").	343.0	104.5	Vein-fracture (< 1 mm) with gypsum. 095/64 S.						
											362.0	110.3	Vein (2 mm) of quartz and pyrite. 140/50 SW.	
											369.0	112.5	Vein (3-4 mm) of gypsum. 065/90	

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
373.0	388.0	113.7	118.3	gypsum alteration strong in this interval.	Medium grained granodiorite gneiss with strong pyroxene alteration in 1.5 m zone of mixed granodiorite and cream coloured aplite- going downward into pure granodiorite gneiss at 388' (118.3 m).	375.0	114.3	Vein (3 mm) of gypsum. 087/80 N.	
						377.0	114.9	Vein ( 1 cm) of fluorite and pyroxene. 000/32 E.	
						380.0	115.8	Vein (3-4 mm) of gypsum. 145/73NE.	
388.0	408.0	118.3	124.4		medium grained granodiorite gneiss. Pyroxene alteration strong.	389.0	118.6	Vein (1-5 cm) of fluorite, K-feldspar and quartz. 055/47 SE.	
						390.0	118.9	Vein (2 cm) of quartz and vugs filled with calcite (less than 5 mm in diametre). 135/76 Sw.	
						401.0	122.2	Vein (5-6 mm) of barren quartz. 097/72S.	
						418.0	127.4	Vein (1-2 cm) of pyroxene crystals up to 2 cm in diametre. 035/65 NW.	
408.0	428.0	124.4	130.5		Pyroxene (10-15% granodiorite gneiss, locally with large clusters of coarse grained pyroxene (veins).	425.0	129.5	Two veins (2-5 mm wide) of quartz, K-feldspar and minor gypsum at margins. 015/50 NW.	
						436.0	132.9	Vein (3-4 mm) of dark grey pyroxene parallel to foliation.	50
428.0	467.0	130.5	142.3		Medium grained granodiorite gneiss (pyroxene 5%, locally greater as alteration) with thin zones of non micaceous pink aplite (pyroxene 1 to 2%).	440.0	134.1	Two discontinuous veins (2-4 mm) of fluorite parallel to S1.	50
						440.0	134.1	Vein of quartz, pyrite (cube ) and minor fluorite. 066/58Se.	
						442.0	134.7	Vein (3-4 mm) of pyroxene and minor fluorite and quartz. 020/64 NW.	
						442.0	134.7	Vein (2 mm) of gypsum. 095/55 S.	
						446.0	135.9	Vein (1 cm+) of pyroxene, calcite and minor fluorite. 065/77 NW.	
						446.0	135.9	Vein (2 mm) of calcite and dolomite. 112/62 NE.	

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						453.0	138.1	Vein (4 mm) of K-feldspar, quartz and pyrite. 066/62 SE.	
						459.0	139.9	Vein (3 cm) of large pyroxene crystals (1-2 cm across). 055/63 NW.	
467.0	516.0	142.3	157.3		Medium grained pyroxene (5-10%) granodiorite gneiss with local pyroxene-rich zones and thin zones (15-20 cm) of pink aplite (pyroxene 1-2%).	468.0	142.6	Vein (5 mm) of barren quartz. 025/71 NW.	
						478.0	145.7	Three to four veins (4 mm wide ) of gypsum. 075/67 NW.	
						487.0	148.4	Vein (3 cm) of fluorite and calcite. 014/49 NW.	
						487.0	148.4	Vein (1-2 mm) of gypsum. 104/74 NE.	
						488.0	148.7	Vein (8 mm) of gypsum (?). Sample S63-81-488. 095/52 N.	
						499.0	152.1	Vein (3-4 mm) of calcite and quartz. 085/65S.	
						502.0	153.0	Vein (3 mm) of quartz and pyroxene. 086/60S.	
						509.0	155.1	Vein (2 mm) of calcite and quartz. 090/50S.	
						511.0	155.8	Vein (1 cm) of quartz, calcite/dolomite. 078/63 SE.	
	516.0		157.3		End of hole.				

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
<b>Project</b>	<b>Client</b>			Eaglet Mines Ltd.					
<b>Drill hole</b>	Structural analysis							<b>Collar Elevation</b>	876.19 m
<b>Coordinates</b>	S64-82							<b>Inclination</b>	90
	93.1 m South							<b>Length</b>	145.1 m
	397.5 m west							<b>Logged by:</b>	JAF

**Note: Assumed strike of foliation of 90°. Core is unsplit for the most part. Level of core is 284' in adit #2.**

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
0.0	80.0	0.0	24.4		Overburden				
80.0	135.0	24.4	41.1	QLG	Pink aplite and muscovite aplite gneiss. Locally pyroxene-bearing (2-4 %) with disseminated fluorite common.	97.0	29.6	Vein-fracture (1-2 mm) with gypsum. 085/47S.	
						127.0	38.7	Two veins (4 mm) of quartz, pyrite, molybdenite and minor fluorite. 082/49 S.	
						133.0	40.5	Vein-fracture with greenish gypsum. 047/85SE.	
135.0	173.0	41.1	52.7		Medium grained pyroxene (5%) granodiorite gneiss and lesser pink aplite gneiss with zones of muscovite-bearing aplite gneiss (5-10 cm wide).	140.0	42.7	Vein (5 mm) of quartz and fluorite parallel to foliation.	65
						140.0	42.7	Vein (3 mm) with greenish gypsum. 042/69 SE.	
						145.0	44.2	Vein (2-3 mm) of quartz and pyrite. 075/37SE.	
						163.0	49.7	Vein (3 mm) of quartz and pyrite. 087/47S..	
						163.0	49.7	Vein (4 mm) of gypsum. 002/75E.	
						165.0	50.3	Vein ( 1 cm ) of fluorite and quartz parallel to foliation	55
						172.0	52.4	Pyroxene concentrated in layers, parallel to S1.	55
173.0	186.0	52.7	56.7		Pink aplite non foliated with fine grained pink muscovite-bearing aplite gneiss at base of interval (1.5 m thick).				

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
186.0	270.0	56.7	82.3		Medium grained pyroxene (4-5%) granodiorite gneiss with thin cream to pink muscovite aplite gneiss zones 40-50- cm thick , commonly 10-15 cm thick..	196.0	59.7	Vein (5-6 mm) of fluorite, quartz and pyroxene. 095/40N.	
						201.0	61.3	Vein-fracture (1 mm) coated with gypsum. 032/70SE.	
						221.0	67.4	Vein-fracture (1 mm) coated with gypsum. 110/36SW.	
						222.0	67.7	Vein (2 cm) of quartz, pyroxene, fluorite and minor calcite parallel to foliation.	55
						231.0	70.4	Vein (3 mm) of barren quartz. 113/46 Sw.	
						237.0	72.2	Vein (2 mm) of gypsum. 025/78 NW.	
						258.0	78.6	Fault- gap in core footage. Altered and broken core 8' (2.4 m).	
270.0	281.0	82.3	85.6		Fine grained locally gneissic pink aplite (muscovite <<1%) with some pyroxene (< 1%) and biotite.	271.0	82.6	Vein of fluorite and pyrite parallel to foliation.	45
						274.0	83.5	Vein-fracture (2 mm) of gypsum. 083/64SE.	
281.0	294.0	85.6	89.6		Pink to cream aplite gneiss with very small amount of muscovite-bearing gneiss passing downward into medium grained granodiorite gneiss zones (15-25 cm thick) with green pyroxene and fluorite alteration.	282.5	86.1	Fracture with thin quartz layer and slicks. Vein: 170/72W; slicks: 52 - >NW	
						310.0	94.5	Vein-fracture (2 mm) with gypsum. 076/75 NW.	
315.0	346.0	96.0	105.5		Pink aplite and muscovite aplite gneiss with fine grained pyroxene (1-2%) and lesser medium grained pyroxene granodiorite gneiss in thin zones (20-25 cm).	319.0	97.2	Vein (2 mm) of gypsum. 056/75 NW.	
						324.0	98.8	Vein (1 mm) of quartz and minor pyrite. 178/55 E.	
						329.0	100.3	Vein (2-3 mm) of dolomite and gypsum. 036/43 NW.	

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						338.0	103.0	Vein-fracture (1 mm) with gypsum. 169/62SW.	
346.0	373.0	105.5	113.7		Medium grained granodiorite gneiss with strong green pyroxene alteration; local thin zones of pink unfoliated aplite, 15-20 cm wide and one zone of aplite 1 m wide (between 356'-359' or 108.5-109.5 m).	354.0	107.9	Vein (2 mm) of gypsum. 083/57S.	
						356.0	108.5	to 415' (126.5 m) core is split.	
						361.0	110.0	Vein (6 mm) of calcite and fluorite parallel to foliation.	55
373.0	436.0	113.7	132.9		White aplite with lesser pink aplite containing some coarse muscovite-bearing zones (pegmatitic K-feldspar and muscovite).	427.0	130.2	Vein (4 mm) of quartz and minor gypsum in cavities. 060/45SE.	
						433.0	132.0	Vein (1 mm) of quartz. 017/37 NW.	
436.0	476.0	132.9	145.1		Medium grained granodiorite gneiss with very thin pink aplite (<10 cm). Dark grey pyroxene and muscovite define foliation. Large clusters of pyroxene as alteration.	443.0	135.0	Vein (5-6 mm) of pyroxene and K-feldspar. 094/52N.	
						458.0	139.6	Vein (1mm) of quartz, pyrite and fluorite. 130/470SW.	
						466.0	142.0	Vein (4-6 mm) of fluorite and K-feldspar. 085/20S.	
						468.0	142.6	Vein (1-5 cm) of pyroxene and pyrite. 038/50NW.	
						471.0	143.6	Vein (2 mm) of gypsum. 088/78S.	
	476.0		145.1		End of hole.				

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>	
				<b>Client</b>	Eaglet Mines Ltd.					
<b>Project</b>	Structural analysis								<b>Collar Elevation</b>	873.80 m
<b>Drill hole</b>	S65-82								<b>Inclination</b>	90
<b>Coordinates</b>	85.5 m south 457.7 m west								<b>Length</b>	172.5 m
								<b>Logged by:</b>	JAF	

**Note: Assumed strike of foliation of 90°. Core is mostly unsplit, locally, fluorite-bearing portions are split.**

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
0.0	60.0	0.0	18.3		Overburden.				
60.0	72.0	18.3	21.9	Snow-shoe	Biotite schist in gradational contact with pink aplite gneiss. Thin aplite sills (2-10 cm) with small apophyses visible.				
72.0	110.0	21.9	33.5	QLG	Fine-grained pink muscovite-bearing aplite gneiss.	74.0	22.6	Vein (4-5 mm) of quartz, pyrite and abundant fluorite. 115/51 SW.	
						74.0	22.6	Vein (2 mm) of K-feldspar, molybdenite and pyrite. 100/48 SW.	
						103.0	31.4	Zones (5 cm wide) of coarse grained pyroxene, calcite and fluorite parallel to foliation.	40
110.0	135.0	33.5	41.1		Pink aplite gneiss with thin (15-20 cm) zones of medium grained pyroxene (5-10%) granodiorite gneiss.	112.0	34.1	Vein-fracture (1 mm) with slicks in gypsum. Vein: 118/83 NE; slicks: 75 -> W.	
						124.0	37.8	Vein (2 mm) of gypsum. Vein 025/77 SE; slicks 43 ->E.	
135.0	290.0	41.1	88.4		Pink aplite gneiss with abundant green pyroxene as alteration (up to 15%)	141.0	43.0	Many veins (2 mm to 10 cm wide) of pure fluorite and some pyroxene. Parallel to foliation	50
						141.0	43.0	Vein (3-4 mm) of fluorite, pyroxene and calcite. 128/76NE.	
						152.0	46.3	Vein (4 mm) of calcite and fluorite. 090/38S.	
						152.0	46.3	Vein (1 cm) of pyroxene and fluorite. 110/84 SW.	
						161.0	49.1	Zones of disseminated pyroxene and fluorite parallel to foliation.	50
						166.0	50.6	Vein (8 mm) of fluorite and minor pyroxene. 000/65E.	
						171.0	52.1	Vein (5 mm) of fluorite, calcite and minor pyroxene. 075/57 NW.	

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						176.0	53.6	Vein (6 mm) of pyrite, fluorite, calcite and sphalerite. 125/30 SW.	
						183.0	55.8	Vein (2 mm) of K-feldspar and quartz. 044/42 NW.	
						192.0	58.5	Vein network of gypsum, 2-3 mm wide. 035/82SE.	
						219.0	66.8	Vein (2 mm) of quartz and minor fluorite and pyrite. 090/40S.	
						221.0	67.4	Vein (4-5 mm) of quartz and minor fluorite and pyrite. 115/32Sw.	
						249.0	75.9	Vein (4 m) of gypsum. 090/75N.	
						286.0	87.2	Vein (3 mm) of K-feldspar and fluorite. 089/30S.	
290.0	480.0	88.4	146.3		Pink muscovite-bearing aplite and medium grains pyroxene (5%) granodiorite gneiss (approximately 40% and 60% respectively).	290.0	88.4	Vein (1.5 cm) of fluorite, calcite and minor pyroxene parallel to foliation.	60
						293.0	89.3	Vein (4 mm) of fluorite, pyroxene and pyrite parallel to foliation.	60
						316.0	96.3	Vein (3 mm) of K-feldspar, pyrite and minor pyroxene.	
						326.0	99.4	Vein (2 mm) of K-feldspar, pyrite and minor pyroxene. 087/54S.	
						343.0	104.5	Vein of quartz, fluorite and minor pyrite. 030/22NW.	
						355.0	108.2	Vein (3 mm) of quartz, fluorite and pyrite. 096/42S.	
						355.0	108.2	Vein (5 mm) of quartz and pyrite. Large clusters of pyrite. 006/74E.	
						372.0	113.4	Vein-fracture (1-2 mm) with gypsum. 071/62 NW.	
						407.0	124.1	Vein (2 mm) of quartz and minor pyrite. 005/47E.	
						408.0	124.4	Vein (2 mm) of gypsum with slickensides. Vein: 100/78SW; Slicks: 73 -> SE (east of sip direction).	
						414.0	126.2	Vein (4 mm) of quartz and minor gypsum. 137/42SW.	
						425.0	129.5	Vein (6-8 mm) of quartz and minor fluorite. 130/57 SW.	



<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						429.0	130.8	Vein (2 mm) of gypsum with slickensides. Vein: 112/75SW; Slicks: 15 -> W.	
						457.0	139.3	Disseminated pyroxene and fluorite parallel to foliation.	45
						461.0	140.5	Vein-fracture (1 mm) with gypsum. 079/67SE.	
480.0	550.0	146.3	167.6		Altered interval. Gypsum and potassium metasomatism has rendered much of the section unusable. Foliation is difficult to measure. Rare recognizable pink aplite, white aplite and lesser pyroxene granodiorite gneiss.	480.0	146.3	To end of hole, the core is split.	
						531.0	161.9	Vein (4 mm) of quartz and calcite. 102/67SW.	
550.0	566.0	167.6	172.5		White to pinkish fine grained aplite with local pyroxene (1-2%).	563.0	171.6	Zone of concentrated pyroxene 10-15 cm wide parallel to foliation.	55
	566.0		172.5		End of hole.				

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
				<b>Client</b>	Eaglet Mines Ltd.				
<b>Project</b>	Structural analysis					<b>Collar Elevation</b>	921.04		
<b>Drill hole</b>	S66-82					<b>Inclination</b>	90		
<b>Coordinates</b>	31.7 m south					<b>Length</b>	190.8		
	299.0 m west					<b>Logged by:</b>	P.B.Read		

**Note: Assumed strike of foliation of 90 °.**

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
0.0	17.0	0.0	5.2		Overburden	17.0	5.2	to 32' (9.8 m) unsplit core.	
17.0	44.0	5.2	13.4		Medium grained (1-2 mm) light to medium grey biotite-muscovite schist with 0-10% garnet porphyroblasts (0-6 mm in diametre). Layered (40%) by thin (1.0 m) hornblende schist and amphibolite zones.	22.0	6.7	Vein (3 mm) of calcite. 290/57SW.	
44.0	170.0	13.4	51.8		Medium grained (1-2 mm) light to medium grey biotite-muscovite schist with rare garnet porphyroblasts.	45.0	13.7	Vein (2 mm) of buff calcite. 305/48 SW.	
						70.0	21.3	to 75.5' (7 m) fault gouge and breccia.	
						80.0	24.4	Vein (60 cm) of quartz, dolomite and pyrite. Orientation unknown.	
						105.0	32.0	First appearance of gypsum veins.	
						170.0	51.8	to 216' (65.8 m) unsplit core.	
170.0	171.0	51.8	52.1		Fine grained (1 mm) biotite (15%) aplitic gneiss sill.				
171.0	172.8	52.1	52.7		Medium grained (2 mm) biotite-muscovite schist.				
172.8	175.0	52.7	53.3		Cream muscovite (5%) aplitic gneiss sill	173.0	52.7	Vein (3 cm) of quartz and pyrite. Unknown orientation.	
175.0	183.0	53.3	55.8		Medium grained (2 mm) medium grey biotite-muscovite schist.	175.0	53.3	Vein (2 mm) of gypsum, calcite and K-feldspar. 340/45SW.	
						175.0	53.3	First appearance of K-feldspar.	
						181.0	55.2	Vein (5 mm) of quartz and K-feldspar parallel to foliation.	20
183.0	187.0	55.8	57.0		Fine grained (1 mm) biotite (5-15%) aplitic gneiss .				
187.0	192.0	57.0	58.5		White aplite with disseminated pyrite and 60 cm of aplitic gneiss.	189.0	57.6	Vein (5 mm) of quartz, and fluorite. Unknown orientation.	

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
192.0	216.0	58.5	65.8		Medium grained (2 mm) biotite-muscovite schist.	189.0	57.6	First appearance of fluorite.	
						197.0	60.0	Fault breccia.	
216.0	218.0	65.8	66.4		Pink aplite gneiss.				
			0.0						
218.0	231.0	66.4	70.4		Medium grained (1-3 mm) biotite schist.	218.0	66.4	to 295' (89.9 m). Unsplit core.	
231.0	288.0	70.4	87.8		White aplitic gneiss cut by numerous gypsum veinlets.	282.5	86.1	Vein (2 mm) of gypsum. 090/90	
288.0	302.0	87.8	92.1		White and pink aplite with disseminated fluorite	295.0	89.9	to 300' (91.4 m) Split core.	
302.0	343.0	92.1	104.5		White and pink gneissic aplite locally muscovite-bearing.	300.0	91.4	to 323' (98.5 m). Unsplit core.	
						313.0	95.4	Disseminated fluorite parallel to foliation.	46
						313.0	95.4	Vein (1 mm) of quartz, pyrite and fluorite. 072/41SE.	
343.0	359.0	104.5	109.4		Medium grained (1-2 mm) cream to pink muscovite gneiss.	323.0	98.5	to 343' (104.5 m) Split core.	
						343.0	104.5	to 365' (111.3 m) Unsplit core.	
						353.0	107.6	Vein (10 cm) of K-feldspar and pyrite. 090/43S.	
359.0	364.0	109.4	110.9		Aplitic gneiss				
364.0	365.0	110.9	111.3		Medium grained (1-3 mm) leucocratic granodiorite gneiss				
365.0	380.0	111.3	115.8		White to pink, slightly gneissic aplite with disseminated fluorite	365.0	111.3	to 370' (112.8 m) Split core	
380.0	400.0	115.8	121.9		Medium grained (1-2 mm) muscovite-bearing gneiss and aplitic muscovite-bearing gneiss.	370.0	112.8	to 406' (123.8 m) Unsplit core	
						397.0	121.0	Vein (4 mm) of K-feldspar, quartz and pyrite. 090/50S.	
400.0	426.0	121.9	129.8		Pink aplite and aplitic gneiss with disseminated fluorite.	406.0	123.8	to 421' (128.3 m) Split core.	48
						419.0	127.7	Disseminated fluorite parallel to foliation.	
426.0	439.0	129.8	133.8		Medium grained (1-3 mm) muscovite gneiss.	421.0	128.3	to 431' (131.4 m) Unsplit core.	
						431.0	131.4	to 451' (137.5 m) Split core.	
						438.5	133.7	Vein (5 mm) of quartz and K-feldspar. 090/45S.	
439.0	452.0	133.8	137.8		Pink aplite with disseminated fluorite.	451.0	137.5	First appearance of pyroxene.	
452.0	465.0	137.8	141.7		Cream and pink gneissic aplite with disseminated fluorite.	451.0	137.5	to 504' (153.6 m) Unsplit core.	

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
465.0	477.0	141.7	145.4		Pink muscovite-bearing gneissic aplite.	467.0	142.3	Vein (2 mm) of pyrite and K-feldspar. 245/60 SW.	
477.0	504.0	145.4	153.6		Medium grained (1-3 mm) pink muscovite gneiss and pink aplitic gneiss zones.	485.0	147.8	to 490' (149.4 m) extensive gypsum veinlets.	
						503.5	153.5	Vein (1 mm) of gypsum parallel to foliation	58
						504.0	153.6	to 574' (175 m) split core.	
504.0	574.0	153.6	175.0		White and minor pink aplite with disseminated fluorite. Sample S66-82-519' (158.2 m): Thin section of white aplite. Sample S66-82-543' (165.5 m) Thin section of white aplite.	574.0	175.0	To 626' (190.8 m) unsplit core.	
574.0	595.0	175.0	181.4		Cream to pink fine grained (1 mm) aplitic gneiss with pyroxene alteration zones. Sample S66-85-583' (177.7 m). Alteration pyroxene Thin section.	583.0	177.7	Disseminated pyroxene parallel to foliation.	50
						586.0	178.6	Vein (2 mm) of gypsum. 065/45SE.	
						587.0	178.9	Vein (1 mm) of gypsum. 085/65SE.	
						587.0	178.9	Vein (1 mm) of K-feldspar and pyrite. 285/61SW.	
						594.0	181.1	Vein (3 mm) of pyrite, calcite and quartz. 064/53 SE.	
595.0	625.0	181.4	190.5		Medium grained (1-3 mm) pyroxene (5%) leuco granodiorite gneiss.	596.0	181.7	Vein (4 mm) of pyrite and quartz. 073/50SE.	
						610.0	185.9	Vein (2 mm) of gypsum. 090/50S.	
						614.0	187.1	Vein (15 mm) of vuggy calcite and quartz. 090/35S.	
625.0	626.0	190.5	190.8		Pink aplitic gneiss with disseminated pyroxene and fluorite.				
	626.0		190.8		End of hole.				

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>	
				<b>Client</b>	Eaglet Mines Ltd.					
<b>Project</b>	Structural analysis								<b>Collar Elevation</b>	919.15 m
<b>Drill hole</b>	S67-82								<b>Inclination</b>	90
<b>Coordinates</b>	36.1 m north								<b>Length</b>	195 m
	292.0 m west								<b>Logged by:</b>	P.B.Read

**Note: Assumed strike of foliation of 90 °.**

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
0.0	20.0	0.0	6.1		Overburden				
20.0	24.0	6.1	7.3		Dark green biotite-bearing hornblende schist.	20.0	6.1	to 250' (76.2 m) Unsplit core.	
24.0	180.0	7.3	54.9		Light to medium grey, medium grained (2 mm) biotite-muscovite schist with local garnet porphyroblasts. Rare thin (20 cm) muscovite aplite layers.	112.0	34.1	Vein (10 cm) of biotite, quartz and K-feldspar parallel to foliation	10
						126.0	38.4	First appearance of gypsum in vug.	
						154.0	46.9	Vein (6 cm) of quartz, biotite and K-feldspar.	
180.0	184.5	54.9	56.2		Heavily sericitized bleached Showshoe mica schist	184.0	56.1	First appearance of fluorite.	
184.5	244.0	56.2	74.4		Pink to white muscovite-bearing aplite with disseminated fluorite	199.0	60.7	Vein (3 mm) of K-feldspar. 050/26SE.	
						199.0	60.7	Vein (6 cm) of K-feldspar and fluorite. 050/20SE.	
						202.0	61.6	Vein (1 mm) of K-feldspar and fluorite. 090/39 S.	
						204.0	62.2	Vein (2 mm) of gypsum. 050/39SE.	
						204.0	62.2	Vein (1 mm) of fluorite and K-feldspar. 090/39S.	
						211.0	64.3	Disseminated fluorite parallel to foliation	45
						221.0	67.4	Vein (2 mm) of fluorite, quartz and K-feldspar. 090/65S.	
244.0	251.0	74.4	76.5		Medium grained (1-3 mm) pink muscovite-bearing gneiss	245.0	74.7	Vein (3 mm) of K-feldspar, pyrite and galena. 063/36SE.	
251.0	260.0	76.5	79.2		Pink to white aplite with disseminated fluorite and pyroxene.	250.0	76.2	to 260' (79.2 m) split core.	

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
260.0	270.0	79.2	82.3		Medium grained (1-3 mm) muscovite-bearing gneiss	260.0	79.2	to 427' (130.1 m) Unsplit core.	
						265.0	80.8	Vein (3 mm) of K-feldspar and pyrite. 335/34SW.	
270.0	313.0	82.3	95.4		Pink to white locally, muscovite-bearing aplitic gneiss.	286.0	87.2	Disseminated fluorite parallel to foliation	43
						296.0	90.2	Vein (1 mm) of pyrite and fluorite. 075/45SE.	
						300.0	91.4	Vein (1-2 mm) of pyrite and fluorite. 052/48SE.	
						303.0	92.4	Vein (2 mm) of galena, pyrite, fluorite and K-feldspar. 090/40S.	
						310.0	94.5	Disseminated fluorite parallel to foliation	49
313.0	321.0	95.4	97.8		Medium grained (1-2 mm) leuco-granodiorite gneiss. Sample S61-82-318' (96.9 m).	318.0	96.9	Vein (15 mm) of chalcedony (?). 297/30SW.	
						320.0	97.5	Definitive alteration pyroxene.	
321.0	341.0	97.8	103.9		Pink to cream gneissic aplite	324.0	98.8	Vein (2 mm) of quartz and pyrite. 290/30SW.	
						330.0	100.6	Disseminated fluorite parallel to foliation	55
						330.0	100.6	Vein (2 mm) of pyrite and quartz. 090/45S.	
						331.0	100.9	Vein (2 mm) of pyrite and quartz. 090/34S.	
341.0	358.0	103.9	109.1		Medium grained (2-4 mm) pyroxene (10%) leuco-granodiorite gneiss. Muscovite gneiss between 345.7 and 347' (105.4-105.8 m). Sample S67-82-354' (107.9 m) typical for thin section	342.0	104.2	Disseminated fluorite parallel to foliation	53
						362.0	110.3	Disseminated fluorite parallel to foliation	52
						391.0	119.2	Vein (6 mm) of pyrite, fluorite and K-feldspar. 045/28SE.	
						399.0	121.6	to 400' vein of quartz, galena and pyrite. Unknown orientation.	
						421.0	128.3	Vein (2 mm) of pyrite and K-feldspar. 090/33S.	
						427.0	130.2	to 432' (131.7 m) Split core.	
						432.0	131.7	to 469' (143 m) Unsplit core.	

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						432.0	131.7	Vein (10 mm) of quartz, K-feldspar and pyrite. 090/47S.	
						459.0	139.9	Vein (4 mm) of quartz and pyrite. 078/40SE.	
						466.0	142.0	Vein (3 mm) of gypsum. 090/78S.	
						469.0	143.0	to 479' (146 m) Split core	
						479.0	146.0	Vein (1 mm) of quartz and pyrite. 295/27SW.	
						479.0	146.0	to 535' (163 m) Unsplit core.	
						481.0	146.6	Vein (1-3 mm) of pyrite, fluorite and quartz. 090/37S.	
					Sample S67-82-484' (147.5 m). Vein truncated by shear zone. Thin section.	484.0	147.5	Vein (4 mm) of quartz, pyrite and fluorite. 090/45S.	
						488.0	148.7	Vein (2 mm) of gypsum. 285/50SW.	
						493.0	150.3	Disseminated fluorite parallel to foliation	48
						493.0	150.3	Vein (1 mm) of quartz, pyrite and K-feldspar. 065/45SE.	
						496.0	151.2	Vein (6 mm) of sphalerite, quartz, galena and pyrite. 070/35 SE.	
						500.0	152.4	Vein (2 mm) of pyrite, quartz and K-feldspar. 290/29SW.	
514.0	529.5	156.7	161.4		Medium grained (1-3 mm) muscovite gneiss.	520.0	158.5	Disseminated K-feldspar and fluorite parallel to foliation	46
						520.0	158.5	Vein (8 cm) of K-feldspar, quartz and pyrite. 335/40SW.	
						524.0	159.7	Vein (3 mm) of chalcedony quartz. 300/55SW.	
						525.0	160.0	Disseminated fluorite and K-feldspar parallel to foliation	47
						525.0	160.0	Vein (4 mm) of K-feldspar and pyrite. 060/37SE.	
						525.0	160.0	Vein (2 mm) of gypsum. 355/90.	
						528.0	160.9	Vein (2 mm) of pyrite and K-feldspar. 053/37SE.	
529.5	560.0	161.4	170.7		Mixed pink aplite and pink muscovite gneiss. Sample S67-82-552' (168.2 m). Thin section for ID.	534.0	162.8	Vein (3 mm) of pyrite, K-feldspar and quartz. 090/38S.	
						535.0	163.1	To 545' (166.1 m) Split core	
						536.0	163.4	Disseminated K-feldspar and fluorite parallel to foliation	55

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						536.0	163.4	Vein (3 mm) of pyrite, K-feldspar, quartz and calcite. 285/50 SW.	
						545.0	166.1	To 561' (171 m) Unsplit core.	
						547.0	166.7	Vein (3 mm) of K-feldspar, galena and pyrite. 090/42 S.	
						552.0	168.3	Vein (1 cm) of chalcedony quartz. 090/52 S.	
560.0	570.0	170.7	173.7		Medium grained (2-4 mm) pink muscovite gneiss.	561.0	171.0	To 641' (195.4 m) Split core.	
570.0	612.0	173.7	186.5		White aplite with coarse disseminated fluorite, and local pink aplite with fluorite. Many gypsum veins.				
612.0	625.0	186.5	190.5		Medium grained (2-3 mm) leuco granodiorite gneiss with many gypsum veins.				
625.0	638.0	190.5	194.5		White aplite with disseminated fluorite.				
638.0	641.0	194.5	195.4		Medium grained leuco granodiorite gneiss with many gypsum veins.				
	641.0		195.4		End of hole.				



<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>	
				<b>Client</b>	Eaglet Mines Ltd.					
<b>Project</b>	Structural analysis								<b>Collar Elevation</b>	869.9 m
<b>Drill hole</b>	S82-82								<b>Inclination</b>	90
<b>Coordinates</b>	156.5 m south								<b>Length</b>	102.4 m
	301.3 m west								<b>Logged by:</b>	JAF

**Note:** Assumed strike of foliation of 90°. Core has been split only where noted.. Level of core in adit #2 is 264' (80.5 m).

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
0.0	22.0	0.0	6.7		Overburden				
22.0	95.0	6.7	29.0	QLG. Biotite altera- tion.	Pink aplite and muscovite-bearing aplitic gneiss, altered, with disseminated fluorite and local biotite (biotite "alteration zone"). Some minor pyroxene granodiorite gneiss locally (20 cm or less in thickness).	34.5	10.5	Fault- gap in core of 5' (1.5 m).	
						47.0	14.3	Vein (1-2 mm) of quartz and gypsum coating. 115/90, slicks: 66 -> 295	
	66.0		20.1		Sample S82-82-66, green pyroxene (?), pyrite, fluorite and K-feldspar.	66.0	20.1	Zone of fluorite, pyrite and fine grained green pyroxene (?) parallel to foliation.	50
95.0	110.0	29.0	33.5		Pink aplite and fine grained (1 mm) muscovite-bearing aplitic gneiss. Granodiorite gneiss absent.	100.0	30.5	Vein (5 mm) of K-feldspar, fluorite and calcite. 074/73 SE.	
						119.0	36.3	to 132' (40.2 m) core is split.	
110.0	136.0	33.5	41.5		Pink aplite with disseminated fluorite. No aplitic gneiss whatsoever.	133.0	40.5	Vein of pyrite, quartz and gypsum. 070/68SE.	
136.0	175.0	41.5	53.3		Biotite as alteration; large biotite flashes and smaller clusters of biotite in random orientation. Biotite is frequently associated with fluorite in veins. The rock is pink aplite with abundant fine grained muscovite-bearing aplite. No granodiorite gneiss, no pyroxene alteration. Muscovite aplite is not muscovite aplite gneiss with coarse muscovite.	136.0	41.5	Vein (4 cm) of massive fluorite and calcite. 110/56 NE.	

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						144.0	43.9	Vein (1 cm) of biotite, fluorite and calcite. 042/54 NW.	
						161.0	49.1	Vein (12 cm) of fluorite and calcite parallel to S1. 110/27 NE.	
175.0	199.0	53.3	60.7		Pink to cream aplite and muscovite aplite gneiss with minor granodiorite gneiss (< 10 cm wide). Pyroxene alteration is present. Also disseminated fluorite and veins of fluorite.	195.0	59.4	Vein (2 mm) of gypsum with slickensides. Vein: 058/53 NW; slicks: down dip.	
199.0	215.0	60.7	65.5		Pink aplite with fine grained pyroxene as alteration.. No muscovite gneiss or muscovite aplite.	211.0	64.3	Vein-fracture (1 mm) with gypsum. 142/68 NE.	
						213.0	64.9	Vein (3 mm) of gypsum. 100/65 SW.	
215.0	225.0	65.5	68.6	gypsum alteration strong at 220 feet.	Pyroxene (< 2%) granodiorite gneiss with cream to pink aplite subordinate (25-30%).	221.0	67.4	Vein (3 mm) of barren quartz. 080/55 SE.	
225.0	230.0	68.6	70.1		Cream coloured aplite gneiss with disseminated fluorite and fine grained pyroxene (?).	227.0	69.2	Vein (2-3 mm) of gypsum with slickensides. Vein: 095/52S. Slicks: dip slip.	
230.0	264.0	70.1	80.5		Cream coloured aplite gneiss with disseminated fluorite and fine grained pyroxene (?).	236.0	71.9	Vein (1 cm) of fluorite, K-feldspar and calcite parallel to foliation.	
						248.0	75.6	Vein of K-feldspar, pyrite, calcite and molybdenite. 078/28SE.	
						256.0	78.0	Vein of fluorite, calcite and pyroxene. 060/54 NW.	
						261.0	79.6	Vein-fracture (1 mm) of gypsum. 125/46 SW.	
264.0		80.5			Adit #2. Medium grained leuco granodiorite gneiss well foliated. No aplite associated with granodiorite at this level.	269.0	82.0	Vuggy calcite (3-4 mm wide) sub-parallel to foliation.	65
						275.0	83.8	Vein (2-3 mm) of quartz and rare pyrite cubes. 093/32S.	
						277.0	84.4	Vein (3 cm) of quartz with fine grained pyrite at margins. Parallel to foliation.	35

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
284.0		86.6		Pyroxene and fluorite alteration	Granodiorite gneiss continues with decrease in aplite component (<15%). The granodiorite is medium grained to coarse grained, strongly foliated. Pyroxene alteration widespread at 284 feet. Thin aplites (pink, non foliated), 10-15 cm wide. At 292 feet granodiorite is locally fined grained (1 mm) with abundant pyroxene and some biotite.	286.0	87.2	Vein (4-5 mm) of pyroxene, K-feldspar and quartz. 060/43 NW.	
						290.0	88.4	Vein (5 mm) of pyroxene, calcite and fluorite parallel to foliation.	40
						298.0	90.8	Vein (1 cm) of pyroxene, K-feldspar and minor fluorite. 045/57 NW.	
						310.0	94.5	Zones (1 cm wide) of foliated biotite and calcite parallel to foliation.	40
						322.0	98.1	Vein (6 mm) of quartz, pyrite and K-feldspar parallel to foliation.	55
						335.0	102.1	Zones of coarse grained green pyroxene in vein 2-3 mm wide parallel to foliation.	46
						335.0	102.1	Vein-fracture (2 mm) of gypsum. 115/42SW.	
	336.0		102.4		End of hole.				

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
					Eaglet Mines Ltd.				
<b>Project</b>	Client Structural analysis							<b>Collar Elevation</b>	756.6 m
<b>Drill hole</b>	S85-82							<b>Inclination</b>	90
<b>Coordinates</b>	323.7 m south 696 m west							<b>Length</b>	111.5 m
								<b>Logged by:</b>	P.B. Read

**Note: Assumed strike of foliation of 054 °. Core mainly unsplit.**

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
0.0	17.0	0.0	5.2		Overburden				
17.0	366.0	5.2	111.6		Cream to locally pink aplitic gneiss	26.0	7.9	Vein (6 mm) of fluorite and pyrite parallel to foliation.	55
						29.0	8.8	Vein (4 cm) of fluorite and pyroxene parallel to foliation.	45
						38.0	11.6	Vein of quartz. 315/27 SW.	
						38.0	11.6	Vein (2 mm) of kaolinite. 082/45 NW.	
						46.0	14.0	Vein (2 mm) of kaolinite. 045/75 SE.	
						58.0	17.7	Vein (2 mm) of fluorite and pyroxene parallel to foliation.	60
						61.0	18.6	Vein (3 mm) of fluorite parallel to foliation.	60
						65.0	19.8	Vein (1 cm) of fluorite parallel to foliation.	50
						76.0	23.2	Vein (1-2 cm) of fluorite and pyroxene parallel to foliation.	55
						76.0	23.2	Vein (1 mm) of kaolinite parallel to foliation.	55
						90.0	27.4	Vein (1 cm) of pyroxene and fluorite parallel to foliation.	45
						104.0	31.7	Vein (3-5 mm) of kaolinite. 060/90.	
						107.0	32.6	Vein (6 mm) of fluorite and pyrite. 010/55SE.	
						118.0	36.0	Vein (5 mm) of K-feldspar, fluorite, pyrite and pyroxene. 090/47S.	
						120.0	36.6	Vein (1.5 cm) of K-feldspar and fluorite parallel to foliation.	55
						148.0	45.1	Vein (1.5 cm) of pyrite and fluorite. 310/38 SW.	
						156.0	47.5	Pyroxene and fluorite parallel to foliation.	60

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						200.0	61.0	Vein parallel to foliation.	55
						200.0	61.0	Vein (3 mm) of fluorite . 035/40SE.	
						208.0	63.4	Vein (3 mm) of kaolinite parallel to foliation.	50
						225.0	68.6	Vein (5 mm) of fluorite. 015/72 NW.	
						232.0	70.7	Zones (2 cm wide) of pyrite, K-feldspar and fluorite parallel to foliation.	65
						232.0	70.7	Vein (5 mm) of fluorite and pyroxene. 080/45 SE.	
						236.0	71.9	Zones (4 cm wide) of clinopyroxene fluorite and K-feldspar parallel to foliation.	60
						254.0	77.4	Foliation parallel to contact between aplite (above) and pink muscovite gneiss.	50
						270.0	82.3	Foliation - contact between aplite (above) and pyroxene granodiorite gneiss.	70
						270.0	82.3	Vein (1 mm) of fluorite. 035/48 SE.	
						273.0	83.2	Vein-fracture (1 mm) with kaolinite. 025/70W.	
						270.0	82.3	Fracture (1 mm) filled with kaolinite. 031/76 SE.	
						287.0	87.5	Vein (5 mm) of kaolinite. 010/65SE.	
						308.0	93.9	Vein (4-5 mm) of pyroxene, fluorite and pyrite. 080/67N.	
						308.0	93.9	Vein (1 mm) of kaolinite. 030/70NW.	
						326.0	99.4	Vein (1 mm) of kaolinite. 045/62SE.	
						327.0	99.7	Vein (1 cm) of fluorite and quartz. 130/53 NE.	
						356.0	108.5	Vein (1 mm) of kaolinite. 055/90.	
						365.0	111.3	Vein of kaolinite. Slickensides: 120/34 SE; vein: 020/65SE.	
	366.0		111.6		End of hole.				



<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						219.0	66.8	Fluorite and K-feldspar disseminated parallel to foliation.	
						236.0	71.9	Two veins (1-3 mm) of kaolinite. 030/48SE and 062/35SE.	
						252.0	76.8	Vein (2 mm) of kaolinite. 325/67SW.	
						267.0	81.4	Three veins (5 mm) of kaolinite. 050/62 SE.	
						299.0	91.1	Contact between pyroxene granodiorite gneiss with muscovite gneiss parallel to foliation.	60
						301.0	91.7	Fluorite layers (2 cm thick) parallel to foliation.	45
						311.0	94.8	Fluorite + pyroxene layers (2 cm thick) parallel to foliation.	50
						313.0	95.4	Vein (2 cm) of fluorite parallel to foliation.	60
						324.0	98.8	Vein (3-4 mm) of kaolinite. 045/90.	
						329.0	100.3	Vein (1 cm) of fluorite. 085/50 NW.	
						341.0	103.9	Vein of fluorite, calcite and muscovite parallel to foliation.	40
						356.0	108.5	Vein (5 mm ) of quartz. 080/50SE.	
						373.0	113.7	Vein (5 mm) of pyrite and fluorite parallel to foliation.	55
	396.0		120.7		End of hole.				

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>	
				<b>Client</b>	Eaglet Mines Ltd.					
<b>Project</b>	Structural analysis								<b>Collar Elevation</b>	757.5 m
<b>Drill hole</b>	S87-82								<b>Inclination</b>	90
<b>Coordinates</b>	261.4 m south								<b>Length</b>	153.3 m
	808.2 m west								<b>Logged by:</b>	PBR/JAF

**Note: Assumed strike of foliation of 054 °. Mainly unsplit core.**

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
0.0	25.0	0.0	7.6		Overburden				
25.0	33.0	7.6	10.1		Pink medium grained (2-4 mm) muscovite gneiss.	27.0	8.2	Vein (2-3 mm) of fluorite and pyrite. 030/60SE.	
33.0	503.0	10.1	153.3		Cream to pink slightly gneissic aplite	33.0	10.1	Contact between muscovite gneiss (above) and aplite parallel to foliation.	60
						63.5	19.4	Vein of quartz and calcite parallel to foliation	55
						73.0	22.3	Two veins (1 and 2 cm thick) of pyroxene and fluorite. Parallel to foliation.	45
						73.0	22.3	Vein (2 mm) of pyrite and fluorite. 140/67 SW.	
						86.0	26.2	Vein (1 mm) of kaolinite. 055/67SE.	
						98.0	29.9	Vein (5-6 mm) of quartz. 040/50SE.	
						99.0	30.2	Vein (5 mm) of quartz and pyrite. 055/45SE.	
						121.0	36.9	Disseminated fluorite parallel to S1. Foliation	55
						128.0	39.0	Disseminated fluorite parallel to S1. Foliation	50
						142.0	43.3	Vein (1-2 mm) of gypsum. 060/74 SE.	
						150.0	45.7	Vein (4 cm) of quartz, fluorite and pyrite. 077/62SE.	
						151.5	46.2	Contact between aplite (above) and muscovite gneiss parallel to foliation.	60
						180.0	54.9	Vein (2 mm) of gypsum. 054/60SE.	
						182.0	55.5	Vein (1-3 mm) of fluorite, pyrite and quartz. 054/50SE.	



<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						187.0	57.0	Vein (2 mm) of pyrite and minor fluorite. 055/32SE.	
						197.0	60.0	Vein (3 mm) of kaolinite parallel to foliation.	70
						237.0	72.2	Vein (0.5-1 cm) of fluorite and pyrite parallel to foliation	25
						240.0	73.2	Kaolinite coated fracture. 160/90.	
						245.0	74.7	Vein (5 mm) of quartz. 054/50SE.	
						260.0	79.2	Vein (1 cm) of pyrite and fluorite parallel to foliation.	55
						260.0	79.2	Vein of pyrite, fluorite and quartz. 054/45SE.	
						266.0	81.1	Vein (4-5 mm) of quartz. 015/50SE.	
						272.0	82.9	Vein (2 mm) of kaolinite. 055/49SE.	
						288.0	87.8	Vein (1 cm) of quartz and pyrite. 315/32SW.	
						298.0	90.8	Disseminated fluorite parallel to S1. Foliation	50
						304.0	92.7	Vein (3-4 mm) of kaolinite. 030/56SE.	
						326.0	99.4	Vein (2 mm) of kaolinite. 055/63SE.	
						346.0	105.5	Vein (3 mm) of kaolinite. 045/65SE.	
						364.0	110.9	Vein (1 cm ) of fluorite. 322/60SW.	
						385.0	117.3	Disseminated fluorite and pyroxene parallel to foliation	60
						465.0	141.7	Vein (2 mm) of kaolinite. 025/60SE.	
						470.0	143.3	Vein (3 mm) of kaolinite. 055/90.	
						482.0	146.9	Vein (1 mm) of kaolinite. 055/40SE.	
						486.0	148.1	Vein (5 mm) of quartz, fluorite and K-feldspar. 084/53NW.	
						491.0	149.7	Vein (2-3 mm) of quartz. 055/20SE.	
	503.0		153.3		End of hole.				

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>	
				<b>Client</b>	Eaglet Mines Ltd.					
<b>Project</b>	Structural analysis								<b>Collar Elevation</b>	759.5 m
<b>Drill hole</b>	S88-82								<b>Inclination</b>	90
<b>Coordinates</b>	226 m south 857.4 m west								<b>Length</b>	135.9 m
								<b>Logged by:</b>	PBR/JAF	

**Note: Assumed strike of foliation of 054 °. Mainly unsplit core.**

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
0.0	32.0	0.0	9.8		Overburden				
32.0	47.0	9.8	14.3		Pink gneissic aplite with zones of medium grained (2 mm) muscovite gneiss	41.0	12.5	Vein (5 mm) of galena, fluorite, sphalerite, pyrite and calcite. 000/42E.	
						44.0	13.4	Vein of pyrite, fluorite and carbonate parallel to foliation. Slickensides on vein down dip.	50
47.0	57.0	14.3	17.4		Cream medium grained (2 mm) pyroxene-fluorite-bearing aplite gneiss. <b>No further lithology logging.</b>	51.0	15.5	Disseminated fluorite, pyrite and pyroxene parallel to foliation.	60
						93.0	28.3	Vein (1-2 cm) of pyroxene, fluorite and pyrite parallel to foliation.	45
						101.0	30.8	Vein (2 mm) of kaolinite. 335/90.	
						130.0	39.6	Vein (1 cm) of fluorite, quartz and pyroxene. 050/60 NW	
						145.0	44.2	Vein (2 mm) of kaolinite. 055/85SE.	
						156.0	47.5	Vein (1 mm) of kaolinite. 055/67SE.	
						191.0	58.2	Vein of galena and fluorite parallel to foliation.	30
						196.0	59.7	Horizontal zones (2 cm thick) of fluorite, K-feldspar and pyrite.	
						211.0	64.3	Disseminated fluorite parallel to foliation.	55
						220.0	67.1	Vein (1-2 mm) of kaolinite. 071/65SE.	
						224.0	68.3	Vein (2-3 cm) of fluorite parallel to foliation.	50
						224.0	68.3	Vein (1-2 mm) of kaolinite. 085/55N.	
						260.0	79.2	Zones (2 cm thick) of fluorite, quartz and pyroxene parallel to foliation	60

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						260.0	79.2	vein of kaolinite. 055/60SE.	
						314.0	95.7	Vein (2-3 mm) of kaolinite. 055/60SE.	
						317.0	96.6	Zones (4 cm thick) of disseminated fluorite, quartz, K-feldspar and pyroxene parallel to foliation.	70
						333.0	101.5	Vein of fluorite, pyrite and calcite. 055/50SE.	
						350.0	106.7	Zone of disseminated pyrite and molybdenite parallel to foliation.	60
						367.0	111.9	Vein of pyrite, calcite and fluorite. 045/45SE.	
						367.0	111.9	Vein (1-1.5 cm) of fluorite (90%) and calcite. 045/60NW.	
						372.0	113.4	Vein of galena, sphalerite, pyrite and fluorite. 050/42 SE.	
						406.0	123.8	Vein of fluorite, calcite, pyrite and quartz. 055/60SE.	
						418.0	127.4	Vein of kaolinite. 055/45SE.	
						424.0	129.2	Vein (1 mm) of kaolinite.	
						444.0	135.3	Vein (1 cm) of quartz parallel to foliation.	55
	446.0		135.9		End of hole.				

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
					Eaglet Mines Ltd.				
<b>Project</b>	<b>Client</b>							<b>Collar Elevation</b>	758.5 m
	Structural analysis							<b>Inclination</b>	90
<b>Drill hole</b>	S89-82							<b>Length</b>	138.9 m
<b>Coordinates</b>	197.7 m south							<b>Logged by:</b>	JAF
	901.0 m west								

**Note: Assumed strike of foliation of 054 °. Mainly unsplit core.**

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
0.0	30.0	0.0	9.1		Overburden				
30.0	37.0	9.1	11.3		Cream aplite gneiss				
37.0	41.0	11.3	12.5		Cream, medium grained (2-3 mm) muscovite gneiss.				
41.0	456.0	12.5	139.0		Cream to pink aplite gneiss.	41.0	12.5	Vein (2-5 mm) of fluorite and pyrite. 145/53NE.	
						48.0	14.6	Two horizontal veins (6 mm and 1.5 cm thick) of fluorite	
						61.0	18.6	Two veins (1 cm) of fluorite. 085/37N.	
						63.0	19.2	Vein (5 mm) of quartz and minor fluorite and pyrite. 170/73E.	
						88.0	26.8	Vein (1.5 cm) of fluorite, quartz, sphalerite and pyrite. 105/83N.	
						98.0	29.9	Vein (5-6 mm) of quartz and fluorite parallel to foliation.	45
						101.0	30.8	Vein (1 cm) of fluorite, quartz and pyrite. 145/28NE.	
						112.0	34.1	Vein of fluorite, pyroxene, pyrite and quartz parallel to foliation.	40
						120.0	36.6	Vein (2 cm) of fluorite and pyrite parallel to foliation.	45
						126.0	38.4	Fracture-coated with chlorite, kaolinite and minor pyrite. 175/67E.	
						150.0	45.7	Fracture (2 mm) coated with kaolinite. 040/75 NW.	
						157.0	47.9	Vein (3 cm) of quartz and fluorite. 170/52E.	
						178.0	54.3	Vein of fluorite, calcite and quartz parallel to foliation.	70
						185.0	56.4	Disseminated fluorite in zone 5 to 6 mm wide parallel to foliation.	50
						211.0	64.3	Veins (1-2 mm) of pyrite, fluorite and minor quartz. 085/50SE.	

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						213.0	64.9	Vein (2 mm) of pyrite and fluorite. 055/60SE.	
						229.0	69.8	Vein of quartz and gypsum. 055/50SE.	
						260.0	79.2	Fractures filled with kaolinite (?) and gypsum. 055/50SE.	
						271.0	82.6	Vein (1 cm) of fluorite parallel to S1. Foliation.	45
						297.0	90.5	Vein (1.5-2 cm) of fluorite, pyrite and quartz parallel to foliation	
						320.0	97.5	Three fractures (8 to 10 cm spacing). Chlorite coated. 055/40SE.	
						336.0	102.4	Three veins (2 mm thick) of quartz, pyrite and fluorite. 140/52SW.	
						347.0	105.8	Disseminated fluorite in zones 2 cm wide parallel to S1. Foliation.	50
						351.0	107.0	Vein (2-4 mm) of fluorite and pyrite. 105/42N.	
						356.0	108.5	Disseminated fluorite parallel to foliation.	50
						361.0	110.0	Vein of pyrite, galena and minor fluorite. 015/56E.	
						366.0	111.6	Thin fractures (< 1mm) filled with quartz, 3-4 cm spacing. 160/56 SW.	
						378.0	115.2	Fluorite coated fracture- 1 mm wide, altered (clay). 113/73SW.	
						388.0	118.3	Disseminated fluorite, pyroxene and pyrite parallel to foliation.	40
						429.0	130.8	Vein (2 mm) of fluorite, quartz and pyrite parallel to foliation.	45
						441.0	134.4	Vein (1.5 cm) of fluorite and minor pyrite. 130/40NE.	
						447.0	136.2	Vein (4 mm) of fluorite, quartz and pyrite. 055/54SE.	
	456.0		139.0		End of hole.				

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
<b>Project</b>		<b>Client</b>		Eaglet Mines Ltd.				<b>Collar Elevation</b>	
<b>Drill hole</b>		S111-85						<b>Inclination</b>	90
<b>Coordinates</b>								<b>Length</b>	
								<b>Logged by:</b>	P.B. Read

**Note: Assumed strike of foliation of 090 °. Core has been split in intervals noted.**

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
0.0	80.0	0.0	24.4		Overburden				
80.0	98.0	24.4	29.9		Fine grained (<1 mm) biotite (7%) aplitic gneiss.	97.0	29.6	First appearance of fluorite.	
						97.0	29.6	Vein (2 mm) of fluorite and quartz. 285/41SW.	
98.0	115.0	29.9	35.1		Medium grained (1-2 mm) biotite pyroxene (5-10%) granodiorite gneiss	106.0	32.3	First appearance of pyroxene alteration	
						107.0	32.6	Vein (1 cm) of quartz. 090/51S.	
115.0	118.0	35.1	36.0		Fine grained (< 1 mm) aplitic gneiss.				
118.0	123.0	36.0	37.5		Medium grained (2 mm) cream muscovite gneiss				
123.0	128.0	37.5	39.0		Medium grained (1-2 mm) biotite (5%) leuco granodiorite gneiss				
128.0	131.0	39.0	39.9		Fine grained (<1 mm) biotite (5%) aplitic gneiss.				
131.0	139.0	39.9	42.4		Pink aplite and gneissic aplite with disseminated pyrite and fluorite.				
139.0	141.0	42.4	43.0		Medium grained (1-2 mm) pyroxene (5%) leuco granodiorite gneiss	141.0	43.0	First appearance of gypsum veins.	
141.0	143.0	43.0	43.6		Pink pegmatitic aplite				
143.0	145.0	43.6	44.2		Medium grained (1-3 mm) leuco granodiorite gneiss	144.0	43.9	Vein (1 cm) of quartz and pyrite. 290/90.	
145.0	147.0	44.2	44.8		White aplite with fluorite.				
147.0	153.0	44.8	46.6		Leucocratic gneiss and biotite-bearing aplite gneiss.				
153.0	170.0	46.6	51.8		Medium grained (2-3 mm) leuco granodiorite.				
170.0	181.0	51.8	55.2		Pink gneissic aplite with local disseminated fluorite.				

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
181.0	218.0	55.2	66.4		Medium grained (2-4 mm) pyroxene (1%) leuco granodiorite gneiss and thin (25 cm) cream aplite sill.	192.0	58.5	Vein (1 mm) of gypsum. 040/55SE.	
218.0	220.0	66.4	67.1		Pink and white aplite dike	218.0	66.4	Contact between aplite and gneiss. 070/49SE.	
222.0	246.0	67.7	75.0		Medium grained (2-4 mm) pyroxene (5-10%) leuco granodiorite gneiss.	222.0	67.7	Vein (1 mm) of pyrite. 023/63 SE.	
246.0	250.0	75.0	76.2		Pink to white aplite with disseminated fluorite.				
250.0	255.0	76.2	77.7		Medium grained (1-3 mm) biotite - pyroxene leuco granodiorite gneiss				
255.0	274.0	77.7	83.5		Pink to white aplite with disseminated fluorite and pyrite.				
274.0	288.0	83.5	87.8		Medium grained (1-3 mm) pyroxene (3%) leuco granodiorite gneiss	285.0	86.9	Vein (5 mm) of gypsum. 085/70NW.	
288.0	292.0	87.8	89.0		Fine grained (1 mm) biotite-pyroxene aplitic gneiss.				
292.0	295.0	89.0	89.9		White aplite with fluorite.				
295.0	329.0	89.9	100.3		Medium grained pyroxene (?) - biotite granodiorite gneiss with thin aplite zones. White and minor pink fine grained pyroxene aplite abundant in layers 10-20- cm wide. Finer grained whitish granodiorite gneiss as a local variant of medium grained granodiorite present. Pyroxene alteration obvious at 325' (99 m).	301.0	91.7	Vein-fracture (1 mm) with gypsum. 305/72SW.	
						322.0	98.1	Vein (4 mm) of greenish gypsum. 332/52SW.	
						325.0	99.1	Vein - pyroxene alteration band- 3 to 4 mm wide.. 048/72 SE.	
						327.0	99.7	Vein (3 mm) of gypsum. 272/84N.	

<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
329.0	353.0	100.3	107.6		Medium grained pyroxene granodiorite gneiss and white pyroxene-bearing aplite. Strong pyroxene alteration with disseminated fluorite and local biotite in fine grained white aplite. Granodiorite gneiss is local, in thin layers (10-15 cm)	349.0	106.4	Vein (2 cm) of fluorite, K-feldspar and biotite parallel to foliation.	55
353.0	371.0	107.6	113.1		Medium grained granodiorite gneiss. No aplite. Biotite+ K-feldspar+ quartz in strongly foliated gneiss.	353.0	107.6	Vein-fracture with gypsum. 056/78SE.	
371.0	435.0	113.1	132.6		Granodiorite gneiss with thin pink aplites gneiss zones (5-20 cm) with fine flashes of biotite+ muscovite+ pyrite outlining foliation. Granodiorite gneiss dominant.	363.0	110.6	Three slightly irregular veins (2-3 mm wide) of gypsum.058/51SE.	
						377.0	114.9	Vein (2 cm) of fluorite, K-feldspar and very minor pyrite. 015/57NW.	
						389.0	118.6	Vein (2 cm) of fluorite, biotite and calcite parallel to foliation.	
						391.0	119.2	Vein-fracture (2 mm) with gypsum. 288/62SW.	
						401.0	122.2	Vein (2-3 mm) of quartz and minor pyrite. 092/46S.	
						415.0	126.5	Vein (6-8 mm) of barren quartz. 088/33S.	
						415.0	126.5	Vein-fracture (2 mm) with gypsum. 082/88NW.	
						420.0	128.0	Vein (4 cm) of vuggy calcite, biotite, quartz and gypsum infilling. 070/43NW.	
						425.0	129.5	Vein-fracture (1-2 mm) with gypsum. 078/70SE.	
435.0	507.0	132.6	154.5		Granodiorite gneiss. Aplite absent. Strongly foliated S1 outlined by biotite + chlorite. Pyroxene alteration locally very intense (+K-feldspar and calcite). Thin K-feldspar layers (2 cm) white in colour are associated with pyroxene + biotite alteration.	439.0	133.8	Vein-fracture (2 mm) with gypsum. 073/90.	



<i>From (feet)</i>	<i>To (feet)</i>	<i>From (metre)</i>	<i>To (metre)</i>	<i>Unit</i>	<i>Lithology</i>	<i>Feet</i>	<i>Metre</i>	<i>Primary tectonic Structure</i>	<i>Angle</i>
						443.0	135.0	Vein (4-5 m) of quartz, calcite and chlorite. 072/68 NW.	
						450.0	137.2	Vein (1-1.5 cm) of fluorite, calcite and pyroxene parallel to foliation.	45
						460.0	140.2	Vein-fracture (1 mm) with gypsum. 060/86NW.	
					465' to 475' (141.7-14.8 m) strong gypsum alteration- irregular veins and fracture filling of gypsum. Local rounded grains of magnetite within alteration bands of chlorite+calcite+pyrite at 486' (148.1 m).	477.0	145.4	Vein-fracture (4 mm) with gypsum. 282/58SW.	
						482.0	146.9	Vein-fracture (2 mm) with gypsum. 012/83NW.	
						495.0	150.9	Vein (1 cm) of fluorite, calcite and chlorite. 305/45NE	
						502.0	153.0	Vein-fracture (1 mm) with gypsum. 052/78NW..	
	507.0		154.5		End of hole.				

Figure 1. Property Location Map



View of Eaglet from Quesnel Lake, 1980's (Eaglet Mines photo)



GEOLOGICAL SURVEY BRANCH  
 ASSESSMENT REPORT  
 27