Lumina Resources Corp

2005 GEOLOGICAL, GEOCHEMICAL, GEOPHYSICAL AND DIAMOND DRILLING REPORT ON THE HUSHAMU PROPERTY

Volume II – Figures

Located in the Northern Vancouver Island Area
Nanaimo Mining Division
NTS 092L/12
50° 39' North Latitude
127° 47' West Longitude

-prepared for-

LUMINA RESOURCES CORP
1550 - 625 Howe Street
Vancouver, BC
Canada V6C 2T6

-prepared by-

Darcy Baker
EQUITY ENGINEERING LTD.
Suite 700, 700 West Pender Street
Vancouver, British Columbia, Canada
V6C 1G8
darcyb@equityeng.bc.ca

April, 2006





SAMPLE	Au_ppm	Ag_ppm	Cu_ppm	Mo_ppm	Pb_ppm
275590	0.62	-1	4,720	90	20
275591	0.84	1	280	30	20
275592	0.69	1	4,160	70	-20
275593	0.81	1	4,610	20	20
275594	0.01	-1	150	-10	20
275595	0.01	-1	60	-10	20
275596	-0.01	-1	190	-10	20
275597	0.32	-1	1,110	30	-20
275598	0.16	-1	140	-10	20
275599	0.73	-1	450	40	-20
275600	0.01	-1	30	-10	-20
279938	0.18	-1	140	-10	-20
279939	0.83	1	9,340	60	-20
279940	0.02	-1	230	-10	-20
279941	0.01	-1	160	-10	-20
279942	0.42	-1	270	20	-20
279943	0.49	2	2,750	100	-20
345852	-0.005	-0.2	23	2	4
379272	0.02	-1	60	40	150
379273	0.04	1	450	680	50
379274	0.04	-1	50	790	40
379275	0.02	1	90	20	80
379276	0.01	1	30	120	90
379277	-0.01	-1	70	-10	20
379293	0.06	-1	20	40	40
379294	0.08	-1	60	190	-20
379295	0.04	-1	110	-10	40
379296	0.19	18	12,050	10	-20
379297	0.07	1	100	-10	20
380003	0.07	1	110	10	-20
380004	0.03	-1	30	40	-20
380069	0.05	1	230	190	110
380070	0.73	2	4,330	-10	-20
380072	0.01	-1	80	-10	-20
380075	0.04	-1	230	-10	30
380076	0.02	1	40	-10	-20
380077	0.01	-1	100	-10	-20
380078	0.02	-1	110	-10	-20
380079	0.01	-1	30	-10	-20
380080	0.51	-1	2,930	10	-20
380081	0.16	-1	30	-10	-20
380082	0.07	-1	30	-10	-20
380083	0.02	1	20	-10	60
380084	0.02	-1	40	-10	-20
380085	0.03	-1	10	10	-20
380086	0.11	1	340	20	-20
380087	0.8	1	4,460	20	-20
380111	0.01	-1	10	-10	-20
380151	0.04	-1	50	110	20
380152	0.01	-1	30	-10	-20
391955	0.01	-1	20	-10	20

Lithologic Units

STRATIFIED ROCKS

QUATERNARY
- QT Glacial Till

LOWER CRETACEOUS
Coal Harbour Group
- lKC Sandstone, siltstone, conglomerate

UPPER TRIASSIC to MIDDLE JURASSIC
Bonanza Group
"Bonanza Volcanics"
- JB Undivided
- JB1 Andesite
- JB2 Andesite lapilli/ash tuff
- JB3 Rhyolite; flow foliated
- JB4 Dacite
- JB5 Basalt
- JB6 Sedimentary rock
- JB7 Biotite hornfels

"Bonanza Volcanics" altered facies
- JBa1 Silica, locally hydrothermal breccia
- JBa2 Silica-pyrophyllite+/-kaolinite+/-dickite+/-diaspore+/-alunite+/-topaz
- JBa3 Silica-pyrophyllite-clay with abundant pyrite
- JBa4 Clay (silica largely absent)
- JBa5 Silica-chlorite-magnetite
- JBa6 Sericite-chlorite-clay+/-pyrite

Parson Bay Formation
- uTP Impure limestone, calcareous mudstone, siltstone, shale, sandstone, volcaniclastic rocks, basaltic lithic tuff, pillow lava

UPPER TRIASSIC
Vancouver Group
Quatsino Formation
- uTQ Bedded to massive micritic to bioclastic limestone

Karmutsen Formation
- uTKls Thin horizons of limestone near the top of the succession
- uTK Aphanitic to coarsely plagioclase-phyric subaerial basalt, commonly amygdaloidal, local pillows and hyaloclastite

INTRUSIVE ROCKS

TERTIARY
Intrusive Dykes, Sills and Stocks
- Tl Basalt

EARLY to MIDDLE JURASSIC
Island Plutonic Suite
- Jl Undivided: medium to coarse grained, equigranular to porphyritic granitoid rocks
- Jl1 Diorite
- Jl2 Quartz feldspar porphyry
- Jl3 Granodiorite
- Jl4 Monzonite

Minor Intrusions
- JBi Mafic intrusions inferred to be "Bonanza" age

Western Forest Products Road

Fault, thrust
Major fault, normal
Outcrop
Claim Boundary
Mag high
Mag Low
Mag Mixed
Drill Hole
Bedding 45
Foliation 45
Joint 45
Dyke 45
2005 Rock Sample (grab,float)
Vein 45
Fold Axis 45
Fault 45
Lineation 45
2005 Silt Sample

0 100 200 300 500
metres

N

LUMINA COPPER CORPORATION

HUSHAMU PROPERTY

NW Expo, Cougar, RedDog GEOLOGY and 2005 SAMPLE LOCATIONS

Date	Scale	Figure
APR 2006	1:5000	7a
U.T.M. Zone: UTM9 - NAD83	Mining District: NANAIMO	
N.T.S.: 92L	State/Province: BC	

EQUITY





SAMPLE	Au_ppm	Ag_ppm	Cu_ppm	Mo_ppm	Pb_ppm	Zn_ppm
279925	0.02	-1	110	-10	50	50
379285	0.01	-1	50	-10	-20	140
379286	0.05	-1	110	70	-20	20
379287	0.08	8	100	240	20	-20
379288	0.01	-1	20	-10	40	-20
379289	-0.01	1	80	-10	30	60
379975	0.01	-1	40	10	40	-20
379976	-0.01	-1	30	10	40	-20
379977	0.23	34	60	-10	740	540
379978	-0.01	-1	30	-10	-20	-20
379980	0.01	-1	120	10	80	360
379981	0.4	1	740	-10	-20	200
379982	0.03	-1	90	-10	-20	190
379983	0.04	-1	150	10	50	90
379984	0.04	-1	120	-10	-20	40
379985	0.05	-1	40	10	-20	70
379986	0.01	-1	90	-10	270	210
379987	-0.01	-1	40	-10	-20	60
379988	-0.01	-1	20	-10	-20	60

Lithologic Units

STRATIFIED ROCKS

QUATERNARY
QT Glacial Till

LOWER CRETACEOUS
Coal Harbour Group
lKC Sandstone, siltstone, conglomerate

UPPER TRIASSIC to MIDDLE JURASSIC
Bonanza Group
"Bonanza Volcanics"
JB Undivided
JB1 Andesite
JB2 Andesite lapilli/ash tuff
JB3 Rhyolite; flow foliated
JB4 Dacite
JB5 Basalt
JB6 Sedimentary rock
JB7 Biotite hornfels

"Bonanza Volcanics" altered facies
JBa1 Silica, locally hydrothermal breccia
JBa2 Silica-pyrophyllite+/-kaolinite+/-dickite+/-diaspore+/-alunite+/-topaz
JBa3 Silica-pyrophyllite-clay with abundant pyrite
JBa4 Clay (silica largely absent)
JBa5 Silica-chlorite-magnetite
JBa6 Sericite-chlorite-clay+/-pyrite

Parson Bay Formation
uTP Impure limestone, calcareous mudstone, siltstone, shale, sandstone, volcaniclastic rocks, basaltic lithic tuff, pillow lava

UPPER TRIASSIC
Vancouver Group
Quatsino Formation
uTQ Bedded to massive micritic to bioclastic limestone
Karmutsen Formation
uTKls Thin horizons of limestone near the top of the succession
uTK Aphanitic to coarsely plagioclase-phyric subaerial basalt, commonly amygdaloidal, local pillows and hyaloclastite

INTRUSIVE ROCKS

TERTIARY
Intrusive Dykes, Sills and Stocks
TI Basalt

EARLY to MIDDLE JURASSIC
Island Plutonic Suite
JI Undivided: medium to coarse grained, equigranular to porphyritic granitoid rocks
JI1 Diorite
JI2 Quartz feldspar porphyry
JI3 Granodiorite
JI4 Monzonite

Minor Intrusions
JBi Mafic intrusions inferred to be "Bonanza" age

Western Forest Products Road

Fault, thrust
Major fault, normal
Outcrop
Claim Boundary
Mag high
Mag Low
Mag Mixed
Drill Hole
45 Bedding
45 Foliation
45 Joint
45 Dyke
2005 Rock Sample (grab,float)
45 Vein
45 Fold Axis
45 Fault
45 Lineation
2005 Silt Sample

0 100 200 300 500
metres

N

LUMINA COPPER CORPORATION

HUSHAMU PROPERTY

SICKLE Area GEOLOGY AND 2005 SAMPLE LOCATIONS

EQUITY

Date	Scale	Figure
APR 2006	1:5000	7b
U.T.M. Zone: UTM9 - NAD83	Mining District: NANAIMO	
N.T.S.: 92L	State/Province: BC	





SAMPLE	Au_ppm	Ag_ppm	Cu_ppm	Mo_ppm	Pb_ppm	Zn_ppm
279919	0.03	-1	160	20	20	20
279920	0.01	-1	60	10	-20	20
279921	0.03	-1	40	20	-20	60
279922	0.01	-1	30	-10	-20	30
279923	0.01	-1	30	-10	-20	20
279924	0.01	-1	120	-10	20	40
379283	0.2	1	2,770	-10	20	90
379285	0.01	-1	50	-10	-20	140
379964	0.04	-1	490	10	-20	120
379965	0.1	-1	140	-10	70	130
379966	0.16	1	1,590	60	-20	30
379967	0.05	-1	690	10	-20	90
379968	0.1	-1	870	20	20	70
379969	0.12	-1	920	10	-20	80
379970	0.13	-1	1,080	10	-20	70
379971	0.08	-1	1,220	10	-20	80
379972	0.04	-1	790	10	-20	80
379973	0.05	-1	420	20	-20	60
379974	0.06	1	620	20	-20	60

Lithologic Units

STRATIFIED ROCKS

QUATERNARY
- QT Glacial Till

LOWER CRETACEOUS
Coal Harbour Group
- lKC Sandstone, siltstone, conglomerate

UPPER TRIASSIC to MIDDLE JURASSIC
Bonanza Group
"Bonanza Volcanics"
- JB Undivided
- JB1 Andesite
- JB2 Andesite lapilli/ash tuff
- JB3 Rhyolite; flow foliated
- JB4 Dacite
- JB5 Basalt
- JB6 Sedimentary rock
- JB7 Biotite hornfels

"Bonanza Volcanics" altered facies
- JBa1 Silica, locally hydrothermal breccia
- JBa2 Silica-pyrophyllite+/-kaolinite+/-dickite+/-diaspore+/-alunite+/-topaz
- JBa3 Silica-pyrophyllite-clay with abundant pyrite
- JBa4 Clay (silica largely absent)
- JBa5 Silica-chlorite-magnetite
- JBa6 Sericite-chlorite-clay+/-pyrite

Parson Bay Formation
- uTP Impure limestone, calcareous mudstone, siltstone, shale, sandstone, volcaniclastic rocks, basaltic lithic tuff, pillow lava

UPPER TRIASSIC
Vancouver Group
Quatsino Formation
- uTQ Bedded to massive micritic to bioclastic limestone

Karmutsen Formation
- uTKls Thin horizons of limestone near the top of the succession
- uTK Aphanitic to coarsely plagioclase-phyric subaerial basalt, commonly amygdaloidal, local pillows and hyaloclastite

INTRUSIVE ROCKS

TERTIARY
Intrusive Dykes, Sills and Stocks
- Tl Basalt

EARLY to MIDDLE JURASSIC
Island Plutonic Suite
- Jl Undivided: medium to coarse grained, equigranular to porphyritic granitoid rocks
- Jl1 Diorite
- Jl2 Quartz feldspar porphyry
- Jl3 Granodiorite
- Jl4 Monzonite

Minor Intrusions
- JBi Mafic intrusions inferred to be "Bonanza" age

Western Forest Products Road

Fault, thrust
Major fault, normal
Outcrop
Claim Boundary
Mag high
Mag Low
Mag Mixed
Drill Hole
45 Bedding
45 Foliation
45 Joint
45 Dyke
2005 Rock Sample (grab,float)
45 Vein
45 Fold Axis
45 Fault
45 Lineation
2005 Silt Sample

0 100 200 300 500
metres

N

LUMINA COPPER CORPORATION

HUSHAMU PROPERTY

Goodspeed_Hep West Area GEOLOGY AND 2005 SAMPLE LOCATIONS

EQUITY

Date	Scale	Figure
APR 2006	1:5000	7c
U.T.M. Zone: UTM9 - NAD83	Mining District: NANAIMO	
N.T.S.: 92L	State/Province: BC	



SAMPLE	Au_ppm	Ag_ppm	Cu_ppm	Mo_ppm	Pb_ppm	Zn_ppm
279926	-0.01	-1	210	-10	50	70
279927	-0.01	-1	330	-10	30	90
279928	-0.01	-1	20	-10	30	40
379290	-0.01	-1	10	-10	-20	30
379291	-0.01	-1	20	-10	20	70
379292	-0.01	-1	10	-10	-20	50
379298	0.01	-1	370	-10	-20	70
379299	0.03	1	1,190	4,400	-20	180
379300	0.01	-1	250	20	-20	40
379989	0.01	2	3,170	-10	-20	80
379990	-0.01	-1	50	-10	-20	50
379991	0.02	-1	70	70	-20	-20
379992	0.02	1	90	-10	40	-20
380059	0.11	-1	50	-10	-20	-20
380060	-0.01	-1	40	-10	-20	30
380061	-0.01	-1	30	-10	-20	40
380101	0.04	17	6,680	140	140	310
380102	-0.01	-1	190	-10	-20	60
380103	0.06	2	580	-10	-20	50

SAMPLE	Au_ppm	Ag_ppm	Cu_ppm	Mo_ppm	Pb_ppm	Zn_ppm
275583	-0.01	-1	190	-10	30	70
275584	-0.01	1	110	10	40	220
275585	-0.01	-1	50	-10	20	90
275586	0.05	3	250	10	250	660
275587	-0.01	-1	50	-10	30	120
275588	-0.01	-1	100	-10	60	300
275589	-0.01	-1	60	-10	30	120
279917	0.32	-1	3,040	30	-20	90
279918	0.13	-1	960	20	20	230
279929	-0.01	-1	10	10	60	150
279930	-0.01	2	10	-10	-20	30
279931	0.01	-1	60	-10	20	90
279932	0.01	-1	80	-10	-20	100
279933	0.01	-1	160	-10	30	150
279934	0.01	-1	100	-10	130	780
379278	0.01	2	510	70	110	640
379281	0.03	1	250	-10	80	100
379282	0.03	-1	40	10	30	20
379378	-0.01	-1	160	-10	-20	360
379379	0.11	18	25,800	30	40	190
379380	0.25	2	1,050	10	50	2,410
379381	0.01	-1	90	-10	20	1,510
379382	-0.01	1	120	-10	-20	140
379383	-0.01	-1	40	-10	-20	70
379384	0.01	4	500	20	350	200
379385	0.07	2	260	20	70	380
379386	0.03	-1	580	110	30	140
379387	-0.01	1	30	-10	-20	100
379388	-0.01	-1	60	-10	-20	90
379389	-0.01	-1	110	-10	-20	110
379390	-0.01	-1	40	-10	-20	80
379391	-0.01	3	3,200	20	30	110
379392	0.01	-1	110	-10	40	60
379393	0.01	-1	90	40	30	40
379394	0.02	-1	80	20	-20	160
379395	0.01	2	2,030	-10	20	680
379396	0.02	5	8,270	20	-20	140
379397	-0.01	1	50	-10	-20	160
379398	0.01	4	4,570	-10	30	290
379399	0.17	-1	150	-10	-20	60
379400	0.03	-1	230	30	-20	180
379958	0.21	1	1,640	30	20	410
379959	0.18	1	1,310	50	20	50
379960	0.08	1	1,690	20	20	320
379961	0.04	1	1,640	20	50	430
379962	0.03	-1	1,380	70	30	410
379963	0.04	1	1,640	70	20	270
379999	-0.01	-1	110	-10	-20	190
380000	-0.01	-1	140	-10	-20	670
380002	-0.01	-1	70	-10	-20	60
380051	-0.01	-1	50	-10	-20	120
380052	0.02	-1	900	10	-20	80
380053	0.03	-1	60	10	-20	70
380115	0.01	-1	10	-10	-20	-20

Lithologic Units

STRATIFIED ROCKS

QUATERNARY
- QT — Glacial Till

LOWER CRETACEOUS
Coal Harbour Group
- IKC — Sandstone, siltstone, conglomerate

UPPER TRIASSIC to MIDDLE JURASSIC
Bonanza Group
"Bonanza Volcanics"
- JB — Undivided
- JB1 — Andesite
- JB2 — Andesite lapilli/ash tuff
- JB3 — Rhyolite; flow foliated
- JB4 — Dacite
- JB5 — Basalt
- JB6 — Sedimentary rock
- JB7 — Biotite hornfels

"Bonanza Volcanics" altered facies
- JBa1 — Silica, locally hydrothermal breccia
- JBa2 — Silica-pyrophyllite+/-kaolinite+/-dickite+/-diaspore
- JBa3 — Silica-pyrophyllite-clay with abundant pyrite
- JBa4 — Clay (silica largely absent)
- JBa5 — Silica-chlorite-magnetite
- JBa6 — Sericite-chlorite-clay+/-pyrite

Parson Bay Formation
- uTP — Impure limestone, calcareous mudstone, siltstone, shale, sandstone, volcaniclastic rocks, basaltic lithic tuff, pillow lava

UPPER TRIASSIC
Vancouver Group
Quatsino Formation
- uTQ — Bedded to massive micritic to bioclastic limestone

Karmutsen Formation
- uTKls — Thin horizons of limestone near the top of the succession
- uTK — Aphanitic to coarsely plagioclase-phyric subaerial basalt, commonly amygdaloidal, local pillows and hyaloclastite

INTRUSIVE ROCKS

TERTIARY
Intrusive Dykes, Sills and Stocks
- TI — Basalt

EARLY to MIDDLE JURASSIC
Island Plutonic Suite
- JI — Undivided: medium to coarse grained, equigranular to porphyritic granitoid rocks
- JI1 — Diorite
- JI2 — Quartz feldspar porphyry
- JI3 — Granodiorite
- JI4 — Monzonite

Minor Intrusions
- JBi — Mafic intrusions inferred to be "Bonanza" age

Western Forest Products Road

Fault, thrust — Major fault, normal — Outcrop — Claim Boundary — Mag high — Mag Low — Mag Mixed

Bedding 45 — Foliation 45 — Joint 45 — Dyke 45 — Vein 45 — Fold Axis 45 — Fault 45 — Lineation 45

0 100 200 300 500
metres

LUMINA COPPER CORPORATION

HUSHAMU PROPERTY

Hushamu Deposit Area GEOLOGY AND 2005 SAMPLE LOCATIONS

Date	Scale	Figure
APR 2006	1:5000	7e
U.T.M. Zone: UTM9 - NAD83	Mining District: NANAIMO	
N.T.S.: 92L	State/Province: BC	

EQUITY

EC-215

400 mN

300 mN

200 mN

100 mN

0 mN

104.0 m (26.0 to 130.0 m)
0.30 g/t Au, 0.28% Cu
70 ppm Mo

HUSHAMU FAULT

CASN

JBa5

FLBX

FLTG

JI2

FLBX
FLBX

FLBX

JB1

EC-215
410.6m

Lithologic Units

STRATIFIED ROCKS

FLBX — Fault Breccia
FLTG — Fault Gouge

QUATERNARY
OVBN — Overburden
QT — Glacial Till

LOWER CRETACEOUS
Coal Harbour Group
IKC — Sandstone, siltstone, conglomerate

UPPER TRIASSIC to MIDDLE JURASSIC
Bonanza Group

"Bonanza Volcanics"		***"Bonanza Volcanics" altered facies***	
JB	Undivided		
JB1	Andesite	JBa1	Silica, locally hydrothermal breccia
JB2	Andesite lapilli/ash tuff	JBa2	Silica-pyrophyllite+/-kaolinite+/-dickite+/-diaspore+/-alunite+/-topaz
JB3	Rhyolite; flow foliated	JBa3	Silica-pyrophyllite-clay with abundant pyrite
JB4	Dacite	JBa4	Clay (silica largely absent)
JB5	Basalt	JBa5	Silica-chlorite-magnetite
JB6	Sedimentary rock	JBa6	Sericite-chlorite-clay+/-pyrite
JB7	Biotite hornfels		

Parson Bay Formation
uTP — Impure limestone, calcareous mudstone, siltstone, shale, sandstone, volcaniclastic rocks, basaltic lithic tuff, pillow lava

UPPER TRIASSIC
Vancouver Group
Quatsino Formation
uTQ — Bedded to massive micritic to bioclastic limestone
Karmutsen Formation
uTKls — Thin horizons of limestone near the top of the succession
uTK — Aphanitic to coarsely plagioclase-phyric subaerial basalt, commonly amygdaloidal, local pillows and hyaloclastite

INTRUSIVE ROCKS

TERTIARY
Intrusive Dykes, Sills and Stocks
TI — Basalt

EARLY to MIDDLE JURASSIC
Island Plutonic Suite
JI — Undivided: medium to coarse grained, equigranular to porphyritic granitoid rocks
JI1 — Diorite
JI2 — Quartz feldspar porphyry
JI3 — Granodiorite
JI4 — Monzonite
Minor Intrusions
JBi — Mafic intrusions inferred to be "Bonanza" age

HOLEID

GEOCHEMISTRY HISTOGRAM KEY

Cu_ppm Histogram 1.0cm = 2000 ppm

Au_ppm Histogram 1.0cm = 2 ppm

Mo_ppm Histogram 1.5cm = 100 ppm

EOH

50 metres

LUMINA RESOURCES CORPORATION

HUSHAMU PROPERTY

Cross Section XS EC-215

Looking 090°

EQUITY

Date	Scale	Figure
APR 2006	1:1000	8a
U.T.M. Zone: UTM9 - NAD83	Mining District: NANAIMO	
N.T.S.: 92L	State/Province: BC	

DACT coarse ash +/-lapilli tuff
w/ BI alt.

Chloritic fg ANDS or ANDS lapilli tuff

EC-137

EC-094

EC-078

EC-176

EC-216

EC-179

EC-203

600 mN

500 mN

400 mN

300 mN

200 mN

100 mN

0 mN

148.4 m (12.2 to 160.6 m)
0.51 g/t Au, 0.31% Cu
61 ppm Mo

EC-137
27.4m

EC-137A
136.2m

EC-094
178m

EC-078
152.4m

EC-216
394.1m

EC-176
256m

EC-179
495.8m

EC-180
465.1m

EC-203
495.6m

50 metres

Lithologic Units

STRATIFIED ROCKS

FLBX Fault Breccia
FLTG Fault Gouge

QUATERNARY
OVBN Overburden
QT Glacial Till

LOWER CRETACEOUS
Coal Harbour Group
IKC Sandstone, siltstone, conglomerate

UPPER TRIASSIC to MIDDLE JURASSIC
Bonanza Group

"Bonanza Volcanics"		***"Bonanza Volcanics" altered facies***	
JB	Undivided		
JB1	Andesite	JBa1	Silica, locally hydrothermal breccia
JB2	Andesite lapilli/ash tuff	JBa2	Silica-pyrophyllite+/-kaolinite+/-dickite+/-diaspore+/-alunite+/-topaz
JB3	Rhyolite; flow foliated	JBa3	Silica-pyrophyllite-clay with abundant pyrite
JB4	Dacite	JBa4	Clay (silica largely absent)
JB5	Basalt	JBa5	Silica-chlorite-magnetite
JB6	Sedimentary rock	JBa6	Sericite-chlorite-clay+/-pyrite
JB7	Biotite hornfels		

Paradise Bay Formation
uTP Impure limestone, calcareous mudstone, siltstone, shale, sandstone, volcaniclastic rocks, basaltic lithic tuff, pillow lava

UPPER TRIASSIC
Vancouver Group
Quatsino Formation
uTQ Bedded to massive micritic to bioclastic limestone
Karmutsen Formation
uTKb Thin horizons of limestone near the top of the succession
uTK Aphanitic to coarsely plagioclase-phyric subaerial basalt, commonly amygdaloidal, local pillows and hyaloclastite

INTRUSIVE ROCKS

TERTIARY
Intrusive Dykes, Sills and Stocks
TI Basalt

EARLY to MIDDLE JURASSIC
Island Plutonic Suite
JI Undivided: medium to coarse grained, equigranular to porphyritic granitoid rocks
JI1 Diorite
JI2 Quartz feldspar porphyry
JI3 Granodiorite
JI4 Monzonite

Minor Intrusions
JBi Mafic intrusions inferred to be "Bonanza" age

HOLEID
Cu_ppm Histogram 1.0cm = 2000 ppm
Au_ppm Histogram 1.0cm = 2 ppm
Mo_ppm Histogram 1.5cm = 100 ppm
EOH

GEOCHEMISTRY HISTOGRAM KEY

LUMINA RESOURCES CORPORATION

HUSHAMU PROPERTY

Cross Section XS EC-216

Looking 090°

Date: APR 2006		Scale: 1:1000	Figure: 8b
U.T.M. Zone: UTM9 - NAD83		Mining District: NANAIMO	
N.T.S.: 92L		State/Province: BC	



Lithologic Units

STRATIFIED ROCKS

FLBX — Fault Breccia
FLTG — Fault Gouge

QUATERNARY
OVBN — Overburden
QT — Glacial Till

LOWER CRETACEOUS
Coal Harbour Group
IKC — Sandstone, siltstone, conglomerate

UPPER TRIASSIC to MIDDLE JURASSIC
Bonanza Group

"Bonanza Volcanics"		***"Bonanza Volcanics" altered facies***	
JB — Undivided			
JB1 — Andesite		JBa1	Silica, locally hydrothermal breccia
JB2 — Andesite lapilli/ash tuff		JBa2	Silica-pyrophyllite+/-kaolinite+/-dickite+/-diaspore+/-alunite+/-topaz
JB3 — Rhyolite; flow foliated		JBa3	Silica-pyrophyllite-clay with abundant pyrite
JB4 — Dacite		JBa4	Clay (silica largely absent)
JB5 — Basalt		JBa5	Silica-chlorite-magnetite
JB6 — Sedimentary rock		JBa6	Sericite-chlorite-clay+/-pyrite
JB7 — Biotite hornfels			

Parson Bay Formation
uTP — Impure limestone, calcareous mudstone, siltstone, shale, sandstone, volcaniclastic rocks, basaltic lithic tuff, pillow lava

UPPER TRIASSIC
Vancouver Group
Quatsino Formation
uTQ — Bedded to massive micritic to bioclastic limestone
Karmutsen Formation
uTKls — Thin horizons of limestone near the top of the succession
uTK — Aphanitic to coarsely plagioclase-phyric subaerial basalt, commonly amygdaloidal, local pillows and hyaloclastite

INTRUSIVE ROCKS

TERTIARY
Intrusive Dykes, Sills and Stocks
TI — Basalt

EARLY to MIDDLE JURASSIC
Island Plutonic Suite
JI — Undivided: medium to coarse grained, equigranular to porphyritic granitoid rocks
JI1 — Diorite
JI2 — Quartz feldspar porphyry
JI3 — Granodiorite
JI4 — Monzonite
Minor Intrusions
JBi — Mafic intrusions inferred to be "Bonanza" age

HOLEID

GEOCHEMISTRY HISTOGRAM KEY

Cu_ppm Histogram 1.0cm = 2000 ppm
Au_ppm Histogram 1.0cm = 2 ppm
Mo_ppm Histogram 1.5cm = 100 ppm

EOH

50 metres

LUMINA RESOURCES CORPORATION

HUSHAMU PROPERTY

Cross Section XS EC-217

Looking 090°

EQUITY

Date: APR 2006	Scale: 1:1000	Figure: **8c**
U.T.M. Zone: UTM9 - NAD83	Mining District: NANAIMO	
N.T.S.: 92L	State/Province: BC	

DACT
lap-tuff
mod. welded
mCL

EC-228

EC-218

N

400 mN

400 mN

Airborne Magnetic High Surface Trace

95.0 m (132.9 to 227.9 m)
1.0 g/t Au, 0.17% Cu
250 ppm Mo

124.9 m (317.8 to 442.7 m)
0.52 g/t Au, 0.11% Cu
46 ppm Mo

300 mN

300 mN

200 mN

200 mN

100 mN

EC-228
235.6m

EC-218
462.4m

Lithologic Units

STRATIFIED ROCKS

FLBX Fault Breccia
FLTG Fault Gouge

QUATERNARY
OVBN Overburden
QT Glacial Till

LOWER CRETACEOUS
Coal Harbour Group
lKC Sandstone, siltstone, conglomerate

UPPER TRIASSIC to MIDDLE JURASSIC
Bonanza Group

"Bonanza Volcanics"		***"Bonanza Volcanics" altered facies***	
JB	Undivided		
JB1	Andesite	JBa1	Silica, locally hydrothermal breccia
JB2	Andesite lapilli/ash tuff	JBa2	Silica-pyrophyllite+/-kaolinite+/-dickite+/-diaspore+/-alunite+/-topaz
JB3	Rhyolite; flow foliated	JBa3	Silica-pyrophyllite-clay with abundant pyrite
JB4	Dacite	JBa4	Clay (silica largely absent)
JB5	Basalt	JBa5	Silica-chlorite-magnetite
JB6	Sedimentary rock	JBa6	Sericite-chlorite-clay+/-pyrite
JB7	Biotite hornfels		

Parson Bay Formation
uTP Impure limestone, calcareous mudstone, siltstone, shale, sandstone, volcaniclastic rocks, basaltic lithic tuff, pillow lava

UPPER TRIASSIC
Vancouver Group
Quatsino Formation
uTQ Bedded to massive micritic to bioclastic limestone
Karmutsen Formation
uTKls Thin horizons of limestone near the top of the succession
uTK Aphanitic to coarsely plagioclase-phyric subaerial basalt, commonly amygdaloidal, local pillows and hyaloclastite

INTRUSIVE ROCKS

TERTIARY
Intrusive Dykes, Sills and Stocks
TI Basalt

EARLY to MIDDLE JURASSIC
Island Plutonic Suite
JI Undivided: medium to coarse grained, equigranular to porphyritic granitoid rocks
JI1 Diorite
JI2 Quartz feldspar porphyry
JI3 Granodiorite
JI4 Monzonite

Minor Intrusions
JBi Mafic intrusions inferred to be "Bonanza" age

HOLEID

GEOCHEMISTRY HISTOGRAM KEY

Cu_ppm Histogram 1.0cm = 2000 ppm

Au_ppm Histogram 1.0cm = 2 ppm

Mo_ppm Histogram 1.5cm = 100 ppm

EOH

50 metres

LUMINA RESOURCES CORPORATION

HUSHAMU PROPERTY

Cross Section
XS EC-218, EC-228

Looking 295°

EQUITY

Date	Scale	Figure
APR 2006	1:1000	8d
U.T.M. Zone: UTM9 - NAD83	Mining District: NANAIMO	
N.T.S.: 92L	State/Province: BC	



Lithologic Units

STRATIFIED ROCKS

FLBX — Fault Breccia
FLTG — Fault Gouge

QUATERNARY
OVBN — Overburden
QT — Glacial Till

LOWER CRETACEOUS
Coal Harbour Group
IKC — Sandstone, siltstone, conglomerate

UPPER TRIASSIC to MIDDLE JURASSIC
Bonanza Group

"Bonanza Volcanics"		***"Bonanza Volcanics" altered facies***	
JB	Undivided		
JB1	Andesite	JBa1	Silica, locally hydrothermal breccia
JB2	Andesite lapilli/ash tuff	JBa2	Silica-pyrophyllite+/-kaolinite+/-dickite+/-diaspore+/-alunite+/-topaz
JB3	Rhyolite; flow foliated	JBa3	Silica-pyrophyllite-clay with abundant pyrite
JB4	Dacite	JBa4	Clay (silica largely absent)
JB5	Basalt	JBa5	Silica-chlorite-magnetite
JB6	Sedimentary rock	JBa6	Sericite-chlorite-clay+/-pyrite
JB7	Biotite hornfels		

Parson Bay Formation
uTP — Impure limestone, calcareous mudstone, siltstone, shale, sandstone, volcaniclastic rocks, basaltic lithic tuff, pillow lava

UPPER TRIASSIC
Vancouver Group
Quatsino Formation
uTQ — Bedded to massive micritic to bioclastic limestone
Karmutsen Formation
uTKls — Thin horizons of limestone near the top of the succession
uTK — Aphanitic to coarsely plagioclase-phyric subaerial basalt, commonly amygdaloidal, local pillows and hyaloclastite

INTRUSIVE ROCKS

TERTIARY
Intrusive Dykes, Sills and Stocks
TI — Basalt

EARLY to MIDDLE JURASSIC
Island Plutonic Suite
JI — Undivided: medium to coarse grained, equigranular to porphyritic granitoid rocks
JI1 — Diorite
JI2 — Quartz feldspar porphyry
JI3 — Granodiorite
JI4 — Monzonite

Minor Intrusions
JBi — Mafic intrusions inferred to be "Bonanza" age

HOLEID
GEOCHEMISTRY HISTOGRAM KEY
Cu_ppm Histogram 1.0cm = 2000 ppm
Au_ppm Histogram 1.0cm = 2 ppm
Mo_ppm Histogram 1.5cm = 100 ppm
EOH

50 metres

LUMINA RESOURCES CORPORATION

HUSHAMU PROPERTY

Cross Section XS EC-219, EC-220

Looking 330°

EQUITY

Date	Scale	Figure
APR 2006	1:1000	**8e**
U.T.M. Zone: UTM9 - NAD83	Mining District: NANAIMO	
N.T.S.: 92L	State/Province: BC	



Lithologic Units

STRATIFIED ROCKS

FLBX — Fault Breccia
FLTG — Fault Gouge

QUATERNARY
OVBN — Overburden
QT — Glacial Till

LOWER CRETACEOUS
Coal Harbour Group
IKC — Sandstone, siltstone, conglomerate

UPPER TRIASSIC to MIDDLE JURASSIC
Bonanza Group
"Bonanza Volcanics"
JB — Undivided
JB1 — Andesite
JB2 — Andesite lapilli/ash tuff
JB3 — Rhyolite; flow foliated
JB4 — Dacite
JB5 — Basalt
JB6 — Sedimentary rock
JB7 — Biotite hornfels

"Bonanza Volcanics" altered facies
JBa1 — Silica, locally hydrothermal breccia
JBa2 — Silica-pyrophyllite+/-kaolinite+/-dickite+/-diaspore+/-alunite+/-topaz
JBa3 — Silica-pyrophyllite-clay with abundant pyrite
JBa4 — Clay (silica largely absent)
JBa5 — Silica-chlorite-magnetite
JBa6 — Sericite-chlorite-clay+/-pyrite

Parson Bay Formation
uTP — Impure limestone, calcareous mudstone, siltstone, shale, sandstone, volcaniclastic rocks, basaltic lithic tuff, pillow lava

UPPER TRIASSIC
Vancouver Group
Quatsino Formation
uTQ — Bedded to massive micritic to bioclastic limestone
Karmutsen Formation
uTKls — Thin horizons of limestone near the top of the succession
uTK — Aphanitic to coarsely plagioclase-phyric subaerial basalt, commonly amygdaloidal, local pillows and hyaloclastite

INTRUSIVE ROCKS

TERTIARY
Intrusive Dykes, Sills and Stocks
TI — Basalt

EARLY to MIDDLE JURASSIC
Island Plutonic Suite
JI — Undivided: medium to coarse grained, equigranular to porphyritic granitoid rocks
JI1 — Diorite
JI2 — Quartz feldspar porphyry
JI3 — Granodiorite
JI4 — Monzonite

Minor Intrusions
JBi — Mafic intrusions inferred to be "Bonanza" age

HOLEID
GEOCHEMISTRY HISTOGRAM KEY
Cu_ppm Histogram 1.0cm = 2000 ppm
Au_ppm Histogram 1.0cm = 2 ppm
Mo_ppm Histogram 1.5cm = 100 ppm
EOH

LUMINA RESOURCES CORPORATION

HUSHAMU PROPERTY

Cross Section XS EC-221

Looking 317°

EQUITY

Date: APR 2006	Scale: 1:1000	Figure: 8f
U.T.M. Zone: UTM9 - NAD83	Mining District: NANAIMO	
N.T.S.: 92L	State/Province: BC	



Lithologic Units

STRATIFIED ROCKS

FLBX — Fault Breccia
FLTG — Fault Gouge

QUATERNARY
OVBN — Overburden
QT — Glacial Till

LOWER CRETACEOUS
Coal Harbour Group
IKC — Sandstone, siltstone, conglomerate

UPPER TRIASSIC to MIDDLE JURASSIC
Bonanza Group
"Bonanza Volcanics"

JB — Undivided
JB1 — Andesite
JB2 — Andesite lapilli/ash tuff
JB3 — Rhyolite; flow foliated
JB4 — Dacite
JB5 — Basalt
JB6 — Sedimentary rock
JB7 — Biotite hornfels

"Bonanza Volcanics" altered facies

JBa1 — Silica, locally hydrothermal breccia
JBa2 — Silica-pyrophyllite+/-kaolinite+/-dickite+/-diaspore+/-alunite+/-topaz
JBa3 — Silica-pyrophyllite-clay with abundant pyrite
JBa4 — Clay (silica largely absent)
JBa5 — Silica-chlorite-magnetite
JBa6 — Sericite-chlorite-clay+/-pyrite

Parson Bay Formation
uTP — Impure limestone, calcareous mudstone, siltstone, shale, sandstone, volcaniclastic rocks, basaltic lithic tuff, pillow lava

UPPER TRIASSIC
Vancouver Group
Quatsino Formation
uTQ — Bedded to massive micritic to bioclastic limestone
Karmutsen Formation
uTKls — Thin horizons of limestone near the top of the succession
uTK — Aphanitic to coarsely plagioclase-phyric subaerial basalt, commonly amygdaloidal, local pillows and hyaloclastite

INTRUSIVE ROCKS

TERTIARY
Intrusive Dykes, Sills and Stocks
TI — Basalt

EARLY to MIDDLE JURASSIC
Island Plutonic Suite
JI — Undivided: medium to coarse grained, equigranular to porphyritic granitoid rocks
JI1 — Diorite
JI2 — Quartz feldspar porphyry
JI3 — Granodiorite
JI4 — Monzonite

Minor Intrusions
JBi — Mafic intrusions inferred to be "Bonanza" age

HOLEID
GEOCHEMISTRY HISTOGRAM KEY
Cu_ppm Histogram 1.0cm = 2000 ppm
Au_ppm Histogram 1.0cm = 2 ppm
Mo_ppm Histogram 1.5cm = 100 ppm
EOH

50 metres

LUMINA RESOURCES CORPORATION

HUSHAMU PROPERTY

Cross Section
XS EC-222

Looking 085°

EQUITY

Date: APR 2006	Scale: 1:1000	Figure: **8g**
U.T.M. Zone: UTM9 - NAD83	Mining District: NANAIMO	
N.T.S.: 92L	State/Province: BC	



Lithologic Units

STRATIFIED ROCKS

FLBX — Fault Breccia
FLTG — Fault Gouge

QUATERNARY
OVBN — Overburden
QT — Glacial Till

LOWER CRETACEOUS
Coal Harbour Group
IKC — Sandstone, siltstone, conglomerate

UPPER TRIASSIC to MIDDLE JURASSIC
Bonanza Group
"Bonanza Volcanics"
JB — Undivided
JB1 — Andesite
JB2 — Andesite lapilli/ash tuff
JB3 — Rhyolite; flow foliated
JB4 — Dacite
JB5 — Basalt
JB6 — Sedimentary rock
JB7 — Biotite hornfels

"Bonanza Volcanics" altered facies
JBa1 — Silica, locally hydrothermal breccia
JBa2 — Silica-pyrophyllite+/-kaolinite+/-dickite+/-diaspore+/-alunite+/-topaz
JBa3 — Silica-pyrophyllite-clay with abundant pyrite
JBa4 — Clay (silica largely absent)
JBa5 — Silica-chlorite-magnetite
JBa6 — Sericite-chlorite-clay+/-pyrite

Parson Bay Formation
uTP — Impure limestone, calcareous mudstone, siltstone, shale, sandstone, volcaniclastic rocks, basaltic lithic tuff, pillow lava

UPPER TRIASSIC
Vancouver Group
Quatsino Formation
uTQ — Bedded to massive micritic to bioclastic limestone
Karmutsen Formation
uTKls — Thin horizons of limestone near the top of the succession
uTK — Aphanitic to coarsely plagioclase-phyric subaerial basalt, commonly amygdaloidal, local pillows and hyaloclastite

INTRUSIVE ROCKS

TERTIARY
Intrusive Dykes, Sills and Stocks
TI — Basalt

EARLY to MIDDLE JURASSIC
Island Plutonic Suite
JI — Undivided: medium to coarse grained, equigranular to porphyritic granitoid rocks
JI1 — Diorite
JI2 — Quartz feldspar porphyry
JI3 — Granodiorite
JI4 — Monzonite

Minor Intrusions
JBi — Mafic intrusions inferred to be "Bonanza" age

HOLEID

GEOCHEMISTRY
HISTOGRAM
KEY

Cu_ppm
Histogram
1.0cm = 2000 ppm

Au_ppm
Histogram
1.0cm = 2 ppm

Mo_ppm
Histogram
1.5cm = 100 ppm

EOH

50 metres

LUMINA RESOURCES CORPORATION

HUSHAMU PROPERTY

Cross Section
XS EC-223

Looking 310°

EQUITY

Date: APR 2006	Scale: 1:1000	Figure: **8h**
U.T.M. Zone: UTM9 - NAD83	Mining District: NANAIMO	
N.T.S.: 92L	State/Province: BC	



Lithologic Units

STRATIFIED ROCKS

FLBX — Fault Breccia
FLTG — Fault Gouge

QUATERNARY
OVBN — Overburden
QT — Glacial Till

LOWER CRETACEOUS
Coal Harbour Group
IKC — Sandstone, siltstone, conglomerate

UPPER TRIASSIC to MIDDLE JURASSIC
Bonanza Group
"Bonanza Volcanics"

JB	Undivided
JB1	Andesite
JB2	Andesite lapilli/ash tuff
JB3	Rhyolite; flow foliated
JB4	Dacite
JB5	Basalt
JB6	Sedimentary rock
JB7	Biotite hornfels

"Bonanza Volcanics" altered facies

JBa1	Silica, locally hydrothermal breccia
JBa2	Silica-pyrophyllite+/-kaolinite+/-dickite+/-diaspore+/-alunite+/-topaz
JBa3	Silica-pyrophyllite-clay with abundant pyrite
JBa4	Clay (silica largely absent)
JBa5	Silica-chlorite-magnetite
JBa6	Sericite-chlorite-clay+/-pyrite

Parson Bay Formation
uTP — Impure limestone, calcareous mudstone, siltstone, shale, sandstone, volcaniclastic rocks, basaltic lithic tuff, pillow lava

UPPER TRIASSIC
Vancouver Group
Quatsino Formation
uTQ — Bedded to massive micritic to bioclastic limestone
Karmutsen Formation
uTKls — Thin horizons of limestone near the top of the succession
uTK — Aphanitic to coarsely plagioclase-phyric subaerial basalt, commonly amygdaloidal, local pillows and hyaloclastite

INTRUSIVE ROCKS

TERTIARY
Intrusive Dykes, Sills and Stocks
TI — Basalt

EARLY to MIDDLE JURASSIC
Island Plutonic Suite
JI — Undivided: medium to coarse grained, equigranular to porphyritic granitoid rocks
JI1 — Diorite
JI2 — Quartz feldspar porphyry
JI3 — Granodiorite
JI4 — Monzonite
Minor Intrusions
JBi — Mafic intrusions inferred to be "Bonanza" age

HOLEID
GEOCHEMISTRY HISTOGRAM KEY
Cu_ppm Histogram 1.0cm = 2000 ppm
Au_ppm Histogram 1.0cm = 2 ppm
Mo_ppm Histogram 1.5cm = 100 ppm
EOH

LUMINA RESOURCES CORPORATION

HUSHAMU PROPERTY

Cross Section
XS EC-224, EC-226
Looking 308°

EQUITY

Date: APR 2006	Scale: 1:1000	Figure: 8i
U.T.M. Zone: UTM9 - NAD83	Mining District: NANAIMO	
N.T.S.: 92L	State/Province: BC	



Lithologic Units

STRATIFIED ROCKS

FLBX — Fault Breccia
FLTG — Fault Gouge

QUATERNARY
OVBN — Overburden
QT — Glacial Till

LOWER CRETACEOUS
Coal Harbour Group
lKC — Sandstone, siltstone, conglomerate

UPPER TRIASSIC to MIDDLE JURASSIC
Bonanza Group
"Bonanza Volcanics" — ***"Bonanza Volcanics" altered facies***

JB	Undivided		
JB1	Andesite	JBa1	Silica, locally hydrothermal breccia
JB2	Andesite lapilli/ash tuff	JBa2	Silica-pyrophyllite+/-kaolinite+/-dickite+/-diaspore+/-alunite+/-topaz
JB3	Rhyolite; flow foliated	JBa3	Silica-pyrophyllite-clay with abundant pyrite
JB4	Dacite	JBa4	Clay (silica largely absent)
JB5	Basalt	JBa5	Silica-chlorite-magnetite
JB6	Sedimentary rock	JBa6	Sericite-chlorite-clay+/-pyrite
JB7	Biotite hornfels		

Parson Bay Formation
uTP — Impure limestone, calcareous mudstone, siltstone, shale, sandstone, volcaniclastic rocks, basaltic lithic tuff, pillow lava

UPPER TRIASSIC
Vancouver Group
Quatsino Formation
uTQ — Bedded to massive micritic to bioclastic limestone
Karmutsen Formation
uTKls — Thin horizons of limestone near the top of the succession
uTK — Aphanitic to coarsely plagioclase-phyric subaerial basalt, commonly amygdaloidal, local pillows and hyaloclastite

INTRUSIVE ROCKS

TERTIARY
Intrusive Dykes, Sills and Stocks
TI — Basalt

EARLY to MIDDLE JURASSIC
Island Plutonic Suite
JI — Undivided: medium to coarse grained, equigranular to porphyritic granitoid rocks
JI1 — Diorite
JI2 — Quartz feldspar porphyry
JI3 — Granodiorite
JI4 — Monzonite

Minor Intrusions
JBi — Mafic intrusions inferred to be "Bonanza" age

50 metres

LUMINA RESOURCES CORPORATION

HUSHAMU PROPERTY

Cross Section
XS EC-229

Looking 360°

EQUITY

Date: APR 2006	Scale: 1:1000	Figure: **8j**
U.T.M. Zone: UTM9 - NAD83	Mining District: NANAIMO	
N.T.S.: 92L	State/Province: BC	



Lithologic Units

STRATIFIED ROCKS

FLBX Fault Breccia
FLTG Fault Gouge

QUATERNARY
OVBN Overburden
QT Glacial Till

LOWER CRETACEOUS
Coal Harbour Group
IKC Sandstone, siltstone, conglomerate

UPPER TRIASSIC to MIDDLE JURASSIC
Bonanza Group
"Bonanza Volcanics"
JB Undivided
JB1 Andesite
JB2 Andesite lapilli/ash tuff
JB3 Rhyolite; flow foliated
JB4 Dacite
JB5 Basalt
JB6 Sedimentary rock
JB7 Biotite hornfels

"Bonanza Volcanics" altered facies
JBa1 Silica, locally hydrothermal breccia
JBa2 Silica-pyrophyllite+/-kaolinite+/-dickite+/-diaspore+/-alunite+/-topaz
JBa3 Silica-pyrophyllite-clay with abundant pyrite
JBa4 Clay (silica largely absent)
JBa5 Silica-chlorite-magnetite
JBa6 Sericite-chlorite-clay+/-pyrite

Parson Bay Formation
uTP Impure limestone, calcareous mudstone, siltstone, shale, sandstone, volcaniclastic rocks, basaltic lithic tuff, pillow lava

UPPER TRIASSIC
Vancouver Group
Quatsino Formation
uTQ Bedded to massive micritic to bioclastic limestone
Karmutsen Formation
uTKls Thin horizons of limestone near the top of the succession
uTK Aphanitic to coarsely plagioclase-phyric subaerial basalt, commonly amygdaloidal, local pillows and hyaloclastite

INTRUSIVE ROCKS

TERTIARY
Intrusive Dykes, Sills and Stocks
TI Basalt

EARLY to MIDDLE JURASSIC
Island Plutonic Suite
JI Undivided: medium to coarse grained, equigranular to porphyritic granitoid rocks
JI1 Diorite
JI2 Quartz feldspar porphyry
JI3 Granodiorite
JI4 Monzonite

Minor Intrusions
JBi Mafic intrusions inferred to be "Bonanza" age

HOLEID

GEOCHEMISTRY HISTOGRAM KEY

Cu_ppm Histogram 1.0cm = 2000 ppm
Au_ppm Histogram 1.0cm = 2 ppm
Mo_ppm Histogram 1.5cm = 100 ppm

EOH

50 metres

LUMINA RESOURCES CORPORATION

HUSHAMU PROPERTY

Cross Section
XS EC-230, EC-231

Looking 291°

EQUITY

Date: APR 2006	Scale: 1:1000	Figure: **8k**
U.T.M. Zone: UTM9 - NAD83	Mining District: NANAIMO	
N.T.S.: 92L	State/Province: BC	



300 mN

200 mN

100 mN

0 mN



Lithologic Units

STRATIFIED ROCKS

FLBX — Fault Breccia
FLTG — Fault Gouge

QUATERNARY
- OVBN — Overburden
- QT — Glacial Till

LOWER CRETACEOUS

Coal Harbour Group
- IKC — Sandstone, siltstone, conglomerate

UPPER TRIASSIC to MIDDLE JURASSIC

Bonanza Group

"Bonanza Volcanics"
- JB — Undivided
- JB1 — Andesite
- JB2 — Andesite lapilli/ash tuff
- JB3 — Rhyolite; flow foliated
- JB4 — Dacite
- JB5 — Basalt
- JB6 — Sedimentary rock
- JB7 — Biotite hornfels

"Bonanza Volcanics" altered facies
- JBa1 — Silica, locally hydrothermal breccia
- JBa2 — Silica-pyrophyllite+/-kaolinite+/-dickite+/-diaspore+/-alunite+/-topaz
- JBa3 — Silica-pyrophyllite-clay with abundant pyrite
- JBa4 — Clay (silica largely absent)
- JBa5 — Silica-chlorite-magnetite
- JBa6 — Sericite-chlorite-clay+/-pyrite

Parson Bay Formation
- uTP — Impure limestone, calcareous mudstone, siltstone, shale, sandstone, volcaniclastic rocks, basaltic lithic tuff, pillow lava

UPPER TRIASSIC

Vancouver Group

Quatsino Formation
- uTQ — Bedded to massive micritic to bioclastic limestone

Karmutsen Formation
- uTKls — Thin horizons of limestone near the top of the succession
- uTK — Aphanitic to coarsely plagioclase-phyric subaerial basalt, commonly amygdaloidal, local pillows and hyaloclastite

INTRUSIVE ROCKS

TERTIARY

Intrusive Dykes, Sills and Stocks
- TI — Basalt

EARLY to MIDDLE JURASSIC

Island Plutonic Suite
- JI — Undivided: medium to coarse grained, equigranular to porphyritic granitoid rocks
- JI1 — Diorite
- JI2 — Quartz feldspar porphyry
- JI3 — Granodiorite
- JI4 — Monzonite

Minor Intrusions
- JBi — Mafic intrusions inferred to be "Bonanza" age

HOLEID

GEOCHEMISTRY
HISTOGRAM
KEY

Cu_ppm
Histogram
1.0cm = 2000 ppm

Au_ppm
Histogram
1.0cm = 2 ppm

Mo_ppm
Histogram
1.5cm = 100 ppm

EOH

50 metres

LUMINA RESOURCES CORPORATION

HUSHAMU PROPERTY

Cross Section
XS EC-232

Looking 360°

EQUITY

Date: APR 2006	Scale: 1:1000	Figure: 8I
U.T.M. Zone: UTM9 - NAD83	Mining District: NANAIMO	
N.T.S.: 92L	State/Province: BC	

Soil Sample Key

- 1969 sampled for Mo and Zn
- 1969 & 72 sampled for Cu
- 1981 RedDog sampled for Au,As,Cu,Mo
- 1982 Utah Mines sampled for Au,As,Hg,Cu,Mo
- 1988/89 Expo sampled for Au,As,Bi,Se,Te
- 1989 Area B sampled for As,Se
- 1989 RedDog South sampled for Au,As,Cu
- 2005 sampled for 32 element ICP



Nahwitti Lake

Kains Lake

HOLBERG

Holberg Inlet

Soil Geochemistry Anomalies

- Copper Gold Moly
- Zinc Silver Lead

Hushamu Soil Geochemistry
Gold ppb

- > 87.2 (98th percentile)
- 45 to 87.2 (95th percentile)
- 27 to 45 (90th percentile)
- 16 to 27 (80th percentile)
- 7 to 16 (50th percentile)
- < 7 ppm or no data

Sample Key

- Soil Sample
- Rock Sample (float)
- Rock Sample > .5g/t Au

LUMINA COPPER CORPORATION

HUSHAMU PROPERTY

Rock and Soil Geochemistry (Gold ppb)

Date	APR 2006	Scale	1:25,000
U.T.M. Zone	UTM9 - NAD83	Mining District	NANAIMO
N.T.S.	92L	State/Province	BC

Figure 9a

Soil Sample Key

- 1969 sampled for Mo and Zn
- 1969 & 72 sampled for Cu
- 1981 RedDog sampled for Au,As,Cu,Mo
- 1982 Utah Mines sampled for Au,As,Hg,Cu,Mo
- 1988/89 Expo sampled for Au,As,Bi,Se,Te
- 1989 Area B sampled for As,Se
- 1989 RedDog South sampled for Au,As,Cu
- 2005 sampled for 32 element ICP



566000 mE
570000 mE
572000 mE
574000 mE
576000 mE
578000 mE
580000 mE
582000 mE
584000 mE
586000 mE
588000 mE
590000 mE

5622000 mN
5620000 mN
5618000 mN
5616000 mN
5614000 mN
5612000 mN
5610000 mN
5608000 mN
5606000 mN

HOLBERG

H o l b e r g I n l e t

Soil Geochemistry Anomalies

- Copper Gold Moly
- Zinc Silver Lead

Hushamu Soil Geochemistry
Copper ppm

- > 225 (98th percentile)
- 126 to 225 (95th percentile)
- 90 to 126 (90th percentile)
- 63 to 90 (80th percentile)
- 33 to 63 (50th percentile)
- < 33 ppm or no data

Sample Key

- Soil Sample
- Rock Sample (float)
- Rock Sample > 3000ppm Cu

LUMINA COPPER CORPORATION

HUSHAMU PROPERTY

Rock and Soil Geochemistry (Copper ppm)

Date	APR 2006	Scale	1:25,000
UTM Zone	UTM9 - NAD83	Mining District	NANAIMO
N.T.S.	92L	State/Province	BC

Figure 9b

Soil Sample Key

- 1969 sampled for Mo and Zn
- 1969 & 72 sampled for Cu
- 1981 RedDog sampled for Au,As,Cu,Mo
- 1982 Utah Mines sampled for Au,As,Hg,Cu,Mo
- 1988/89 Expo sampled for Au,As,Bi,Se,Te
- 1989 Area B sampled for As,Se
- 1989 RedDog South sampled for Au,As,Cu
- 2005 sampled for 32 element ICP

Nahwitti Lake

Kains Lake

HOLBERG

Holberg Inlet

Soil Geochemistry Anomalies

- Copper Gold Moly
- Zinc Silver Lead

Hushamu Soil Geochemistry Molybdenum ppm

- > 34 (98th percentile)
- 19 to 34 (95th percentile)
- 12 to 19 (90th percentile)
- 7 to 12 (80th percentile)
- 3 to 7 (50th percentile)
- < 3 ppm or no data

Sample Key

- Soil Sample
- Rock Sample (float)
- Rock Sample > 60ppm Mo

LUMINA COPPER CORPORATION

HUSHAMU PROPERTY

Rock and Soil Geochemistry (Moly ppm)

Date	APR 2006	Scale	1:25,000
UTM Zone	UTM9 - NAD83	Mining District	NANAIMO
N.T.S.	92L	State/Province	BC

Figure 9c


Soil Sample Key
1969 sampled for Mo and Zn
1969 & 72 sampled for Cu
1981 RedDog sampled for Au,As,Cu,Mo
1982 Utah Mines sampled for Au,As,Hg,Cu,Mo
1988/89 Expo sampled for Au,As,Bi,Se,Te
1989 Area B sampled for As,Se
1989 RedDog South sampled for Au,As,Cu
2005 sampled for 32 element ICP
566000 mE
570000 mE
572000 mE
574000 mE
576000 mE
578000 mE
580000 mE
582000 mE
584000 mE
586000 mE
588000 mE
590000 mN
5622000 mN
5620000 mN
5618000 mN
5616000 mN
5614000 mN
5612000 mN
5610000 mN
5608000 mN
5606000 mN
Nahwitti Lake
Kains Lake
HOLBERG
H o l b e r g I n l e t
Red Dog
Soil Geochemistry Anomalies
Copper Gold Moly
Zinc Silver Lead
Hushamu Soil Geochemistry
Zinc ppm
> 194 (98th percentile)
119 to 194 (95th percentile)
86 to 119 (90th percentile)
59 to 86 (80th percentile)
32 to 59 (50th percentile)
< 32 ppm or no data
Sample Key
Soil Sample
Rock Sample (float)
Rock Sample > 500 ppm Zn
LUMINA COPPER CORPORATION
HUSHAMU PROPERTY
Rock and Soil Geochemistry
(Zinc ppm)
EQUITY
Date APR 2006
Scale 1:25,000
Figure 9d
U.T.M. Zone UTM9 - NAD83
Mining District NANAIMO
N.T.S. 92L
State/Province BC