


Smiley NW



Ministry of Energy & Mines
Energy & Minerals Division
Geological Survey Branch


**ASSESSMENT REPORT
TITLE PAGE AND SUMMARY**

TITLE OF REPORT [type of survey(s)] <u>PROSPECTING</u>		# TOTAL COST <u>2100</u>
AUTHOR(S) <u>J. T. SHEARER, M.Sc., P. Geo</u>	SIGNATURE(S) 	
NOTICE OF WORK PERMIT NUMBER(S)/DATE(S)	YEAR OF WORK <u>2010</u>	
STATEMENT OF WORK - CASH PAYMENT EVENT NUMBER(S)/DATE(S)	<u>EVENT # 4779472</u>	
PROPERTY NAME	<u>SMILEY NW</u>	
CLAIM NAME(S) (on which work was done)	<u>SMILEY Northwest 592881</u>	
COMMODITIES SOUGHT	<u>LIMESTONE</u>	
MINERAL INVENTORY MINFILE NUMBER(S), IF KNOWN	<u>N/A</u>	
MINING DIVISION	<u>Nanaimo</u>	NTS <u>92L/7W</u>
LATITUDE	<u>50 ° 24 ' 10 "</u>	LONGITUDE <u>126 ° 49 ' 50 "</u> (at centre of work)
OWNER(S)	1) <u>J. T. SHEARER</u> 2) _____	
MAILING ADDRESS	<u>Unit 5 - 2330 TYNER ST. PORT COQUITLAM, B.C.</u>	
OPERATOR(S) [who paid for the work]	<u>V3C 221</u>	
1) <u>Homegold Resources Ltd</u>	2) _____	
MAILING ADDRESS	<u>As above</u>	
PROPERTY GEOLOGY KEYWORDS (lithology, age, stratigraphy, structure, alteration, mineralization, size and attitude):	<u>Triassic Quaternary Limestone intruded by "granite" to the north.</u>	
REFERENCES TO PREVIOUS ASSESSMENT WORK AND ASSESSMENT REPORT NUMBERS	<u>Assessment Report</u>	

(OVER)



TYPE OF WORK IN THIS REPORT	EXTENT OF WORK (IN METRIC UNITS)	ON WHICH CLAIMS	PROJECT COSTS APPORTIONED (incl. support)
GEOLOGICAL (scale, area)			
Ground, mapping _____			
Photo interpretation _____			
GEOPHYSICAL (line-kilometres)			
Ground			
Magnetic _____			
Electromagnetic _____			
Induced Polarization _____			
Radiometric _____			
Seismic _____			
Other _____			
Airborne _____			
GEOCHEMICAL			
(number of samples analysed for ...)			
Soil _____			
Silt _____			
Rock _____			
Other _____			
DRILLING			
(total metres; number of holes, size)			
Core _____			
Non-core _____			
RELATED TECHNICAL			
Sampling/assaying _____			
Petrographic _____			
Mineralographic _____			
Metallurgic _____			
PROSPECTING (scale, area) _____		592881	2100
PREPARATORY/PHYSICAL			
Line/grid (kilometres) _____			
Topographic/Photogrammetric (scale, area) _____			
Legal surveys (scale, area) _____			
Road, local access (kilometres)/trail _____			
Trench (metres) _____			
Underground dev. (metres) _____			
Other _____			
TOTAL COST			2100





Contact Us ▶ Help ⓘ
Printer Version 🖨️

B.C. HOME

Mineral Titles

Mineral Titles Online

**Mineral Claim
Exploration and
Development
Work/Expiry Date
Change**

Mineral Claim Exploration and Development Work/Expiry Date Change

Confirmation

Recorder: SHEARER, JOHAN THOM (124452) Submitter: SHEARER, JOHAN THOM (124452)
Recorded: 2010/JUL/29 Effective: 2010/JUL/29
D/E Date: 2010/JUL/29

- Select Input Method
- Select/Input Tenures
- Input Lots
- Link Event Numbers
- Data Input Form
- Upload Report
- Review Form Data
- Process Payment
- Confirmation

Confirmation

If you have not yet submitted your report for this work program, your technical work report is due in 90 days. The Exploration and Development Work/Expiry Date Change event number is required with your report submission. **Please attach a copy of this confirmation page to your report.** Contact Mineral Titles Branch for more information.

Smiley NW

Event Number: 4779472
Work Type: Technical Work
Technical Items: Geological, Preparatory Surveys, Prospecting
Work Start Date: 2010/APR/25
Work Stop Date: 2010/JUL/15
Total Value of Work: \$ 2100.00
Mine Permit No:

Summary of the work value:

Tenure Number	Claim Name/Property	Issue Date	Good To Date	New Good To Date	# of Days Forward	Area in Ha	Applied Work Value	Sub-mission Fee
592881	SMILEY NORWEST	2008/oct/14	2010/jul/30	2014/jul/30	1461	82.17	\$ 1903.59	\$ 131.56

- [Main Menu](#)
- [Search for Mineral / Placer / Coal Titles](#)
- [Search for Reserve Sites](#)
- [View Mineral Tenures](#)
- [View Placer Tenures](#)
- [View Coal Tenures](#)
- [MTO Help Tips](#)
- [Free Miner Landowner Notification](#)

Exit this e-service ▶

Financial Summary:

Total applied work value: \$ 1903.59
PAC name: J Shearer
Debited PAC amount: \$ 0.0
Credited PAC amount: \$ 196.41
Total Submission Fees: \$ 131.56
Total Paid: \$ 131.56

Please print this page for your records.

The event was successfully saved.

Please use **Back** button to go back to event confirmation index.

Back



GEOLOGICAL and PROSPECTING ASSESSMENT REPORT

ON THE SMILEY NW PROJECT

**BC Geological Survey
Assessment Report
31800**

**TENURE #592881
NIMPKISH LAKE AREA, VANCOUVER ISLAND, B.C.
NANAIMO MINING DISTRICT
N.T.S. 92L/7W (92L.036+046)
LATITUDE 50°24'10", LONGITUDE 126°49'50"**

for

**Homegold Resources Ltd.
Unit 5-2330 Tyner Street,
Port Coquitlam, B.C.
V3C 2Z1
Phone: 604-944-6102**

by

**J. T. Shearer, M.Sc., P.Geo.
Unit 5-2330 Tyner Street,
Port Coquitlam, B.C.
V3C 2Z1**

Phone: 604-970-6402 / Fax: 604-944-6102

Website: www.HomegoldResourcesLtd.com

E-mail: jo@HomegoldResourcesLtd.com

September 1, 2010

Fieldwork completed between April 25, 2010 and July 15, 2010

TABLE OF CONTENTS

	<u>Page</u>
LIST OF FIGURES and TABLES	ii
SUMMARY	iii
INTRODUCTION	1
LOCATION and ACCESS	2
FIELD PROCEDURES	2
LIST of CLAIMS.....	4
HISTORY	5
REGIONAL GEOLOGY	8
LOCAL GEOLOGY and 2010 PROGRAM	13
PREVIOUS DIAMOND DRILLING	15
CONCLUSIONS and RECOMMENDATIONS	17
REFERENCES	18
APPENDICES	
Appendix I Statement of Qualifications	20
Appendix II Statement of Costs	21

LIST OF ILLUSTRATIONS and TABLES

ILLUSTRATIONS

		<u>Following Page</u>
FIGURE 1	Location Map	iv
FIGURE 2	Google Image, General Area	3
FIGURE 3	Claim Map.....	6
FIGURE 4	Detail Claim Map.....	7
FIGURE 5	Regional Geology, 1:63,240	9
FIGURE 5a	Detail Regional Geology	10
FIGURE 6	2010 Travers Area, 1:5,000	14

TABLES


		<u>Page</u>
TABLE I	List of Claims.....	4

SUMMARY

- 1) The Smiley Norwest (total 4 cells) Mineral Claim was located in 2009 to cover a potential of an extensive zone of bleached white limestone.
- 2) The claims are west of Beaver Cove, 24 km south of Port McNeil.
- 3) Previous work for high brightness filler CaCO_3 including limited diamond drilling that was done for Industrial Fillers (Pluess Stauffer, OMYA) in the late 1980's.
- 4) The claims are underlain by intrusive despite being shown on the Regional Map as having Quatsino.
- 5) High brightness (up to 91.21%) and purity (up to 56% CaO)(99.68% CaCO_3) have been obtained from preliminary sampling to the south.
- 6) Work in 2010 shows that the area mapped previously along the access road as limestone is actually intrusive.
- 7) Future work should include (a) detail geological mapping along zone 100m wide from intrusive contact, (b) reconnaissance magnetometer lines throughout the property to identify the presence of blind intrusive bodies or dykes and (c) further wide spaced short diamond drill holes along the intrusive contact to test for higher brightness zones.

Respectfully submitted,

J. T. Shearer, M.Sc., P.Geo.



INTRODUCTION

The Smiley property was staked in October 2009 to cover the potential of fairly pure, white limestone, which had been known from rock exposures on old logging roads in the general area. Recent logging by WFP has exposed the area along the mapped limestone-intrusive contact.

The present program consisted of prospecting in the later part of 2010. Prospecting and mapping at a scale of 1:5,000 was completed in 2010 along the overgrown access roads.

Previously, the area was examined by Achermann and Duncan G. Ogden for Industrial Fillers and by David Coffin for Vanguard Consulting between June 15 and 19, 1988. A short diamond drilling program was conducted to the west of the property between August 2 and August 10, 1988. Some regional geological mapping was completed by Howard Brown for Pleuss Stauffer in 1984.

Initial discussions have taken place with WFP Logging on the possibility of using the private deep water dock facilities at Beaver Cove. In the past the Kelsy Bay-Beaver Cove Ferry used the ramp and the Nimpkish Iron operation also loaded barges at Beaver Cove.

LOCATION and ACCESS

The property is located on the east of Nimpkish lake and approximately 6 km southwest of the deep harbour at Beaver Cove, on Vancouver Island's Northeast coast. Port McNeil, the closest supply point to the property, lies approximately 10 air-km or 24 road-km to the northwest. Port McNeil is capable of providing accommodation, contract excavators and the other usual requirements for an exploration program.

Access to the property is gained by driving south from Port McNeil along B.C. Highway 19 (Island Highway) for a distance of 14 km, turning east onto the Beaver Cove access road. A series of branch logging roads provide access to most parts of the claim group.

The Smiley NW property occupies a portion of the transition between the lowlands of Vancouver Island's northeast coast and the rugged mountain ranges to the south. Elevations on the property range from 25 metres to 600 metres a.s.l. Much of the property is a north facing side hill with an average slope of 12°. The drainage has a trellis pattern but creeks can be expected to flow usually during run-off periods due to the limestone bedrock.

The claims are within TFL 37 owned by Western Forest Products (WFP), who operate numerous camps, the largest being Woss where the Forestry Engineering office is located. A unique feature of TFL 37 is the still operating logging railway, which transports logs to the sorting and shipping facility at Beaver Cove.

FIELD PROCEDURES

Sample locations (see Appendix III) were established using a Garmin GPS Unit. The field data was downloaded into the Garmin Mapsource program for plotting.

A
N



Figure 2 Google Image General Area

LIST of CLAIMS

The property consists of one claim totalling 4 cells in Table 1 and Figure 4.

TABLE I
List of Claims

Claim Name	Tenure #	Size (ha)	Cells	Date Located	Current Anniversary Date*	Owner
Smiley Norwest	592881	82.17	4	October 14, 2008	July 30, 2014	J. T. Shearer

* after common dating and application of assessment work documented in this report.

Mineral title in British Columbia is acquired by locating claims in the proscribed manner as outlined in the MINERAL ACT and regulations. Title is maintained by filing appropriate assessment work in the amount of \$4 per ha for the first 3 years and \$8 per ha thereafter.

Under the present status of mineral claims in British Columbia, the consideration of industrial minerals requires careful designation of the products end use. An industrial mineral is a rock or naturally occurring substance that can be mined and processed for its unique qualities and used for industrial purposes (as defined in the *Mineral Tenure Act*). It does not include "Quarry Resources". Quarry Resources includes earth, soil, marl, peat, sand and gravel, and rock, rip-rap and stone products that are used for construction purposes (as defined in the *Land Act*). Construction means the use of rock or other natural substances for roads, buildings, berms, breakwaters, runways, rip-rap and fills and includes crushed rock. Dimension stone means any rock or stone product that is cut or split on two or more sides, but does not include crushed rock.

HISTORY

The area has long been known for its timber production. Several skarn copper-magnetite showings were found in 1929 southeast of the Smiley NW Claims along Kinman Creek and Smith Creek.

There are several assessment reports available on the area near the Smiley NW Group as follows:


Assessment Report

Number

094	Menzies, M., and Brynelsen, B. O., 1953: Trenching and Mapping for Noranda.
10986	Quin, and DeCarle, 1983: Input EM and Airborne Magnetometer 33.7 line km also plotted on a 1:10,000 orthophoto with total magnetics and horizontal coil EM anomalies for Mintek Resources
12348	Morton J. W., 1984: Geochemistry for Mintek Resources
18850	Soux, C. and Coffin, D., 1988: Diamond Drill Program Report for Industrial Fillers Ltd. (Pleuss Stauffer) two 150m short holes, widely spaces.

Geological mapping was carried out by Pleuss Stauffer geologist, Howard Brown in several places on the northern Vancouver Island. A reduced summary version of Brown's mapping is shown in Soux and Coffin (1988).

To the south of Nimpkish Lake a magnetite skarn produced several million tonnes in the late 1950's and early 1960's from the Klannick Iron Deposit. Mineralization in the general area was originally discovered around 1900.



SMILEY NW




Figure 4 Detail Claim Map

REGIONAL GEOLOGY

The Nimpkish-Beaver Cove Area was most expertly mapped by H. Gunning in the field seasons 1929 to 1931 who established a more detailed stratigraphy and named the Karmutsen Formation and Bonanza Group.

These maps were published by Hoadley (1953) along with Memoir 272 (Geology and Mineral Deposits of the Zeballos-Nimpkish Area, Vancouver Island, B.C.). More recently Mueller and Roddick completed 1:250,000 mapping of the 29L sheet for the Geological Survey of Canada and published Paper 74-8 on the general Area (Muller, Northcote and Carlise, 1974).

The area is primarily composed of intermediate volcanic sequences of the Karmutsen Formation conformably overlain by Quatsino Formation Limestone. A major antiformal structure occurs from which exposes Triassic Parson Bay mixed sedimentary rock and Lower Jurassic Bonanza Group intermediate to felsic volcanic sequences. Rock units generally trend to the northwest, displaying a series of open folds with gentle dips east and west.


All of the above units have been intruded by members of the intermediate to felsic Island Intrusions of Upper Jurassic age. These intrusions have caused both skarn and other hydrothermal metal deposits at numerous locations on Vancouver Island.

Major faults tend to lie sub-parallel to the fold structures, although cross-faulting has been mapped.

Hoadley (1953) describes the Quatsino Formation (page 17) as follows:

“The Quatsino Formation consists almost entirely of limestone, with a few thin flows of andesite or basalt. The limestone is fine to coarsely crystalline, and ranges from white to black, with various intermediate colours. Towards the base, it tends to be exceedingly fine grained, and grey and brownish or buff colours are characteristic. Midway of the formation the colours are predominantly white or grey, but towards the top the limestone becomes dark grey to black, due to a varying quantity of carbonaceous matter, and the formation grades upward into argillites and impure limestones of the overlying Bonanza group. Even at the top, however, light grey or even white beds are interbedded with the darker varieties in the upper part of the formation but in the lower part, where white to brownish grey and buff colours predominate, it is poorly preserved. In the upper part, too, the beds are generally thin, thicknesses of ½ inch and less being common and formation 2 or 3 feet uncommon. The formation as a whole is dominantly a high-calcium limestone. The rock is too jointed in many places to serve as a building stone, but where the beds are least deformed and well removed from intrusions, as from Beaver Cove to Bonanza Lake, it could be extracted in blocks sufficiently large for ordinary structural purposes. Within a mile or two of bodies of the Coast intrusions, the limestone may be highly contorted and extremely jointed and fractured, cut by many acidic dykes, and partly altered to lime-silicate minerals, iron oxides, magnetite and hematite, and by sulphides of copper, iron, zinc, and lead.”

The lower part of the Quatsino limestone is well exposed on the east side of Nimpkish Lake, 2 miles from the outlet. At its base there is a small fault, trending 070° east, which throws the



SMILEY NW




Figure 5a Detail Regional Geology

underlying volcanic rocks up against the limestone. The volcanic rocks, which include andesite, amygdaloidal basalt, and sheared agglomerate, are exposed for 500 yards or more to the south and are underlain by at least 50 feet of grey and white mottled limestone, which at its base becomes argillaceous and well bedded and rests conformably on a slightly sheared and altered amygdaloidal flow. A second smaller bed of limestone lies conformably in these volcanic rocks a few hundred yards farther south. At this locality, the lower part of the Quatsino Formation is composed of interbedded limestone and volcanic flows.

For about a mile on the east side of Nimpkish Lake opposite Halfway Islands, the rocks at and near the base of the Quatsino formation are exposed at low water (Hoadley, 1953). There, the top of the underlying volcanic group is rolling and irregular and remnants of the overlying Quatsino limestone have been preserved in one or two saucer-shaped low-lying areas. The relations between the limestone and underlying volcanic rocks are complex. In one place, 1km due south of Halfway Islands, an irregular, 3-foot bed of light grey, fine-grained, limestone, some distance below the base of the Quatsino Formation, is overlain and underlain by andesitic lavas, and is contorted and slightly faulted. Farther south are amygdaloidal basalts and a peculiar fragmental rock, the latter consisting of grey to greenish or brownish dense limestone nodules or rounded fragments, rarely more than 1 inch or 2 inches in diameter, in a matrix of green and reddish andesite and basaltic fragments from ¼ inch to 18 inches in diameter, some of them resembling bombs. This rock might be termed a breccia, but it has the appearance of having been formed by incorporation of volcanic ejectamenta in a calcareous mud, possibly with the addition of a few angular fragments of limestone (Hoadley, 1953).

Farther south, at the first good expose of its base, in this locality, the Quatsino Formation was found by Hoadley to be underlain by andesitic flows containing several irregular gobs, up to 5 feet across, of limestone, the whole intruded by irregular and curving andesite dykes. The Quatsino limestone overlies this material and dips gently westward but contains irregular to lenticular dyke-like masses of andesite.

At one place on the small peninsula northeast of Halfway Islands, the base of the limestone is again well exposed. There, the limestone is apparently lying on green to purplish andesite flows and fragmental rocks, but it is intruded by numerous dykes of similar appearance to the lavas. Also, the dykes contain many large and small fragments of limestone. The limestone itself is massive or poorly and irregularly bedded. Farther south, the underlying andesite and amygdaloidal basaltic volcanic rocks are exposed for almost 900 feet to the small point east of the north end of Halfway Islands. There, pure white, crystalline, massive limestone, banded in grey shades for 8 feet above the base, overlies green, rusty, pyritic andesite, the contact striking 030° and dipping 30° southeast (Hoadley, 1953).

Most of the intrusive rocks of Vancouver Island form part of the Coast intrusions, which were emplaced during Jurassic or Cretaceous time and which now occupy much of the Coast Mountain area of British Columbia. They are holocrystalline, igneous rocks that range in colour from pink and brown to grey and dark greenish grey, and in composition from basic to acidic, with rocks of the granite clan predominating. They form sills, dykes, stocks and batholithic bodies in the Vancouver group and are of great economic significance in that most of the mineral deposits of the region are believed to be genetically related to them.

On northern Vancouver Island, these intrusive rocks are largely confined to long, narrow, northwesterly trending belts separated by somewhat wider belts of Upper Triassic volcanic and sedimentary rocks. The areas of intrusive rocks are, in detail, irregular and discontinuous. Regionally, however, they form bands 2 to 5 miles in width that can be traced along the strike of the volcanic rocks for many miles (Hoadley, 1953).

Elsewhere in British Columbia, there are sources of white limestone, most notably at Benson Lake (about 20 km directly west of the Smiley Claims) operated by IMASCO and several producers on Texada Island.

Texada Island has produced high quality white limestone from small deposits over the course of its history. There are no extensive white limestone deposits on Texada island (Mathews and MacCammon, 1957); however, there are workable deposits situated on the island. The Blubber Bay quarries of Pacific Lime and their subsequent owners mined white limestone and stockpile it for specialty markets. The white limestone sells for a premium and so was able to be selectively mined. Beale Quarries Limited also produced white limestone from a body south of Quarry No. 5 in the vicinity of Lot 499.

White limestone was produced from Lot 500 south of Van Anda and south of the Lafarge quarry on Lot 499. The stone was pulverized and bagged for shipment on the property until Fred Beale opened a stucco plant in the old smelter building in Van Anda. From the 1940's until 1959 Lot 500 supplied the stucco plant with white limestone until Imperial Limestone Company Limited gained control of the operation. J. A. Jack & Sons Incorporated of Seattle, Washington own Imperial Limestone. The limestone is shipped to the Seattle processing plant and sold for agricultural limestone, stucco, chicken grit and other pulverized limestone products.

Imperial Limestone built a crushing and barge load-out installation at Butterfly Bay (Spratt Bay). In 1975 the stucco plant in Van Anda was shut down and the building destroyed. Imperial built a new pulverizing and bagging plant at Butterfly Bay as a result. The plant was eventually phased out when freight costs became too high to operate it. All stone processing is now carried out in Seattle.

The largest white limestone body is at Texada Quarrying Ltd. (formerly Ideal Cement) Paxton Lake Zone. The Paxton Lake deposit has been developed on 3 wide levels but has recently been inactive due to, temporarily, low priced white limestone sourced out of southeast Alaska. This deposit, despite it's very high brightness, is now closed due to high cost of freight. The origin of the white limestone is controversial. The genesis of the white rock may be due to metasomatism, stratigraphic control, hydrothermal alteration or volcanic intrusives. The white colour is probably the result of the bleaching of black limestone by hydrothermal fluids percolating along a system of vertical joints.

LOCAL GEOLOGY and 2010 PROGRAM

The Smiley NW property is underlain on the Northeast corner by Quatsino Formation limestone in conformable contact with undifferentiated Karmutsen Formation basalt and andesite, all of which has been intruded by a northwesterly trending body of coarse grained biotite quartz monzonite. Thin sills and dykes of fine grained diabase cut the limestone but were not seen to cut the monzonite.

Previous work on the property divided the limestone into Upper and Lower members. The Upper member is medium to dark grey in colour and occasionally contains silica. Interbeds of white weathering, off white to light grey limestone are also present. The Lower member is generally white to light grey and fine grained, except where recrystallized and has thin beds of dark grey and cherty material. Pyritic lens both conform to and cross bedding.

Bedding in the limestone generally trends northerly. A synclinal axis runs through the centre of Smiley 4 in the lower Limestone, passing east of Smiley 5 along the top of a small ridge of Upper Limestone. Dips flatten quickly away from the axis in either direction, indicating a fairly broad, shallow structure.

The pyritic lens are within areas, which have been replaced by vitreous to cloudy silica, with blebs and poorly formed crystals of pyrite filling random fracture planes. They are defined by remnant bedding planes and by fractures trending northeasterly, sub-parallel to the limestone/monzonite contact. The lens are most prominent in the southern part of the property. Pyritic lens increase with proximity to the volcanic/limestone contact and proximity to the monzonite body. They appear to be the result of hydrothermal fluids, which moved along the planes of weaknesses during intrusion of the monzonite body.

The intrusive-limestone contact is will exposed on the Island Highway on Smiley 1 mineral claim, Figure 7, at a point 28.3 km south of the Port McNeil-Highway junction. Minor rusty weathering skarn has developed along the contact within the intrusive. Small sill-like bodies of intrusive were also noted within the limestone a short distance from the contact. The contact on the highway is oriented 140°/65° NW.

The intrusive are well exposed in the northeast portion of the claims. Hoadley (1953) characterizes the pluton east and southeast of Nimpkish Lake as essentially granodiorite, although parts of it are quartz monzonite and in places it approaches granite in composition. In a few thin sections, especially those of the granites, interstitial micrographic intergrowths of quartz and alkali feldspar were observed (Hoadley, 1953). Alteration of the feldspars to sericite, zoisite and albite is common. Green hornblende is the dominant ferromagnesian constituent, but in places is exceeded by dark brown biotite, in ragged flakes. Some of the biotite is derived from the hornblende and both biotite and hornblende have been altered in part to chlorite (Hoadley, 1953).

Throughout this entire area, the intrusive rocks are Lithologically very similar and except for the more basic border phases all belong to the granite clan, with granodiorite, quartz monzonite and granite the most common types.




FIGURE 6

Figure 6 2010 Traverse Area Smiley NW Project

PREVIOUS DIAMOND DRILLING (to the Southwest of Property)

In 1988, two 150 metre BQ diamond core holes, PT-88-1 and 88-2, were completed by Pluess Stauffer to the southwest of the Smiley Northwest property. The holes were spotted at the road accessible sites located approximately 750 metres apart at about the same elevation. Diamond drill hole PT88-1 was spotted 700m @ Az. = 295° from the monzonite contact. PT-88-2 was spotted at 750m @ Az. = 345° laterally and 5m vertically lower from PT-88-1.

PT-88-1 was collared in then cut 134.5 metres of generally light grey to white limestone, with one 8.5m section of grey limestone centred at 41 metres. The section from 17m to 27.5m contains what appear to be three andesitic dykes, which have been silicified and pyritized; the dykes represent 75% of this section.

The section from 134.5m to 137.5m contains 1m of amygdaloidal andesite followed by 2m of white limestone. The section from 137.5m to 152.5m (bottom of hole) contained greenish grey andesite, which has been altered to chlorite and epidote in places.

PT-88-2 was collared in, then cut, 65 metres of generally light grey to grey limestone. From 65m to 88m the hole cut alternating lens of generally light grey to white limestone and intermediate volcanics; several of the contacts have been altered by hydrothermal fluids. From 88m to 152m (bottom of hole) the hole cut greenish grey andesite, which has been altered, to chlorite and epidote in places. The limestone/volcanic contact has been altered to silica and pyrite for a length of 5 metres.

Both holes indicated that the limestone/volcanic contact is flat or dipping very gently along the section Az. = 295°, which is consistent with a general strike WNW-ESE. The calculated dip based on this assumption is approximately -5° to the south or southeast. Until fill-in data is available, the assumption should be simply that the contact has a shallow dip in a southerly direction (Soux & Coffin, 1988).

The north-south trending synclinal axis mapped in limestone does not appear to be representative of the contact orientation. This is probably a result of either a) location of one or both of the holes over a local rise in the paleotopography, or b) discrepancy resulting from the movement of intervening faults.

The two diamond drillholes were completed by Homegold in 2001 to the southwest of the Smiley NW property. Hole NIMP-01-02 was collared approximately 200m northwest of the intrusive contact not far from Highway 19. On surface down to 2.95m is a very white mostly medium to finely crystalline limestone. Traces of pyrite were observed along minor high angle fractures. However, below 2.95m, a short section 2.95m to 4.92m of medium grey limestone was encountered. A very minor amount of intrusive dyke, which had been stretched and boudinaged with rounded fragments between 3.59m and 3.62m.

White limestone appears again between 4.92m and 9.55m. Below 9.55m to 24.23m is a light grey limestone, which is characterized by aligned vuggy sections, which appear to be related to whiter layers or laminae. Near the bottom of the hole (24.23-27.43m) is white limestone. In hole MIMP-01-02 the whiter sections are distinctly finer grained.

In hole NIMP-01-03, which is closer to the intrusive-carbonate contact (approximately 50m north of contact), the white limestone section is thicker (to 19.51m) and more continuous. The white section is also slightly coarse crystalline in Hole NIMP-01-03 than the distinctly finer grained white limestone farther removed from the intrusive contact.

Traces of dyke fragments are also noted in hole NIMP-01-03 at 3.20m as 3mm wide rounded greenish lenses, which suggests considerable plastic flow. A dark green andesitic dyke was encountered between 19.51m and 21.18m as a uniformly dark green, very fine grained intrusive with minor pyrite along fracture surfaced. The limestone below the dyke is noticeably darker grey than the upper limestone interval and also finer crystalline. Dark chloritic coated slickensides throughout the lower limestone unit gives an even darker overall impression. Minor sparry calcite lenses were noted at the end of Hole at 24.38m.

Work in 2010 on the Smiley NW Project showed that the access road, Figure 6, showed only intrusive outcrops along the entire length of the road despite having the regional geology map indicating that the limestone contact should have been exposed.

CONCLUSIONS and RECOMMENDATIONS


Work in 2010 failed to encounter any limestone along the overgrown access road. Only intrusive rocks were observed.

Diamond drilling in 1988 and 2001 encountered the Karmutsen contact higher than would have been expected from an interpretation of surface mapping. This may be because of a local rise(s) in the paleotopography. The apparent dip, from drill intersections, of the contact at a shallow angle to the south is influenced by intervening faults, and requires further testing to ensure its reliability.

Analytical sampling of the property, especially proximal to the monzonite, should include analysis of the hydrothermal alteration for precious and other metal content. Similar alteration of these units elsewhere contains economic gold mineralization.

The general condition of the limestone/intrusive contact could be tested by the drilling of one hole on section with 88-01 and 88-2, from an existing road location approximately 850m north of 88-2. This hole would be collared near the Upper/Lower contact, thereby testing a complete section of the later. A series of holes should also be drilled around NIMP-01-03 in order to test continuity of section over shorter distances to the east. This information could then be used to enhance the present structural interpretation prior to in fill drilling.

Respectfully submitted,



J. T. Shearer, M.Sc., P.Geo.
Homegold Resources Ltd.

REFERENCES

- Brown, H. J., June 1984:
Geology of the Port McNeill (sic) Quarry Area Map Only; Private Report.
- Dolmage, V., 1919:
Quatsino Sound and Certain Mineral Deposits of the West Coast of Vancouver Island, B.C., Geological Survey of Canada, Summary Report, 1918.
- Fischl, P., 1992:
Limestone and Dolomite Resources in British Columbia. British Columbia Mineral Resources Division, Geological Survey Branch Open File 1992-18, 150 p.
- Gunning, D. F., 1991:
Rocks in Motion, Imasco, Surrey, B.C., Conference Proceeding, Industrial Minerals Forum, 1991. Page 167-169.
- Gunning, H. C., 1930:
Geology & Mineral Deposits of the Quatsino-Nimkish Area, Vancouver Island, Geological Survey of Canada, Summary Report 1929 Pt. A P94-143.
- Gunning, H. C., 1929/31 and Hoadley, J. W., 1952:
Geology of Nimpkish Map Sheet @ 1" = 1 mile; GSC Map 1029A.
- Hoadley, J. W. 1953:
Geology and Mineral Deposits of the Zeballos-Nimpkish Area, Vancouver Island, B.C., Geological Survey of Canada, Memoir 272, 82 pp.
- McCammon, J. W., 1968:
Limestone Deposits at the North End of Vancouver Island, Mines and Petroleum Resources Report, 1968. M.M.A.R. Page 312-318.
- Mathews, W. H. and McCammon, J. W., 1957:
Calcareous deposits of Southwestern British Columbia. British Columbia Department of Mines Bulletin No. 40, 105p.
- Muller, J. E., Northcote, K. E. and Carlise, D., 1974:
Geology and Mineral Deposits of Alert-Cape Scott Map Area, Vancouver Island, B.C. Geological Survey of Canada, Paper 74-8, 77 p.
- Muller, J. E., 1973 and Roddick, J. A., 1980:
Geology of Alert Bay – Cape Scott @ 1:250,000, map 1552A.
- Shearer, J. T., 1998:
Mining Permit Application Summary on the South Slesse Limestone Quarry MX7-114, for I.G. Machine & Fibers Ltd., Dated January 10, 1998, 23 pages.

1999:

Diamond Drilling Report on the Davies Bay Limestone Deposit Filed for Assessment Credit for Tilbury Cement Ltd.

2000:

Diamond Drilling Report on the Ravens Bay-Will Claims Tex Limestone Deposit filed for Assessment Credit, for Chemical Lime Corp. Inc.

2001:

Geological, Prospecting and Diamond Drilling Assessment Report on the White Limestone Project, Smiley Property, July 16, 2001, 37pp. Assessment Report 26686

Soux, C. and Coffin, D., 1988:

Diamond Drill Program Report on Tsulton Property for Industrial Fillers Ltd. (Pluess Stauffer), Assessment Report 17759, 28pp.

Webster, I. C. L. and Ray, G. E., 1990:

Geology and Mineral Occurrences of Northern Texada Island NTS 92F/9, 10, 15. British Columbia Ministry of Energy, Mines and Petroleum Resources, British Columbia Geological Survey Open File 1990-3, 1 sheet.

APPENDIX I

STATEMENT of QUALIFICATIONS

SEPTEMBER 1, 2010


Appendix I

STATEMENT OF QUALIFICATIONS

I, JOHAN T. SHEARER, of 3572 Hamilton Street, in the City of Port Coquitlam, in the Province of British Columbia, do hereby certify:

1. I am a graduate of the University of British Columbia (B.Sc., 1973) in Honours Geology, and the University of London, Imperial College (M.Sc., 1977).
2. I have over 35 years experience in exploration for base and precious metals and industrial mineral commodities in the Cordillera of Western North America with such companies as McIntyre Mines Ltd., J.C. Stephen Explorations Ltd., Carolin Mines Ltd. and TRM Engineering Ltd.
3. I am a fellow in good standing of the Geological Association of Canada (Fellow No. F439) and I am a member in good standing with the Association of Professional Engineers and Geoscientists of British Columbia (Member No. 19,279). I am also a fellow of the Geological Society (London) and Society of Economic Geologists (SEG).
4. I am an independent consulting geologist employed since December 1986 by Homegold Resources Ltd. at #5-2330 Tyner St., Port Coquitlam, B.C.
5. I am the author of a report entitled "Geological and Prospecting Report on the Smiley NW Claims, Nanaimo Mining Divisions" dated September 1, 2010.
6. I have visited the property between July 25, 2000 and June 3, 2001. I have carried out mapping and sample collection and am familiar with the regional geology and geology of nearby properties. I have become familiar with the previous work conducted on the Smiley claims by examining in detail the available reports and maps and have discussed previous work with persons knowledgeable of the area.
7. I have an Open Pit Supervisor Ticket (#98-3550) for daily supervision duties.
8. I own an interest in the Smiley Claims and own Homegold Resources Ltd.

Dated at Port Coquitlam, British Columbia, this 1st day of September 2010.



J.T. Shearer, M.Sc., F.G.A.C., P.Geo.
Quarry Supervisor #98-3550
September 1, 2010

APPENDIX II

STATEMENT of COSTS

SEPTEMBER 3, 2010

APPENDIX II

STATEMENT of COSTS
SMILEY NW PROJECT

Wages & Benefits	GST	Total without GST
J. T. Shearer, 1 day @ \$700/day	\$ 35.00	700.00
Ron Olynyk, 1 day @ \$400/day	20.00	400.00
Wages Sub-Total	\$55.00	1,100.00
Expenses		
Truck Rental, fully equipped 4x4, 2 days @ \$98.50/day		197.00
Fuel		175.00
Ferries		98.00
Hotel, Meals & Camp Supplies		120.00
Map Preparation and Drafting		100.00
Report Preparation		700.00
Word Processing and Reproduction		150.00
Expenses Sub-Total		\$ 1,540.00
Total	\$ 55.00	\$ 2,640.00

Event # 4779472
Filed \$2,100.00 July 29/10
Applied \$1,903.59