



ASSESSMENT REPORT TITLE PAGE AND SUMMARY

TITLE OF REPORT: Assessment report for 2012 rock chip sampling on the Wilcum1-7 claims, Wilcum High Peak, Kimberley, South-East British Columbia, Canada.

TOTAL COST: \$ 30,907.36

AUTHOR(S): Luke van der Meer
SIGNATURE(S): "signed"

NOTICE OF WORK PERMIT NUMBER(S)/DATE(S): N/A
STATEMENT OF WORK EVENT NUMBER (5430397) /DATE (2013/FEB/05):

YEAR OF WORK: 2012

PROPERTY NAME: WILCUM HIGH PEAK

CLAIM NAME(S) (on which work was done):

WILCUM1-7: 522473, 522474, 522475, 522476, 777142, 777222, 777262.

COMMODITIES SOUGHT: LEAD, ZINC, COPPER, GOLD.

MINERAL INVENTORY MINFILE NUMBER(S), IF KNOWN: 082FNE052, 082FNE062, 082FNE066, 082FNE088, 082FNE152, 082FNE161.

MINING DIVISION: FORT STEELE

NTS: Map 082F 09E

BCGS: 082 F060 and 070

LATITUDE: 49 ° 36 ' 16 "

LONGITUDE: 116 ° 08 ' 05 " (at centre of work)

UTM Zone: **EASTING:** **NORTHING:**

OWNERS: 201966 LEITH, WILLIAM CUMMING 50.0%
146010 WALLACH, DAVID ARNOLD 50.0%


MAILING ADDRESS:

**P.O. Box 29191
Kelowna,
BC V1W 4A7
250-801-7744**

OPERATOR: DW EXPLORATION

MAILING ADDRESS:


**P.O. Box 29191
Kelowna,
BC V1W 4A7
250-801-7744**



REPORT KEYWORDS (PURCELL SUPERGROUP, PROTEROZOIC, THOLEITIC GABBRO, GRANOPHYRE, MASSIVE SULPHIDES, PYRITE, PYRRHOTITE)

REFERENCES TO PREVIOUS ASSESSMENT WORK AND ASSESSMENT REPORT NUMBERS:

TYPE OF WORK IN THIS REPORT	EXTENT OF WORK (in metric units)	ON WHICH CLAIMS	PROJECT COSTS APPORTIONED (incl. support)
GEOLOGICAL (scale, area)			
Ground, mapping		777142, 777262	\$ 10,102.45
Photo interpretation			
GEOPHYSICAL (line-kilometres)			
Ground			
Magnetic			
Electromagnetic			
Induced Polarization			
Radiometric			
Seismic			
Other			
Airborne			
GEOCHEMICAL (number of samples analyzed for ...)			
Soil			
Silt			
Rock	10 samples	777142, 777262	\$ 600.00
Other			
DRILLING (total metres, number of holes, size, storage location)			
Core			
Non-core			
RELATED TECHNICAL			
Sampling / Assaying	10 samples	777142, 777262	\$ 10,102.45
Petrographic			
Mineralographic			
Metallurgic			
PROSPECTING (scale/area)	14km2	777142, 777262	\$ 10,102.45
PREPATORY / PHYSICAL			
Line/grid (km)			
Topo/Photogrammetric (scale, area)			
Legal Surveys (scale, area)			
Road, local access (km)/trail			
Trench (number/metres)			
Underground development (metres)			
Othe			
		TOTAL COST	\$30,907.36



**ASSESSMENT REPORT FOR 2012 ROCK SAMPLING AND
PROSPECTING ON THE WILCUM CLAIMS—WILCUM HIGH PEAK
PROJECT, SOUTH-EASTERN BRITISH COLUMBIA, CANADA.**

FORT STEEL MINING DIVISION, SOUTH-EASTERN BRITISH COLUMBIA, CANADA

**B.C.G.S. 082 F060 and 070
N.T.S. Map 082F 09E
Latitude 49° 36' 16", Longitude 116° 08' 05"**

**For Owner-Operator:
DW Exploration
P.O. Box 29191
Kelowna, BC
BC V1W 4A7, Canada
250-801-7744**



**Author:
Luke van der Meer, PGeo, BSc-Geology
222-991 Hornby St
Vancouver, BC
V6Z 1V3, Canada**

**BC Geological Survey
Assessment Report
34079**

15 May 2013

EXECUTIVE SUMMARY

The Wilcum property is located in the Purcell Mountains approximately 30 km west-northwest of Cranbrook, BC. The property comprises a total of 3,620.181 ha (8945.662 acres) straddling the St. Mary's River, immediately east of St. Mary's Lake. Access to the property is readily available for 2WD vehicle to, and around the periphery, of the property along existing, well maintained logging roads. Older logging and exploration roads and trails are potentially available; however, they will likely require rehabilitation to re-establish them for use.

The Wilcum property is located approximately 15 km southwest of the former Sullivan Mine, a world class sedimentary exhalative deposit that produced until closure on December 21, 2001. The Sullivan Mine produced more than 160 million tons of ore grading 6.5% lead, 5.6% zinc and 67 g/t silver. Minor by-products included bismuth, cadmium, gold, indium, iron, sulphur and antimonial lead and tin concentrate.

The Sullivan Horizon (locus of the Sullivan Mine) is located at the contact between the lower and middle Aldridge formations, at the Lower - Middle contact ("LMC"). The stratigraphy underlying the property belongs the Proterozoic Aldridge Formation, specifically, the lower Aldridge Formation and, therefore, covers prospective strata in the footwall of the Sullivan Horizon.

A total of 41 man days were spent on field explorations by Dr W Lieth during 2012. During his field investigations Dr Leith constructed nine hand-trenches and test-pits (approximately 1-2ft deep) from which to collect prospective rock samples from excavated out crop and sub-cropping rocks.

Numerous undescribed rock samples were collected from nine pits located adjacent to the main haul road that transects the property and at the base of out cropping and sub cropping rocks of the underlying Aldridge Formation.

10 samples were selected by Dr Leith and Mr Wallach for analysis. These samples were numbered and dispatched to Acme Analytical Laboratories (Vancouver) Ltd. Two samples from Trench 1 and 3 contained anomalous values for Copper.

The Wilcum property covers prospective stratigraphy correlated to the Lower Aldridge Formation in the footwall of the Sullivan Horizon. Historically, the LMC has been the preferred horizon of interest for exploration; however, the Sullivan is, to date, unique among SEDEX deposits in that it is the only economic SEDEX occurrence identified to date in the Aldridge Formation. SEDEX deposits elsewhere in the world occur in clusters throughout the host stratigraphy.

For this reason, the LMC is the primary horizon of interest. However, sulphides (predominantly pyrrhotite) are ubiquitous in the Aldridge Formation and are abundant in the Lower Aldridge Formation. Furthermore, smaller massive sulphide bodies have been identified at a variety of levels within the Aldridge Formation, including within, and adjacent to the Wilcum property. Some of these occurrences have been interpreted as sedimentary exhalative in origin, including the Kootenay King mine (Past Producer). In addition, base metal massive sulphide veins or Proterozoic age have been identified, including those within the Northstar Corridor (Stemwinder, Northstar, Quantrelle). Proterozoic age base

metal mineralization occurrences include the St. Eugene Mine (Past Producer), the Vine Vein and the Fors occurrence. These occurrences are interpreted to confirm base metal sources active in the Proterozoic during deposition of the Aldridge Formation and may include sedimentary exhalative style mineralization.

TABLE OF CONTENTS

EXECUTIVE SUMMARY	2
TABLE OF CONTENTS	4
LIST OF FIGURES	5
LIST OF TABLES	5
1. INTRODUCTION.....	6
2. LOCATION AND ACCESS	8
3. PHYSIOGRAPHY AND CLIMATE.....	8
4. MINERAL TENURE AND CLAIM STATUS.....	9
5. EXPLORATION HISTORY.....	10
6. TERMS OF REFERENCE.....	12
7. REGIONAL GEOLOGY	12
Proterozoic Purcell Supergroup Stratigraphy	12
Aldridge Formation	13
Middle Aldridge.....	13
Laminated Siltstone markers	13
Upper Aldridge.....	13
Intrusives.....	13
Proterozoic Moyie Sills.....	14
Structure	14
Mineralization	15
Sullivan Mine Type	15
8. LOCAL GEOLOGY	17
Aldridge Formation.....	17
9. PROPERTY GEOLOGY	17
Granophyre	18
Iron-rich Bedrock Horizon	18
Sulphide-rich Sedimentary Float (minor base metals).....	19
Intrusives.....	19

10.	2012 WORK PROGRAM	20
11.	RESULTS.....	21
12.	CONCLUSIONS.....	22
13.	RECOMENDATIONS.....	23
14.	STATEMENT OF COSTS	24
15.	REFERENCES	26
16.	APPENDICIES	27
	APPENDIX 1: Certificate of the Author.....	
	APPENDIX 2: Rock sampling locations and analytical results.	
	APPENDIX 3: Sample Maps and Diagrams	
	Sample Location.....	
	Copper in rocks.....	
	Silver in rocks.....	
	Lead in rocks.....	
	Zinc in rocks.....	

LIST OF FIGURES

Figure 1: Property location map, South Eastern British Columbia, BC.....	6
Figure 2: property location map and claim outlines, West of Kimberly, BC, Canada.....	7
Figure 3: Local property location map, West of Kimberly, BC, Canada. Showing: claim boundaries, mineral occurrences, and primary drainages	9

LIST OF TABLES

Table 1: Summary list of the Wilcum claim group, Kimberley area, BC. Source: Mineral Titles Online (MTO).	9
Table 2: Summarized results for the 2011 rock sampling program	21

1. INTRODUCTION

The Wilcum property is located in the Purcell Mountains approximately 28 km west-northwest of Cranbrook, BC (Fig. 1 and 2). The properties comprises a total of 3,620.181 ha (8945.662 acres), straddling the St. Mary's River, immediately east of St. Mary's Lake (Fig. 3). Access to the property is readily available for 2WD vehicle to, and around the periphery, of the property along existing, well maintained logging roads. Older logging and exploration roads and trails are potentially available; however, they will likely require rehabilitation to re-establish them for use.

The Wilcum property is located approximately 15 km southwest of the former Sullivan Mine, a world class sedimentary exhalative deposit that produced until closure on December 21, 2001.

The Sullivan Mine produced more than 160 million tons of ore grading 6.5% lead, 5.6% zinc and 67 g/t silver. Minor by-products included bismuth, cadmium, gold, indium, iron, sulphur, antimony, lead, and tin concentrate.

The Sullivan Horizon (locus of the Sullivan Mine) is located at the contact between the lower and middle Aldridge formations, at the Lower - Middle contact ("LMC"). The stratigraphy underlying the property belongs the Proterozoic Aldridge Formation, specifically, the lower Aldridge Formation and, therefore, covers prospective strata in the footwall of the Sullivan Horizon.

A total of 41 man days were spent on field explorations by Dr W Lieth during 2012. During his field investigations Dr Leith constructed nine hand-trenches and test-pits (approximately 1-2ft deep) from which to collect prospective rock samples from excavated out crop and sub-cropping rocks.

10 samples were selected by Dr Leith and Mr Wallach for analysis. These samples were dispatched to Acme Analytical Laboratories (Vancouver) Ltd. all samples were analyzed by Aqua Regia digestion Ultratrace ICP-MS analysis for standard Full Suite of 53 elements.

Minor anomalous copper was uncovered in trench 1 and trench 3, these anomalous values continue, an as yet, poorly-defined trend of weak mineralization within samples collected from Dr Lieths hand trenching over the previous years.



Figure 1: Property location map, South Eastern British Columbia, BC.

The Wilcum property cover prospective stratigraphy correlated to the Lower Aldridge Formation in the footwall of the Sullivan Horizon. Historically, the LMC has been the preferred horizon of interest for exploration; however, the Sullivan is, to date, unique among SEDEX deposits in that it is the only economic SEDEX occurrence identified to date in the Aldridge Formation. SEDEX deposits elsewhere in the world occur in clusters throughout the host stratigraphy.




Figure 2: property location map and claim outlines, West of Kimberly, BC, Canada.

The LMC is the primary horizon of interest, sulphides (predominantly pyrrhotite) are however ubiquitous in the Aldridge Formation and are abundant in the Lower Aldridge Formation. Furthermore, smaller massive sulphide bodies have been identified at a variety of levels within the Aldridge Formation, including within, and adjacent to the Wilcum property. Some of these occurrences have been interpreted as sedimentary exhalative in origin, including the Kootenay King mine (Past Producer). In addition, base metal massive sulphide veins or Proterozoic age have been identified, including those within the Northstar Corridor (Stemwinder, Northstar, Quantrelle). Proterozoic age base metal mineralization occurrences include the St. Eugene Mine (Past Producer), the Vine Vein and the Fors occurrence. These occurrences are interpreted to confirm base metal sources active in the Proterozoic during deposition of the Aldridge Formation and may include sedimentary exhalative style mineralization.

2. LOCATION AND ACCESS

The Wilcum property is located in the Purcell Mountains, approximately 28 kilometers west northwest of Cranbrook, British Columbia (Fig. 1 and 2). The claims comprising the property (Fig.3) are located in the Fort Steele Mining Division and extend west from Hell roaring Creek south of St. Mary's River and between Matthew Creek and Resort Creek north of St. Mary's River, centered at approximate UTM coordinates 562900 E, 5494900 N (Latitude 49° 36' 16", Longitude 116° 08'05"). The nearest major centre is the city of Cranbrook, from which exploration programs can be easily supported and supplied. The applicable 1:20,000 TRIM (Terrain Resource and Inventory Management) maps are 082F 060 and 070.

Vehicular access to the property is available from the main St. Mary haul road, south of St. Mary's River, and the St. Mary Lake road north of St. Mary River, to the central portion of the property.

Logging roads along Hell roaring Creek and Angus Creek provide access to the southwest portion of the property; Matthew Creek provides access to the northeast portion of the property. Additional access along Sawmill Creek (a tributary of Perry Creek) provides limited access to the southeast portion of the property. It may prove feasible to rehabilitate old logging and exploration trails from Sawmill Creek and along Pudding and Bannock creeks to gain access to the south central portion of the property. Helicopter support is also available from Cranbrook.

3. PHYSIOGRAPHY AND CLIMATE

The coniferous forest consists predominantly of pine, fir and larch which have been actively logged over the past 30 years. A number of clear-cuts are present throughout the property in various stages of regeneration.

Relief on the property is generally moderate at lower to middle elevation areas, with high relief areas at upper elevations (Fig. 3). Elevation ranges from approximately 950 m along St. Mary River to 2280 m at the northwest corner of the property. Due to the location of the property within the core of the Purcell Mountains east of Kootenay Lake, the area is generally subject to moderately heavy accumulations of snow during the winter months. As a result, the property is available for exploration from mid-May to late October. However, 4WD vehicle supported diamond drilling can take place later into the year despite snow due the relatively extensive and reasonably well maintained network of logging roads.

4. MINERAL TENURE AND CLAIM STATUS

The Wilcum property consists of 7 mineral tenures (Table 1, Fig. 3). All claim information was verified using the BC Government’s Mineral Titles Online (MTO) website and is current as of the compilation of this report. These mineral tenures are currently held in 50/50 ownership between Mr D Wallach (50%) and Dr W Leith (50%).




Figure 3: Local property location map, West of Kimberly, BC, Canada. Showing: claim boundaries, mineral occurrences, and primary drainages.

OWNERS: **201966** LEITH, WILLIAM CUMMING 50.0%

146010 WALLACH, DAVID ARNOLD 50.0%

The property encompasses a total area of approximately 3,620.181 ha (8945.662 acres). The tenures are located immediately north and east of St. Mary Lake and straddle the St. Mary River, approximately 30km west of Kimberley BC.

Table 1: Summary list of the Wilcum claim group, Kimberley area, BC. Source: Mineral Titles Online (MTO).

Tenure Number	Claim Name	Owner	Tenure Type	Tenure Sub Type	Map Number	Issue Date	Good To Date	Status	Area (ha)
522473	WILCUM2	146010 (50%)	Mineral	Claim	082F	2005/nov/21	2014/Sep/07	GOOD	502.267
522474	WILCUM3	146010 (50%)	Mineral	Claim	082F	2005/nov/21	2014/Sep/07	GOOD	523.013
522475	WILCUM4	146010 (50%)	Mineral	Claim	082F	2005/nov/21	2014/Sep/07	GOOD	522.826
522476	WILCUM5	146010 (50%)	Mineral	Claim	082F	2005/nov/21	2014/Sep/07	GOOD	502.148
777142	WILCOM 1	146010 (50%)	Mineral	Claim	082F	2010/may/21	2014/Sep/07	GOOD	523.175
777222	WILCUM 6	146010 (50%)	Mineral	Claim	082F	2010/may/21	2014/Sep/07	GOOD	523.345
777262	WILCUM 7	146010 (50%)	Mineral	Claim	082F	2010/may/21	2014/Sep/07	GOOD	523.407

5. EXPLORATION HISTORY

The following summary has been taken (verbatim) from Kennedy and Anderson (1999), and Walker (2009), and has been adapted in the latter, by the author to reflect the most recent exploration conducted on and immediately adjacent to the current claim block.

Prior to 1979 Cominco Ltd. recognized the presence of a large fragmental located just west of St. Mary Lake which is at equivalent stratigraphy to the Sullivan orebody. The fragmental is exposed at lower elevations on both sides of the St. Mary valley. Most of Cominco's early work on this exploration target has not been reported on in publicly available assessment reports.

- **1979** – Cominco Ltd. (AR 7676) reported on diamond drill hole C79-1 which tested the Clair fragmental on the south side of the St. Mary valley.
- **1981** – Cominco Ltd. (AR 10,311) reported on diamond drill hole C81-1 which tested the Clair fragmental on the north side of the St. Mary valley.
- **1984** – BP-Selco (Assessment Reports 13,108 & 13,871) conducted a program of soil sampling and geological mapping on the Pinetree claims as a follow-up program to a tin anomaly detected by regional stream sampling in 1983. The soil survey outlined a 500m by 100m tin in soil anomaly roughly coincident with the eastern extrapolation of the later discovered stratiform zinc Horn horizon. Follow-up detailed sampling showed that some pegmatite dikes in the survey area are anomalous in tin.
- **1985** – Cominco Ltd. (AR 13,828) conducted a UTEM survey over the Chair fragmental and the strike extent of the lower-middle Aldridge contact on the south side of the St. Mary River.
- **1987** – Esso Minerals Ltd. (AR 16,971) worked on the MAC claims which covered St. Mary Lake and small areas to the north, south and west. Their work included geological mapping over the Clair fragmental which occurs at the same stratigraphy as the Sullivan orebody. They collected and analyzed 32 stream sediment samples, 2 heavy mineral samples and 40 rock samples. One sample from a gabbro boulder with quartz veining returned .083 oz/ton Au, 0.1 oz/ton Ag, 8823 ppm,Cu and 159 ppm Zn.
- **1991** – The Darlin claims were staked based on a prospecting discovery of zinc-mineralized float along the lower Angus Creek Forestry Road. Chapleau Resources Ltd. and Barkhor Resources Ltd. Optioned the property and in turn optioned it to Kokanee Explorations Ltd. who conducted a work program of line cutting, soil geochemistry, surface geophysics and 3025.6 meters of diamond drilling in 13 holes, D91-1 to D91-12 and D91-14. Some of the drill holes were tested by a borehole UTEM survey. The soil geochemistry and surface and downhole geophysical surveys have not been publicly reported on although the writer has reviewed the data in company files. Assessment report 22,799 (Pighin, 1992) reports only on diamond drill holes D91-1,2,3 and 6. No Report was filed on the remaining 9 drill holes. Kokanee Explorations Ltd staked the Horn claims; a large block surrounding the Darlin property and extending for about 8 km to the west, to cover previously identified soil geochem anomalies they were aware of.
- **1992** – Minnova Inc started their evaluation of the Horn property with a program of 61 km of line cutting, geological mapping, prospecting, 521 soil samples, 363 rock sample analyses, 49 core sample analyses, 12 km of ground Pulse EM geophysics, 22 km of ground VLF-EM / Mag

geophysics and 2085.1 meters of diamond drilling in 6 holes, H92-1 to H92-6. Five of these holes, H92-1 to 5 were drilled on or very close to the present Burn claims and were reported on by Burge,

- **1993** – (Assessment Report 22,799) Prospecting located a bedrock exposure of the stratiform zinc mineralization which is now referred to as the Horn horizon. Two of the drill holes successfully tested the Horn horizon, while other holes tested geologic and geophysical targets. One hole, H92-4, the easternmost hole drilled, intersected approximately 100 m of strong albite-chlorite-pyrite alteration. The alteration zone may be related to a fault structure but its similarity to hangingwall alteration at the Sullivan orebody requires a careful evaluation.
- **1993** – Metall Mining Corporation (formerly Minnova) continued their evaluation of the Horn property with a program of 10 km of line cutting, geological mapping, 188 soil samples, 66 rock sample analyses, 105 core sample analyses, 11 km of ground Pulse EM geophysics and 1438 meters of diamond drilling in 2 holes in the western part of the property. The western two drill holes tested a Pulse EM anomaly and intersected a thick, complex section of the Clair fragmental which was not expected in the drill holes based on surface geology. Fracture zinc and lead mineralization is present in both drill holes. These results showed that the Clair fragmental extended farther south than was previously known and that the fragmental was better mineralized in these drill holes than in Cominco's 1979 drill hole which tested the fragmental on the south side of the St. Mary River. Metall then conducted a down hole EM geophysical survey which detected a strong off hole anomaly.
- **1994** – Consolidated Ramrod Gold Corp. (formerly Kokanee Explorations Ltd.) Deepened diamond drill hole H92-3 which was originally drilled to test a geophysical anomaly but was in position to be deepened to test the Horn horizon zinc mineralization. The hole was deepened from 343.2m to 752.1 m for a total of 408.9 m (Pighin, 1995, AR 23,911) and intersected three narrow zones of zinc mineralization, confirming a significant area extent to the zinc mineralized Horn horizon.
- **1996** – Quest International Resources Ltd. (formerly Consolidated Ramrod Gold Corp.) drill tested the off hole EM anomaly detected by Metall Mining Corp. in 1993 on the western Horn property. The hole was drilled to 745.7 m and intersected a narrower zone of fragmental than in DDH H93-7 and 8, with minor base metal sulfides and graphite in brecciated fault zones and shears.
- **1996** – C, Kenedy on behalf of Sedex Mining Corp (AR24596) conducted prospecting and sampling on the adjacent/overlapping historical PMR claims block, directly to the north of the current Wilcum Claims.
- **1996** – F, Fairclough (AR24295) conducted prospecting and sampling on the adjacent/overlapping historical Horn, and Clair claims block, directly to the west of the current Wilcum Claims. This prospecting targeted extensions to the mineralized Horn horizon.
- **1997** – C Cowie on behalf of MINER RIVER RESOURCES, and EAGLE PLAINS RESOURCES LTD (AR24907) conducted reconnaissance mapping and sampling on the historical and overlapping Bootleg Claim Group to evaluate the property for possible Sullivan-type base metal mineralization. Silt sampling of the main property drainages, and reconnaissance prospecting was undertaken. A total of 22 rock samples and 27 stream sediment samples were collected. Stream sediments yielded anomalous results in northern drainages, whilst three samples of

rusty Aldridge quartzite float taken within 300m of the anomalous drainage returned high metal values. RBBLR-01 returned a value of 17.0g/t Ag and 3096ppm Pb, RBBLR-02 returned a value of 1020ppm As, and RBBLR-03 had geochemical values of 220ppm Ba and 1024ppm Zn.

- **2008** – Dynamic Exploration Ltd conducted field explorations on a historical grouping of the Wilcum Claims which overlap the current claims, and collected 122 soil samples from along the main St. Mary Haul Road through the approximate centre of the property. Samples were collected from a variably developed “B Horizon”, with many of the samples taken from the top of road cut exposures. Sample depths ranged from 5 cm to 50 cm. Results returned from this preliminary soil program were disappointingly low for precious and base metals, further work was recommended.

6. TERMS OF REFERENCE

At no point has the author ever visited the Wilcum High Peak property, nor has any firsthand knowledge of the property geology or mineral occurrences that occur upon it.

This assessment report has been prepared by the author using documents and information provided by the Owner-Operator for this purpose. While reasonable care has been taken in the preparation of this report, the Author cannot guarantee the accuracy or completeness of all supporting documentation.

Dr W Leith (FMC 201966) was solely responsible, and himself conducted all field prospecting and hand trenching at his own discretion during 2012. Dr W Leith with the assistance of co-owner and operator Mr D Wallach selected and prepared samples for analysis.

7. REGIONAL GEOLOGY

Proterozoic Purcell Supergroup Stratigraphy

The Proterozoic Purcell Supergroup, host of the Sullivan Mine, constitutes a thick prism of dominantly clastic sediments exceeding 10,000 metres in thickness, with the base unexposed. The oldest exposed strata is represented by the Fort Steele Formation fluvial/deltaic sequences of quartz arenite, quartz wacke and mudstone, comprising a sequence at least 200 metres thick. Fine-grained clastic beds at the top of the Fort Steele formation grade into very rusty-weathering, fine-grained quartz wacke and mudstone of the Aldridge Formation (1433 Ma +/- 10 Ma), at least 5000 metres thick in the Purcell Mountains. The Aldridge Formation grades upward over 300 metres through a sequence of carbonaceous mudstone with minor beds of grey and green mudstone and fine-grained quartz wacke to the 1800 metre thick Creston Formation, composed of grey, green and maroon quartz wacke and mudstone with minor white arenite.

The following has been taken from Höy (1993), as compile by Walker (2009):

Aldridge Formation

“Within the Purcell Mountains, it has been subdivided into three main divisions: the lower Aldridge comprises rusty weathering siltstone, quartz wacke and argillite; the middle Aldridge, grey weathering quartz wacke and siltstone interbedded with silty argillite; and the upper Aldridge, rusty to dark weathering laminated argillite and silty argillite .

Middle Aldridge

The middle Aldridge comprises more than 2000 metres of dominantly well-bedded, medium to locally coarse-grained quartz arenite, wacke and siltstone.

A continuous section ... is not exposed in the Purcell Mountains; the most complete section, between the Moyie and Cranbrook faults, is broken by a number of faults. In general, the basal part comprises interbedded quartz wacke and arenite with only minor sections of silty argillite. Exposures of the basal part are typically grey weathering; however, in recent manmade exposures ... these units are typically rusty weathering. Within the upper part of the middle Aldridge, quartz arenite and quartz wacke beds become thinner and less pure, and the proportion of bedded siltstone and argillite increases. The upper part of the middle Aldridge comprises a number of distinct cycles of massive, grey quartz arenite beds that grade upward into an interlayered sequence of quartz wacke, siltstone and argillite, and are capped by siltstone and argillite. The contact with the upper Aldridge is placed above the last bed of massive grey quartz arenite.

Laminated Siltstone markers

The marker units are sequences of laminated dark, and siltstone, up to several metres thick, in which each laminae can be matched in precise detail for distances up to several hundred kilometres. The pattern of each laminae is each sequence in unique and hence recognition of a specific sequence of laminae allows accurate positioning of isolated outcrops or drill intersections within the thick middle Aldridge succession. At least fourteen of these marker sequences are recognized. Locally, the markers are interrupted by turbidity deposits, or partly or totally removed due to erosion by turbidity currents. ...

Upper Aldridge

The upper Aldridge Formation comprises about 500 ... metres of dominantly medium to dark grey siltstone, argillaceous siltstone and argillite. It is generally rusty weathering, thin bedded and thinly laminated. Thin graded ailtite-argillite couplets and lenticular bedding with tan siltstone lenses in argillite are common bed-forms; syneresis cracks are commonly observed near the top of the upper Aldridge.

The contact of the upper Aldridge with the Creston Formation is relatively abrupt, and is placed where green tinged siltite layers first appear. Elsewhere, a massive, thick-bedded siltstone or wacke marks the base of the Creston Formation.

Intrusives

The following has been taken from Höy (1993), as compile by Walker (2009):

Proterozoic Moyie Sills

The Moyie Sills (or Intrusives) comprise laterally extensive gabbro (to dioritic) sills which are restricted to the lower Aldridge and the lower part of the middle in the Purcell Mountains. The sills comprise up to 30 percent of the lower to middle Aldridge stratigraphic succession, having an aggregate thickness in excess of 2000 metres, with the abundance decreasing upwards relative to the abundance of thick-bedded A-E turbidites.

In the Lamb Creek area west of Moyie Lake, an aggregate thickness of approximately 1300 metres of sills is interlayered with 2800 metres of lower and middle Aldridge sedimentary rock.

Moyie sills form an extensive suite of basaltic rocks that intruded lower and middle Aldridge turbidites and siltstones. Although it has been proposed that Moyie sills are coeval with deposition of upper Aldridge or Creston rocks, or perhaps with the Nicol Creek lavas, contact relationships between sills and Aldridge rocks indicate that some sills were extruded at very shallow depths in unconsolidated, water-saturated sediments. Others with fine-grained chilled margins have contact metamorphosed the country rocks. As these sills are interpreted to be part of a continuous magmatic event, they record an igneous/thermal event of regional extent during deposition of lower and middle Aldridge rocks. Hence, a Middle Proterozoic uranium-lead date of 1445 Ma from zircons in the Lumberton sill west of Cranbrook defines the minimum age of deposition of lower and basal middle Aldridge.

Structure

The following has been summarized from Höy (1993), as compile by Walker (2009):

Rocks of the Purcell Supergroup have been affected by several separate phases of deformation, ranging from Middle Proterozoic through to Paleocene. The North American craton underwent two phases of extension, a compressional orogeny and subsequent continental rifting, followed by development of a miogeocline. Thrusting and folding associated with development of the Foreland Fold and Thrust belt took place from Cretaceous to Paleocene time and was followed by Eocene extension.

The earliest deformation was associated with extension in the Middle Proterozoic which resulted in block faulting along the margin of the Purcell Basin, coincident with deposition of the Fort Steele and Aldridge formations. Movement along growth faults is interpreted to have ceased by upper middle to upper Aldridge time.

A late Middle to early Upper Proterozoic (1300 to 1350 Ma) compressional event, the East Kootenay orogeny, has been interpreted based upon evidence for deformation and metamorphism prior to deposition of lower Paleozoic miogeoclinal strata. This event was associated with folding, development of a regional cleavage and granitic intrusions (i.e. 1305 ± 52 Ma Hellroaring Creek stock). Localized high grade metamorphic areas (i.e. Mathew Creek) are related to this tectonic event which is interpreted to have terminated Belt Purcell sedimentation.

The extensional Goat River orogeny occurred during deposition of the Windermere Supergroup (800 to 900 Ma) and is characterized by large-scale block faulting during and perhaps immediately prior to

deposition of strata. The Windermere Supergroup is comprised of a basal conglomerate (Toby Formation) overlain by immature clastic and carbonate sediments of the Horsethief Creek Group.

The Toby Formation consists of "... predominantly conglomerates and breccias, interpreted to have been deposited in fan sequences adjacent to active fault scarps in large structural basins. Locally, up to 2000 metres of underlying Belt-Purcell rocks have been eroded from uplifted blocks, providing a sediment source ... in adjacent basins" (Höy 1993).

The earlier tectonic events may record incipient rifting, with development of block-faulted, intracratonic structural basins, whereas by early Paleozoic time continental separation had occurred as platformal and miogeoclinal sediments were deposited on a western continental margin. The Laramide orogeny (Late Jurassic to Paleocene) resulted in the horizontal, northeast directed compression of Proterozoic strata and the overlying Paleozoic miogeoclinal prism onto the North American craton. Easterly verging thrust faults and folds developed with normal faults and westerly verging back thrusts and normal faults, resulting in a complex structural pattern. Two major faults, St. Mary and Moyie faults, have had a significant role in the structural history and fabric of the region, controlling facies and thickness changes in Proterozoic and Paleozoic strata.

A final episode of north-trending, west-dipping normal faulting took place in the Late Tertiary. The Rocky Mountain Trench is the most prominent and is a listric normal fault having dip-slip separation of at least 5 to 10 kilometres. However, strike slip separation is interpreted to be minimal based on stratigraphic correlations across the trench.

Mineralization

The following has been summarized from Höy (1993), as compile by Walker (2009):

Sullivan Mine Type

The Sullivan orebody is a conformable iron-lead-zinc sulphide lens enclosed by metasedimentary rocks of the Middle Proterozoic (Helikian) Aldridge Formation, the basal formation of the Purcell Supergroup. The regional metamorphic grade is upper greenschist facies.

The orebody occurs near the top of the Lower Aldridge Formation, at the transition from the Lower to Middle Aldridge Formation ("Lower - Middle Contact or LMC"), and has the shape of an inverted and tilted saucer. The maximum dimensions are approximately 2000 metres north-south and 1600 metres east-west. In general, bedding dips flat to gently east in the western portion of the orebody, moderately east to northeast in the central portion, and gently east to northeast in the eastern portion.

The orebody attains a maximum thickness of 100 metres approximately 100 metres northwest of its geographic centre, and thins outward in all directions (averaging 21 metres in thickness). To the east, it thins gradually to a sequence of pyrrhotite-laminated mudstone 3 to 5 metres thick that persists laterally for some distance. To the north, the orebody thins less gradually and is truncated by the Kimberley fault. To the west, the orebody thins abruptly and is cut by dyke-like apophyses of the foot-wall gabbro. The gabbro (correlated to the Middle Proterozoic Moyie Intrusions) lies beneath the orebody and is typically concordant about 500 metres below its eastern edge. To the west, the gabbro

rapidly cuts up-section through the host stratigraphy transgresses to the footwall of the orebody near its western margin. Farther west, the gabbro cuts back down-section to approximately its original stratigraphic position. To the south, within the limit of economic mineralization, thickness changes are generally irregular and abrupt.

The Kimberley fault dips 45° to 55° north and truncates the ore zone to the north. The fault juxtaposes strata, the fault juxtaposes strata of the Creston and Kitchener formations against those of the Lower Aldridge and is, therefore, interpreted to have over 3000 metres of stratigraphic displacement. "Sullivan-type" faults cut the orebody with a consistent west side down, normal displacement ranging from a few metres to 30 metres displacement. The Sullivan orebody consists of sulphide-rich rock, with more than 70 per cent sulphide content, in thick, gently dipping conformable units enclosed by interbedded, unaltered to altered quartz wacke and mudstone. In the western portion of the orebody, massive pyrrhotite containing occasional wispy layers of galena is overlain by sulphide-rich rock in which conformable layering consists of pyrrhotite, sphalerite, galena and pyrite are intercalated with beds of clastic sedimentary rock. The ore passes outward on the north, east and south to delicately-bedded sulphide-bearing rock interbedded with fine-grained clastic sedimentary rocks. Eastward across a transition zone, the orebody is composed of five distinct conformable units of well-bedded sulphide-rich rock interbedded with clastic sedimentary rock. Each sulphide bed thins eastward from the transition zone. The transition zone is commonly only a few metres or tens of metres wide. Three bedded sulphide sequences occur above the main orebody, particularly in the area of the transition zone. Locally, these constitute ore. Much of the orebody is underlain by locally derived intraformational conglomerate, which is more than 80 metres thick in the west and thins to the east. Footwall rocks are cut by tabular bodies of chaotic breccia containing blocks of conglomerate and bedded sedimentary rock; these extend downward unknown distances from the sulphide footwall in the west. Footwall mineralization consisting of thin conformable laminae, veins and locally intense fracture-filling is common in the west and very rare in the east.

Pyrrhotite and pyrite (average ratio of 7:3) are the most abundant sulphides in the Sullivan orebody. Galena and sphalerite are the principal ore minerals. Minor but economically important minerals include tetrahedrite, pyrargyrite, boulangerite and arsenopyrite. Cassiterite is an important minor constituent in the western part of the orebody. Minerals constituting less than 1 per cent include chalcopyrite, jamesonite, magnetite and less abundant scheelite and stannite. Principal non-sulphide minerals are quartz and calcite with abundant tourmaline, chlorite, muscovite, albite, pale brown to reddish-brown mica, garnet, tremolite, epidote, actinolite, cordierite and hornblende. Either quartz or calcite may make up 50 to 70 per cent of the non-sulphide suite, chlorite 30 per cent and the other minerals up to about 20 per cent.

8. LOCAL GEOLOGY

The structure of the area is dominated by the Purcell Anticlinorium, a broad anticlinal structure which exposes strata of the Purcell Supergroup. The western limb of the anticlinorium is host to several regionally significant faults, having considerable east side down, dip-slip displacement and resulting in duplication of the Purcell Supergroup strata. The property is influenced by the major northeast trending St. Marie River Fault approximately 2 km to the south. The St. Marie Fault juxtaposes the strata of the lower Aldridge Formation against strata of the Cambrian Eager Formation along Hellroaring Creek and, therefore, has considerable dip-slip and stratigraphic displacement, representing in excess of 4.6 km of vertical displacement (Brown 1998, Höy 1993). The Aldridge Formation in the hangingwall in the area of the Wilcum property is comprised predominantly of the lower Aldridge Formation, with subordinate exposures of the middle

Aldridge Formation.

The Sullivan Mine (and Sullivan Horizon) is located at the Lower - Middle Contact ("LMC"). This stratigraphic contact has been mapped westward from the mine (Høy and Jackaman 2004) and passes immediately northwest of the Wilcum property, placing the strata underlying the property in the footwall of the LMC, in the lower Aldridge Formation. The strata of the lower Aldridge have been intruded by a considerable cumulative thickness of tholeiitic basalt, comprising the Moyie Intrusive Suite, only slightly subordinate to the clastic lithologies of the sedimentary host strata. Based on a review of isopach maps, Lydon et al. (2008) proposed that the source vent the prolific and volumetrically extensive Moyie Intrusive Suite was located in the general vicinity of the St. Mary drainage. As such, the proposed source vent may have provided a conduit for basic, magmatic fluids from the mantle resulting in precipitation of the Sullivan mine and the peripheral, Proterozoic base metal occurrences North Star Corridor.

9. PROPERTY GEOLOGY

No singular property geology, or deposit geology narrative exists nor is there focus on a particular mineral showing, a number of which occur within the current claims. In the absents of this, the combined works of Hoy (1993), Kennedy and Anderson (1999), and various other authors go to form the understanding of the property geology and mineral prospectivity for the new claim grouping; the Wilcum High Peak project area as it is known in the context of this report.

The following has been taken from Kennedy and Anderson (1999), as compiled by Walker (2009). This text best summarizes local property scale features and may serve as a prospecting guide. Furthermore the reader is referred to figure 4 in Walker (2009) for the most up-to-date compiled map representation of the property scale geology.

Previous exploration programs have confirmed the existence of exhalative, iron-rich, base metal horizons in stratigraphy mapped as Lower Aldridge sediments. Though these conformable zones are narrow (widest intercept 0.10 m) they have been traced successfully by soil geochemistry and short hole

drilling. These strataform lenses have a known strike length in excess of 2 km. The area prospected straddles a steep northwest striking syncline which has been delineated by previous geological mapping.

The major prospecting difficulty encountered on the (Wilcum property). is the lack of sustained outcrop. This factor is very evident on the northwestern part of the property (south of the St. Mary River). Here most outcrops encountered are gabbros dominating topographic highs. Topographic highs are sporadic which makes gabbro correlation a best guess effort.

Granophyre

Possibly the most intriguing discovery; was that of two confirmed occurrences of granophyre. Granophyre is recognized within the gabbro complex (mine sill) which underlies the Sullivan orebody, 14 km to the northeast. Both outcrops prospected had zones in which clasts were noted, these clasts, though altered and similar in makeup to the matrix, do show faint but distinguishable boundaries. The granophyre is mostly made of quartz, biotite, and feldspar with varying amounts of iron sulphide. The outcrop ... has a number of very iron-rich zones in which both pyrrhotite and pyrite are abundant, chalcopyrite is also quite common. Of most potential significance is that this outcrop has abundant disseminations of arsenopyrite in the more iron-rich zones. Another positive feature is the existence of quartz vein float containing massive sulphide lenses. The massive sulphide is pyrrhotite, chalcopyrite along with minor amounts of arsenopyrite. Where these veins were encountered in outcrop they were striking in a northwest direction, similar to weak but prevalent shearing noted within the package. The granophyre in this area seems to be structurally controlled as it pinches out to the east while thickening to the west before being lost in the overburden. A poorly exposed outcrop of Lower Aldridge phyllitic siltstone and muscovite schist, with minor fine-grained quartzite seems to be in contact with the granophyre hangingwall. An actual contact was not seen owing to the poor exposure. One narrow quartzite package within this sedimentary sequence was obviously more iron-rich. The abundant pyrrhotite section was accompanied by increased silicification, sericite, albite and chlorite alteration. Some rare disseminations of zinc mineralization were noted within this alteration. Again, due to poor exposure this horizon could not be traced along-strike east or west.

The granophyre encountered on Route 4 has a 5 m wide pegmatite sill on its hangingwall with thin bedded phyllitic siltstones and muscovite schists on its footwall. The granophyre is in excess of 20 m wide and contains abundant blebs and clusters of pyrrhotite. The more iron-rich zones contain disseminations of fine grained chalcopyrite, no arsenopyrite was noted. One hand sample from the outcrop did contain a small amount of lime green pyromorphite crystals in a quartz-rich vug. The pegmatite hosts rare blebs of galena in zones with yellow pyromorphite crust, this mineralization occurs close to the pegmatite granophyre contact.

Iron-rich Bedrock Horizon

Northeast of the granophyre on Route 4 you contact a quartzite dominated sequence of silicified pyrrhotite-rich rocks. This package overlies thin bedded phyllites and mica schists. These thin bedded rocks are rusty, sheared and slightly folded. Narrow chloritic limonite-rich quartz veins are commonly associated with more iron-rich shear plains within this stratigraphy. The pyrrhotite-rich quartz sequence is hosted within a 50 m thick sediment panel. Individual iron-rich quartzite beds reach widths in excess

of 2 m with the majority 0.5 m or less. Pyrrhotite content within these beds exceeds 10% by volume. Rare chalcopyrite and arsenopyrite were the only other sulphides noted with the pyrrhotite. This sedimentary horizon may represent the contact between Middle and Lower Aldridge stratigraphy. These rocks strike northwest and dip moderately to the northeast. They occupy the western flank of the northerly trending synform. A further support for Middle-Lower Aldridge contact is that these rocks overlay the granophyre complex, this is a similar situation to what occurs at the Sullivan Corridor.

Sulphide-rich Sedimentary Float (minor base metals)

Two types of sedimentary float containing pyrrhotite and base metal mineralization was seen during the prospecting program.

- The first type of float is similar to that found in outcrop, silicified fine grain quartzite with chlorite, sericite, albite and moderate to heavy disseminations of pyrrhotite. Some of the iron is obviously banded.
- The other type of float is coarse grained quartz sand with actinolite, sericite and chlorite. The pyrrhotite is in blebs and clusters with thin threads of iron-pyrite. Minor amounts of sphalerite and galena are found along fractures or more rarely with the pyrrhotite.

Intrusives

Gabbro is the most abundant intrusive encountered on the property and also is the most dominant rock type on all traverse routes. Most gabbro is of standard type, dark green, fine to coarse grained, predominant hornblende, plagioclase with accessory mafic minerals. Of most interest are the iron-rich gabbros which have pyrrhotite with minor amounts of iron-pyrite and chalcopyrite. These unusual rusty outcrops are often found in association with granophyres or adjacent to structures.

Two float occurrences of abundant pieces of lamprophyre were seen along Route 2. The northern occurrence was lamprophyre made up of coarse matrix material, having clusters of ragged green mica. The other occurrence south on Route 2 is a more standard type lamprophyre, a very micaceous fine to medium grain rock. Samples tested from both locations proved non-magnetic.

Next to the gabbros, pegmatite sills and/or dykes are the most prevalent intrusive encountered on the property. Most sills or dykes are quite thin, usually less than 2 m in thickness. As mentioned previous, the largest pegmatite encountered was a sill in the granophyre complex, this sill is in excess of 5 m. The pegmatites consist of very coarse feldspar, quartz, muscovite and occasionally minor tourmaline. The 5 m wide sill had the only base metal mineralization noted. All pegmatites had minor amounts of disseminated limonite or pyrite. Some of the mineralization occurred with very discontinuous narrow quartz veining.

10. 2012 WORK PROGRAM

A total of 41 man days were spent on field explorations by Dr W Lieth during 2012. During his field investigations Dr Leith constructed nine hand-trenches and test-pits (approximately 1-2ft deep) from which to collect prospective rock samples from excavated out crop and sub-cropping rocks. Dr Lieth also carried out reconnaissance field prospecting on the claim block adjacent to, and to help better direct his excavations.

Numerous undescribed rock samples were collected from nine pits located adjacent to the main haul road that transects the property and at the base of out cropping and sub cropping rocks of the underlying Aldridge Formation. Sample locations are illustrated in appendix 3 and were recorded using hand-held GPS which are regarded as being accurate to within 10 m.

10 samples were selected by Dr Leith and Mr Wallach for analysis. These samples were numbered and dispatched to Acme Analytical Laboratories (Vancouver) Ltd. all samples were analyzed by Aqua Regia digestion Ultratrace ICP-MS analysis for standard Full Suite of 53 elements.

11. RESULTS

The results for the rock sample analysis are summarized in the table below:

Table 2: Summarized results for the 2012 rock sampling program.

Sample ID	EAST	NORTH	ZONE	DESCRIPTION	DATE SAMPLED	ROCK DESCRIPTION	Long-Lat	Cu PPM	Pb PPM	Zn PPM	Ag PPB
2161301	560883.3	5496317	Z11N	Site 1	24-Feb-2013	n/a	49.6163 N - 116.157 W	160.53	2.28	38.1	52
2161302	561039.8	5496440	Z11N	Site 2	16-Mar-2013	n/a	49.6174 N - 116.155 W	84	3.35	33.6	28
2161303	561194.5	5496540	Z11N	Site 3a	26-Apr-2013	Cu & maybe Ge, In, Ga	49.6183 N - 116.153 W	6.44	2.95	11.4	13
2161304	561359.5	5496628	Z11N	Site 3b	26-Apr-2014	Cu & maybe Ge, In, Ga	49.6191 N - 116.151 W	127.62	1.68	39.7	87
2161305	561516	5496751	Z11N	Site 4	8-May-2013	n/a	49.6202 N - 116.148 W	2.76	4.12	23.9	11
2161306	561682	5496851	Z11N	Site 5	14-Jun-2013	may be kimberlite	49.6211 N - 116.146 W	2.65	4.42	51.3	10
2161307	561869.4	5496937	Z11N	Site 6	12-Jul-2013	may be kimberlite	49.6218 N - 116.143 W	2.95	10.42	12.6	24
2161308	562042	5496980	Z11N	Site 7	22-Aug-2013	n/a	49.6222 N - 116.141 W	2.72	2.45	5.9	9
2161309	562237.1	5497020	Z11N	Site 8	1-Sep-2013	n/a	49.6225 N - 116.138 W	2.36	3.44	22.5	11
2161310	561194.5	5496540	Z11N	Site 9	7-Oct-2013	n/a	49.6183 N - 116.153 W	2.04	3.15	5.9	5

12. CONCLUSIONS

The first priority for further exploration at the property should be to expand Dr Leith's excavations around previously anomalous trenching to confirm the presence of any possible massive sulphide Cu, Au, Ag bearing rocks. If it is possible to establish a trend to the mineralization discovered this should be the second priority for ongoing work on the property.

Furthermore Walker (2009) has summarized most appropriately that 'previous programs within and immediately adjacent to, the property have reported the presence of massive sulphide lenses, both at surface and in drill holes'. These represent key features of interest for future evaluation of the property for Sedex potential.

Ongoing explorations are required to further develop this property; ultimately drill targets need to be established to affect the potential discovery of the Sedex type deposit on the Wilcum High Peak property.

To facilitate drill targeting there are a range of conventional exploration techniques that can be utilized to develop targets. Beyond the scope of the work currently being conducted by the owners, there is a demonstrated model for making successful Sedex Type discoveries; Exploration methods to be considered, compiled, and continued should be stream, soil and rock geochemical profiling, detailed where necessary to reflect prospective areas of interest according to additional and updated compilation and mapping of the property geology, structure, alteration and mineralization. Geophysical exploration techniques are also highly recommended to facilitate exploration under the cover which blankets significant portions of the property, and otherwise hampers field explorations.

It is the author's opinion that many of the previous recommendations made by historical authors are still valid and apply to the project in its current context; these recommendations should be coalesced into a unified exploration strategy to continually develop an ongoing exploration program for the Wilcum High Peak property.

13. RECOMENDATIONS

- Conduct a complete and comprehensive compilation of all historic data that has been collected though out the project history.
 - The data review should include but not be limited to verification of any geophysics, compilation of a geochemical database for rock, soil, and stream samples, and compilation of all meaningful historical mapping.
 - Additionally compilation and plotting of drill hole data and analytical results should be made so as to tie them in to surface data to extend control for projection of massive sulphide horizons.
 - All data should be compiled into a GIS database to facilitate and direct future project explorations in the area.
- Complete 'Ground truthing' and resampling of all previously identified MINFILE occurrences and other significant showing located by previous authors.
- Comprehensive and where appropriate, detailed geological mapping should be conducted.
 - Additional lithological/stratigraphic ground truthing should be conducted over the entire claim block in an attempt to confirm and correlate the location of mineralized occurrences and local lithologies of interest. This could include geochemical profiling of potential Sedex host lithologies.
 - Walker (2009) recommends mapping the distribution of Moyie Intrusive exposures so as to provide stratigraphic and/or structural control for projection of massive sulphide horizons across the property.
 - Other prospective exploration targets in the search for sedex deposits are fault-bounded sub-basins, since hydrothermal exhalations were controlled by fluid movement along these faults. Synsedimentary faults can be identified by the presence of synsedimentary fault breccias, which are composed of sedimentary fragments cemented by more sedimentary material.
- Stream silt samples could also be collected from along the creeks draining the property in an attempt to identify those having anomalous geochemistry.
 - This could be facilitated by the use of portable XRF to help rapidly, and cost effectively, deduce the presence of soil geochemistry anomalies.
- Additional and comprehensive Soil geochemical samples should be collected from a-b soil horizons along traverse lines that transect local stratigraphy, and structures and surrounding mineral showing where appropriate.
 - This could be facilitated by the use of portable XRF to help rapidly, and cost effectively, deduce the presence of soil geochemistry anomalies.
 - Trenching could be utilized to further develop larger soil geochemistry anomalies and perhaps further to develop potential drilling targets.
- Geophysical Exploration should also be considered; the sulphide horizons are large and considerably more conductive and denser than the host sedimentary rocks. Techniques to be considered should include airborne and ground surveys for magnetic, gravity and electromagnetic properties, as well as ground-based induced-polarization surveys.

14. STATEMENT OF COSTS

Personnel (Name)/ Position	Field Days (list actual days)	Days	Rate	Subtotal*
DR WILIAM C LEITH/ PROSPECTOR (\$47.33ph)	February: 24-28 (48hrs)	5	\$ 47.33ph	\$2,271.84
DR WILIAM C LEITH/ PROSPECTOR (\$47.33ph)	March : 16-19 (44hrs)	4	\$ 47.33ph	\$2,082.52
DR WILIAM C LEITH/ PROSPECTOR (\$47.33ph)	April: 26-30 (45hrs)	5	\$ 47.33ph	\$2,129.85
DR WILIAM C LEITH/ PROSPECTOR (\$47.33ph)	May : 8-11 (52hrs)	4	\$ 47.33ph	\$2,461.16
DR WILIAM C LEITH/ PROSPECTOR (\$47.33ph)	June : 14-18 (51hrs)	5	\$ 47.33ph	\$2,413.83
DR WILIAM C LEITH/ PROSPECTOR (\$47.33ph)	July : 12-16 (47hrs)	5	\$ 47.33ph	\$2,224.51
DR WILIAM C LEITH/ PROSPECTOR (\$47.33ph)	August: 22-25 (42hrs)	4	\$ 47.33ph	\$1,987.86
DR WILIAM C LEITH/ PROSPECTOR (\$47.33ph)	September: 1-5 (53hrs)	5	\$ 47.33ph	\$2508.49
DR WILIAM C LEITH/ PROSPECTOR (\$47.33ph)	October: 7-10 (47hrs)	4	\$ 47.33ph	\$2224.51
429hrs		41.0		\$20,304.57
Office Studies	List Personnel			
Database compilation	LUKE VAN DER MEER/ GEOLOGIST	0.5	\$ 550.00	\$275.00
Report preparation	LUKE VAN DER MEER/ GEOLOGIST	1.0	\$ 550.00	\$550.00
		1.5		\$825.00
Ground Exploration Surveys	Area in Hectares/List Personnel			
Reconnaissance	DR WILIAM C LEITH/ PROSPECTOR	1400	ha	
Geochemical Surveying	Number of Samples	No.	Rate	Subtotal
Rock	laboratory costs for 10 samples	10.0	\$ 60.00	\$600.00
				\$600.00

Transportation		No.	Rate	Subtotal
fuel	fuel costs accrued (vehicle and generator)	2,000	\$ 1.30	\$2,600.00
				\$2,600.00
Accommodation & Food		Rate	Days	Subtotal
Camp	costs accrued	90.00	41.00	\$3,690.00
Meals	costs accrued	70.00	41.00	\$2,870.00
				\$6,560.00
Equipment Rentals		No.	Rate	Subtotal
Field Gear (Specify)	misc cost accrued		\$ -	\$17.79
				\$17.79
Total				\$30,907.36

15. REFERENCES

- Brown, D.A. 1998. Geological Compilation of Grassy Mountain (East Half) and Moyie Lake (West Half) Map Areas, Southeastern British Columbia, British Columbia Ministry of Energy and Mines Geoscience Map 1998-3, 1:50,000 scale map.
- Downie, C.C. 1997. GEOLOGICAL REPORT on the BOOTLEG PROPERM Fort Steele Mining Division, Southeastern British Columbia. Assessment Report 24907 filed March 18, 1997.
- Fairclough, F. 1996. HORN PROPERTIES UNTITLED REPORT, Geological Mapping and Prospecting, Fort Steele Mining Division, B.C., Assessment Report 24295 filed February 14, 1996.
- Goldsmith, B. 1985 Preliminary Geological Mapping, P.C. MINERAL CLAIM GROUP, FORT STEELE MINING DIVISION ARGYLE AND ALKI CREEKS ST. MARY LAKE AREA, B.C. Assessment Report 14423 filed January 20, 1985.
- Höy, T. and Jackaman, W. 2004. Geology of the St. Mary Mapsheet, NTS 82F/09; British Columbia Ministry of Energy, Mines and Petroleum Resources Geoscience Map 2004-01.
- Höy, T. 1993. Geology of the Purcell Supergroup in the Fernie West-Half Map Area, Southeastern British Columbia, British Columbia Ministry of Energy, Mines and Petroleum Resources Bulletin 84, 157 p.
- Kennedy, C. and Anderson, D. 1999. Geological Mapping and Prospecting Assessment Report -Burn Property, Fort Steele Mining Division, B.C., Assessment Report 25,808, filed January, 1999.
- Lydon, J.W, Thomas, M.D. and Currie, M.L. 2008. Ni-Cu-PGE potential of the East Kootenays: a case for optimism, Geological Survey of Canada poster presented at Association for Mineral Exploration British Columbia Roundup.
- Mark, D. G. 1985. GEOPHYSICAL REPORT ON AIRBORNE MAGNETIC AND VLF-EM SURVEYS OVER THE ST. MARY LAKE PXOPERTY ST, MARY LAKE, KIMBERTXY AREA FORT STEELE WINING DIVISION BRITISH COLUMBIA. Assessment Report 13682 filed April 4, 1985.
- Royer, G.A. 1985. GEOLOGICAL/GEOPHYSICAL REPORT ON THE BOOTLEG CLAIM GROUP ST. MARY LAKE AREA, FORT STEELE M.D., BRITISH COLUMBIA. Assessment Report 14358 filed November 27, 1985.
- Walker, R.T. 2009. Assessment Report for the Wilcum Property Soil Sampling, Fort Steele Mining Division, BC. Assessment Report , filed May 18th, 2009.

16. APPENDICIES

APPENDIX 1: Certificate of the Author.

APPENDIX 2: Rock sampling locations and analytical results.

APPENDIX 3: Sample Maps and Diagrams

Sample Location

APPENDIX 1


Certificate of the Author.

I Luke van der Meer of 308-1685 W13th Ave, Vancouver, British Columbia do hereby certify that:

- I am a contracting geologist employed by VDM Geological Consulting at 308-1685 W13th Ave, Vancouver, British Columbia, Canada.
- I graduated from the University of Otago in 2001 with a Bachelor's degree in Science majoring in Geology and Geography.
- I am currently within the application process for Membership to the Professional Association of Engineers and Geoscientists of British Columbia.
- I have worked as a geologist for 10 years since my graduation from university, primarily within exploration and mining in Australia, New Zealand, Mongolia and Canada.
- I have been contracted by DW Exploration to compile an Assessment Report on the 2012 work program at the Wilcum High Peak Property.
- At no time have I conducted a site visit therefore this assessment report has been prepared by the author using documents and information provided by the Owner-Operator for this purpose. While reasonable care has been taken in the preparation of this report, the Author cannot guarantee the accuracy or completeness of all supporting documentation.
- Other than my capacity as a contracting geologist to DW Exploration, I have not received nor do I expect to receive and interest, directly or indirectly in the property described in this report; the Wilcum High Peak Property.

Dated at Vancouver, British Columbia, this 15th of May 2013.

Luke P.A. van der Meer, PGeo, B.Sc-Geology



**APPENDIX 2: Rock
sampling
locations and
analytical results.**

	Sample ID	EAST	NORTH	ZONE	DESCRIPTION	DATE SAMPLED	ROCK DESCRIPTION	Long-Lat
1	2161301	560883.3	5496317	Z11N	Site 1	24-Feb-2013	n/a	49.6163 N -116.157 W
2	2161302	561039.8	5496440	Z11N	Site 2	16-Mar-2013	n/a	49.6174 N -116.155 W
3	2161303	561194.5	5496540	Z11N	Site 3a	26-Apr-2013	Cu & maybe Ge, In, Ga	49.6183 N -116.153 W
4	2161304	561359.5	5496628	Z11N	Site 3b	26-Apr-2014	Cu & maybe Ge, In, Ga	49.6191 N -116.151 W
5	2161305	561516	5496751	Z11N	Site 4	8-May-2013	n/a	49.6202 N -116.148 W
6	2161306	561682	5496851	Z11N	Site 5	14-Jun-2013	may be kimberlite	49.6211 N -116.146 W
7	2161307	561869.4	5496937	Z11N	Site 6	12-Jul-2013	may be kimberlite	49.6218 N -116.143 W
8	2161308	562042	5496980	Z11N	Site 7	22-Aug-2013	n/a	49.6222 N -116.141 W
9	2161309	562237.1	5497020	Z11N	Site 8	1-Sep-2013	n/a	49.6225 N -116.138 W
10	2161310	561194.5	5496540	Z11N	Site 9	7-Oct-2013	n/a	49.6183 N -116.153 W



www.acmelab.com

Acme Analytical Laboratories (Vancouver) Ltd.
9050 Shaughnessy St Vancouver BC V6P 6E5 CANADA
PHONE (604) 253-3158

Client: **DW Exploration**
5241 Cobble Crescent
Kelowna BC V1W 5C3 Canada

Submitted By: Dave Wallach
Receiving Lab: Canada-Vancouver
Received: April 03, 2013
Report Date: April 22, 2013
Page: 1 of 2

CERTIFICATE OF ANALYSIS

VAN13001075.1

CLIENT JOB INFORMATION

Project: Wilcum High Peak
Shipment ID:
P.O. Number
Number of Samples: 10

SAMPLE PREPARATION AND ANALYTICAL PROCEDURES

Method Code	Number of Samples	Code Description	Test Wgt (g)	Report Status	Lab
R200-250	10	Crush, split and pulverize 250 g rock to 200 mesh			VAN
1F06	10	1:1:1 Aqua Regia digestion Ultratrace ICP-MS analysis	30	Completed	VAN

SAMPLE DISPOSAL

RTRN-PLP Return
RTRN-RJT Return

ADDITIONAL COMMENTS

Acme does not accept responsibility for samples left at the laboratory after 90 days without prior written instructions for sample storage or return.

Invoice To: DW Exploration
5241 Cobble Crescent
Kelowna BC V1W 5C3
Canada

CC:



This report supersedes all previous preliminary and final reports with this file number dated prior to the date on this certificate. Signature indicates final approval; preliminary reports are unsigned and should be used for reference only. All results are considered the confidential property of the client. Acme assumes the liabilities for actual cost of analysis only. Results apply to samples as submitted. *** asterisk indicates that an analytical result could not be provided due to unusually high levels of interference from other elements.



www.acmelab.com

Acme Analytical Laboratories (Vancouver) Ltd.
 9050 Shaughnessy St Vancouver BC V6P 6E5 CANADA
 PHONE (604) 253-3158

Client: **DW Exploration**
 5241 Cobble Crescent
 Kelowna BC V1W 5C3 Canada

Project: Wilcum High Peak
 Report Date: April 22, 2013

Page: 2 of 2

Part: 1 of 1

CERTIFICATE OF ANALYSIS

VAN13001075.1

	Method Analyte Unit MDL	WGHT	1F30	1F30	1F30	1F30	1F30	1F30	1F30	1F30	1F30	1F30	1F30	1F30	1F30	1F30	1F30	1F30	1F30	1F30	1F30	
		Wgt	Mo	Cu	Pb	Zn	Ag	Ni	Co	Mn	Fe	As	U	Au	Th	Sr	Cd	Sb	Bi	V	Ca	
		kg	ppm	ppm	ppm	ppm	ppb	ppm	ppm	ppm	%	ppm	ppm	ppb	ppm	ppm	ppm	ppm	ppm	ppm	ppm	%
		0.01	0.01	0.01	0.01	0.1	2	0.1	0.1	0.1	1	0.01	0.1	0.1	0.2	0.1	0.5	0.01	0.02	0.02	ppm	2
2161301	Rock	1.87	0.29	160.5	2.28	38.1	52	13.6	15.7	380	3.39	3.7	0.2	2.6	1.5	8.8	0.06	0.09	0.03	173	1.40	
2161302	Rock	0.42	0.31	84.00	3.35	33.6	28	3.1	8.9	335	2.88	2.1	0.3	0.9	3.5	19.4	0.06	0.19	0.03	52	1.11	
2161303	Rock	0.20	0.11	6.44	2.95	11.4	13	1.4	0.3	245	0.19	1.7	3.4	0.3	1.2	0.9	0.10	0.02	0.02	<2	0.08	
2161304	Rock	0.26	0.25	127.6	1.68	39.7	87	13.2	10.0	305	2.00	1.5	0.1	2.1	0.8	11.2	0.07	0.16	0.04	63	1.12	
2161305	Rock	0.79	0.07	2.76	4.12	23.9	11	0.8	0.6	655	0.22	1.4	3.1	0.6	13.6	12.1	0.16	0.03	0.50	<2	0.37	
2161306	Rock	0.37	0.11	2.65	4.42	51.3	10	1.6	1.8	516	1.21	1.0	1.8	<0.2	16.9	18.7	0.01	0.03	0.10	12	0.11	
2161307	Rock	0.79	0.08	2.95	10.42	12.6	24	0.7	0.3	312	0.22	2.4	1.6	0.6	2.4	2.0	0.07	0.04	0.19	<2	0.10	
2161308	Rock	0.29	0.20	2.72	2.45	5.9	9	4.4	11.5	40	0.70	2.9	0.3	0.5	1.1	1.2	0.02	0.04	0.04	<2	<0.01	
2161309	Rock	1.07	0.07	2.36	3.44	22.5	11	0.8	0.8	501	0.21	0.9	3.4	*	13.6	10.5	0.09	<0.02	0.66	<2	0.26	
2161310	Rock	1.30	0.36	2.04	3.15	5.9	5	5.5	14.6	52	0.80	5.8	0.5	0.3	0.7	1.3	0.02	0.05	0.10	<2	<0.01	



www.acmelab.com

Acme Analytical Laboratories (Vancouver) Ltd.

9050 Shaughnessy St Vancouver BC V6P 6E5 CANADA

PHONE (604) 253-3158

Client: DW Exploration
5241 Cobble Crescent
Kelowna BC V1W 5C3 Canada

Project: Wilcum High Peak
Report Date: April 22, 2013

Page: 2 of 2

Part: 2 of 1

CERTIFICATE OF ANALYSIS

VAN13001075.1

Method	1F30	1F30	1F30	1F30	1F30	1F30	1F30	1F30	1F30	1F30	1F30	1F30	1F30	1F30	1F30	1F30	1F30	1F30	1F30	1F30	1F30
Analyte	P	La	Cr	Mg	Ba	Ti	B	Al	Na	K	W	Sc	Tl	S	Hg	Se	Te	Ga	Cs	Ge	
Unit	%	ppm	ppm	%	ppm	%	ppm	%	%	%	ppm	ppm	ppm	%	ppb	ppm	ppm	ppm	ppm	ppm	
MDL	0.001	0.5	0.5	0.01	0.5	0.001	1	0.01	0.001	0.01	0.1	0.1	0.02	0.02	5	0.1	0.02	0.1	0.02	0.1	
2161301	Rock	0.049	2.6	1.1	0.66	22.1	0.112	1	1.33	0.229	0.14	<0.1	12.0	<0.02	0.06	<5	0.2	0.05	5.4	0.28	0.1
2161302	Rock	0.093	5.0	0.5	0.37	17.9	0.094	2	1.02	0.152	0.09	<0.1	9.0	<0.02	<0.02	<5	<0.1	0.02	4.7	0.31	0.2
2161303	Rock	0.031	2.1	0.9	<0.01	6.9	0.001	3	0.21	0.046	0.09	0.1	0.3	0.05	<0.02	<5	<0.1	<0.02	0.7	0.57	<0.1
2161304	Rock	0.031	1.3	1.2	0.63	33.1	0.116	1	1.12	0.165	0.10	<0.1	8.1	0.05	<0.02	<5	<0.1	0.07	3.0	1.24	0.2
2161305	Rock	0.188	25.4	1.6	0.03	31.5	0.006	6	0.54	0.011	0.39	0.6	0.8	1.17	<0.02	8	<0.1	0.03	1.5	46.42	0.2
2161306	Rock	0.046	57.3	1.2	0.17	99.3	0.034	<1	0.44	0.039	0.23	0.1	1.4	0.17	<0.02	<5	<0.1	<0.02	3.4	1.43	<0.1
2161307	Rock	0.046	2.6	0.6	<0.01	15.5	<0.001	3	0.16	0.063	0.08	<0.1	0.2	0.05	<0.02	<5	<0.1	0.05	0.5	0.75	<0.1
2161308	Rock	0.010	21.2	1.7	<0.01	3.4	<0.001	<1	0.04	0.005	0.02	<0.1	0.6	<0.02	<0.02	<5	<0.1	<0.02	0.2	0.12	<0.1
2161309	Rock	0.131	26.3	1.5	0.03	33.5	0.005	7	0.54	0.010	0.40	0.5	0.9	1.20	<0.02	7	<0.1	0.04	1.5	45.49	0.2
2161310	Rock	0.011	8.5	1.5	<0.01	4.3	<0.001	<1	0.03	0.005	0.02	<0.1	1.3	<0.02	<0.02	<5	<0.1	0.03	0.2	0.19	<0.1

CERTIFICATE OF ANALYSIS

VAN13001075.1

Method	Analyte	1F30	1F30	1F30	1F30	1F30	1F30	1F30	1F30	1F30	1F30	1F30	1F30	1F30	
		Hf	Nb	Rb	Sn	Ta	Zr	Y	Ce	In	Re	Be	Li	Pd	Pt
Unit		ppm	ppm	ppm	ppm	ppm	ppm	ppm	ppm	ppb	ppm	ppm	ppb	ppb	
MDL		0.02	0.02	0.1	0.1	0.05	0.1	0.01	0.1	0.02	1	0.1	0.1	10	2
2161301	Rock	0.05	0.18	3.4	0.4	<0.05	0.6	9.56	6.5	0.03	3	0.3	3.7	<10	<2
2161302	Rock	0.07	0.35	3.7	0.4	<0.05	0.6	12.13	12.4	<0.02	3	<0.1	3.0	<10	<2
2161303	Rock	<0.02	0.41	12.0	1.7	<0.05	0.3	1.81	4.8	<0.02	<1	0.5	1.2	<10	<2
2161304	Rock	0.04	0.16	11.3	0.2	<0.05	0.5	5.25	3.5	<0.02	<1	<0.1	3.4	<10	<2
2161305	Rock	<0.02	0.39	267.2	21.5	0.09	0.3	32.19	57.1	<0.02	<1	1.6	287.6	<10	<2
2161306	Rock	0.06	2.38	26.7	0.6	<0.05	1.5	9.67	87.0	0.03	<1	0.2	21.9	<10	<2
2161307	Rock	0.07	0.06	7.9	0.6	<0.05	1.8	1.30	6.4	<0.02	<1	0.6	1.2	<10	<2
2161308	Rock	<0.02	0.03	1.1	<0.1	<0.05	0.7	1.55	43.3	<0.02	<1	<0.1	<0.1	<10	<2
2161309	Rock	<0.02	0.31	254.3	20.0	0.11	0.3	26.99	56.5	<0.02	<1	0.8	277.1	<10	<2
2161310	Rock	0.02	0.03	1.5	<0.1	<0.05	0.7	1.64	19.5	<0.02	<1	<0.1	0.9	<10	<2

QUALITY CONTROL REPORT

VAN13001075.1

Method	WGHT	1F30	1F30	1F30	1F30	1F30	1F30	1F30	1F30	1F30	1F30	1F30	1F30	1F30	1F30	1F30	1F30	1F30	1F30	1F30	1F30
Analyte	Wgt	Mo	Cu	Pb	Zn	Ag	Ni	Co	Mn	Fe	As	U	Au	Th	Sr	Cd	Sb	Bi	V	Ca	
Unit	kg	ppm	ppm	ppm	ppm	ppb	ppm	ppm	ppm	%	ppm	ppm	ppb	ppm	ppm	ppm	ppm	ppm	ppm	ppm	
MDL	0.01	0.01	0.01	0.01	0.1	2	0.1	0.1	1	0.01	0.1	0.1	0.2	0.1	0.5	0.01	0.02	0.02	2	0.01	
Pulp Duplicates																					
2161308	Rock	0.29	0.20	2.72	2.45	5.9	9	4.4	11.5	40	0.70	2.9	0.3	0.5	1.1	1.2	0.02	0.04	0.04	<2	<0.01
REP 2161308	QC		0.24	2.46	2.62	7.0	8	4.4	10.9	40	0.70	2.5	0.4	0.4	1.2	1.2	0.03	0.03	0.05	<2	<0.01
Reference Materials																					
STD DS9	Standard		12.69	102.5	124.3	291.2	1776	38.2	7.1	563	2.25	24.6	2.7	107.8	6.5	70.5	2.36	5.73	6.84	38	0.70
STD DS9 Expected			12.84	108	126	317	1830	40.3	7.6	575	2.33	25.5	2.69	118	6.38	69.6	2.4	4.94	6.32	40	0.7201
BLK	Blank		<0.01	<0.01	<0.01	<0.1	<2	<0.1	<0.1	<1	<0.01	0.2	<0.1	<0.2	<0.1	<0.5	<0.01	<0.02	<0.02	<2	<0.01
Prep Wash																					
G1	Prep Blank		0.10	4.90	3.57	96.4	27	2.6	3.8	569	1.89	0.8	2.5	1.1	6.5	55.6	0.24	0.05	0.09	35	0.48
G1	Prep Blank		0.09	4.55	3.34	104.8	16	2.6	3.6	566	1.78	0.7	2.0	0.7	6.1	51.2	0.23	<0.02	0.07	33	0.44



www.acmelab.com

Acme Analytical Laboratories (Vancouver) Ltd.
 9050 Shaughnessy St Vancouver BC V6P 6E5 CANADA
 PHONE (604) 253-3158

Client: **DW Exploration**
 5241 Cobble Crescent
 Kelowna BC V1W 5C3 Canada

Project: Wilcum High Peak
 Report Date: April 22, 2013

Page: 1 of 1

Part: 2 of 1

QUALITY CONTROL REPORT


VAN13001075.1

Method		1F30	1F30	1F30	1F30	1F30	1F30	1F30	1F30	1F30	1F30	1F30	1F30	1F30	1F30	1F30	1F30	1F30	1F30	1F30	1F30
Analyte		P	La	Cr	Mg	Ba	Ti	B	Al	Na	K	W	Sc	Tl	S	Hg	Se	Te	Ga	Cs	Ge
Unit		%	ppm	ppm	%	ppm	%	ppm	%	%	%	ppm	ppm	ppm	%	ppb	ppm	ppm	ppm	ppm	ppm
MDL		0.001	0.5	0.5	0.01	0.5	0.001	1	0.01	0.001	0.01	0.1	0.1	0.02	0.02	5	0.1	0.02	0.1	0.02	0.1
Pulp Duplicates																					
2161308	Rock	0.010	21.2	1.7	<0.01	3.4	<0.001	<1	0.04	0.005	0.02	<0.1	0.6	<0.02	<0.02	<5	<0.1	<0.02	0.2	0.12	<0.1
REP 2161308	QC	0.010	22.5	1.5	<0.01	3.8	<0.001	<1	0.04	0.005	0.02	<0.1	0.7	<0.02	<0.02	<5	<0.1	<0.02	0.2	0.13	0.1
Reference Materials																					
STD DS9	Standard	0.078	13.5	110.9	0.59	279.9	0.106	2	0.92	0.084	0.39	2.8	2.4	4.93	0.16	197	4.4	5.00	4.5	2.32	0.1
STD DS9 Expected		0.0819	13.3	121	0.6165	295	0.1108		0.9577	0.0853	0.395	2.89	2.5	5.3	0.1615	200	5.2	5.02	4.59	2.37	0.1
BLK	Blank	<0.001	<0.5	<0.5	<0.01	<0.5	<0.001	<1	<0.01	<0.001	<0.01	<0.1	<0.1	<0.02	<0.02	<5	<0.1	<0.02	<0.1	<0.02	<0.1
Prep Wash																					
G1	Prep Blank	0.071	13.8	4.5	0.49	161.8	0.116	2	0.91	0.088	0.48	<0.1	2.7	0.31	<0.02	<5	<0.1	<0.02	4.9	3.56	0.2
G1	Prep Blank	0.065	13.1	4.6	0.46	156.6	0.109	2	0.83	0.077	0.46	<0.1	2.6	0.31	<0.02	<5	<0.1	<0.02	4.5	3.54	0.1

QUALITY CONTROL REPORT

VAN13001075.1

Method		1F30	1F30	1F30	1F30	1F30	1F30	1F30	1F30	1F30	1F30	1F30	1F30	1F30	1F30
Analyte		Hf	Nb	Rb	Sn	Ta	Zr	Y	Ce	In	Re	Be	Li	Pd	Pt
Unit		ppm	ppm	ppm	ppm	ppm	ppm	ppm	ppm	ppm	ppb	ppm	ppm	ppb	ppb
MDL		0.02	0.02	0.1	0.1	0.05	0.1	0.01	0.1	0.02	1	0.1	0.1	10	2
Pulp Duplicates															
2161308	Rock	<0.02	0.03	1.1	<0.1	<0.05	0.7	1.55	43.3	<0.02	<1	<0.1	<0.1	<10	<2
REP 2161308	QC	<0.02	0.03	1.3	<0.1	<0.05	0.8	1.67	49.2	<0.02	<1	<0.1	<0.1	<10	<2
Reference Materials															
STD DS9	Standard	0.07	1.47	31.3	6.2	<0.05	1.9	5.73	26.7	2.20	59	5.4	22.5	111	348
STD DS9 Expected		0.08	1.33	33.8	6.4	0.004	2	5.97	25.4	2.2	61	5.4	25.2	120	350
BLK	Blank	<0.02	<0.02	<0.1	<0.1	<0.05	<0.1	<0.01	<0.1	<0.02	<1	<0.1	<0.1	<10	<2
Prep Wash															
G1	Prep Blank	0.08	0.85	45.0	0.6	<0.05	1.2	5.63	26.5	0.03	<1	0.6	31.6	<10	<2
G1	Prep Blank	0.06	0.84	42.6	0.5	<0.05	1.0	4.94	24.8	0.02	1	0.1	31.7	<10	<2



Legend


- ◆ Sample Location
- ▭ Wilcum License
- ▭ Other License
- Road/Track
- River/Stream
- Lake
- Quaternary Cover

DW Exploration

**Wilcum High Peak Property
2012 Hand Trench Locations
Locations By**

Date: 17/05/2013
 Author: Ivdm
 Office: van
 Drawing: Ivdm
 Scale: 1:25000
 Projection: UTM Zone 11 (NAD 83)

0 250 500 1,000
metres



WILCOM 1

WILCUM3

WILCUM4

WILCUM2

WILCUM5

WILCUM 6

WILCUM 7

87 ppb
13 ppb
52 ppb
28 ppb
10 ppb
9 ppb
24 ppb
11 ppb
11 ppb
5 ppb

