

Ministry of Energy and Mines
BC Geological Survey

**ASSESSMENT REPORT
TITLE PAGE AND SUMMARY**

TITLE OF REPORT [type of survey(s)]	TOTAL COST
Geological Report on the J Property, 150 Mile House Area, Cariboo Mining Division, British Columbia	\$3690.45

AUTHOR(S) David Bridge, P. Geo. SIGNATURE(S) David Bridge

NOTICE OF WORK PERMIT NUMBER(S)/DATE(S) _____ YEAR OF WORK 2017

STATEMENT OF WORK - CASH PAYMENT EVENT NUMBER(S)/DATE(S) 5673585 (2017/Nov/13)

PROPERTY NAME J Property

CLAIM NAME(S) (on which work was done) Tenure (1047973)

COMMODITIES SOUGHT Copper, silver,

MINERAL INVENTORY MINFILE NUMBER(S), IF KNOWN 093A 066

MINING DIVISION Cariboo NTS 093A/12E

LATITUDE 52 ° 32 ' 17 " LONGITUDE 121 ° 44 ' 13 " (at centre of work)

OWNER(S)
1) Jedway Enterprises Ltd. 2) _____

MAILING ADDRESS
104-19286 21st Avenue
Surrey, BC V3S 3M3

OPERATOR(S) [who paid for the work]
1) Jedway Enterprises Ltd. 2) _____

MAILING ADDRESS
104-19286 21st Avenue
Surrey, BC V3S 3M3

PROPERTY GEOLOGY KEYWORDS (lithology, age, stratigraphy, structure, alteration, mineralization, size and attitude):
Nicola Group, basaltic volcanic rocks and volcanoclastic rocks, Redbed copper, High grade copper veins

REFERENCES TO PREVIOUS ASSESSMENT WORK AND ASSESSMENT REPORT NUMBERS 885, 4683, 12589, 20792, 25261, 25960, 26614, and 32283

TYPE OF WORK IN THIS REPORT	EXTENT OF WORK (IN METRIC UNITS)	ON WHICH CLAIMS	PROJECT COSTS APPORTIONED (incl. support)
GEOLOGICAL (scale, area)			
Ground, mapping	1:500 2 ha	1047973	\$2090.45
Photo interpretation	1:9146 150 ha	1047973	\$1600.00
GEOPHYSICAL (line-kilometres)			
Ground			
Magnetic _____			
Electromagnetic _____			
Induced Polarization _____			
Radiometric _____			
Seismic _____			
Other _____			
Airborne _____			
GEOCHEMICAL			
(number of samples analysed for ...)			
Soil _____			
Silt _____			
Rock _____			
Other _____			
DRILLING			
(total metres; number of holes, size)			
Core _____			
Non-core _____			
RELATED TECHNICAL			
Sampling/assaying _____			
Petrographic _____			
Mineralographic _____			
Metallurgic _____			
PROSPECTING (scale, area) _____			
PREPARATORY/PHYSICAL			
Line/grid (kilometres) _____			
Topographic/Photogrammetric (scale, area) _____			
Legal surveys (scale, area) _____			
Road, local access (kilometres)/trail _____			
Trench (metres) _____			
Underground dev. (metres) _____			
Other _____			
TOTAL COST			\$3690.45

Jedway Enterprises Ltd.
104-19286 21st Ave, Surrey, BC V3S 3M3

Geological Report on the J Property, 150 Mile House Area, Cariboo
Mining Division, British Columbia

Trim 093A052

BC Geological Survey
Assessment Report
37266

Latitude 52°32'17"North
Longitude 121°44'13" West

Program: June 19 and June 21, 2017

Author: David Bridge, P.Geo
1580-132B Street, Surrey, BC, V4A 6J2

Date: February 6, 2018

Revised on July 30, 2018 with addition of a geological mapping program

Summary:

The J Property is located on BCGS Map 093P052 and it is roughly 50 kilometers north of 150 Mile House in the Cariboo Mining Division on the Cariboo Plateau. The area of interest is roughly centered at Latitude 52°32'17" North and Longitude 121°44'13" West. A geological program of air photograph interpretation at a scale of approximately 1: 9146 was done using Google Earth photos and BC Trim maps from www.MapPlace2.gov.bc.ca covering the J Property where redbed copper and vein copper veins occurs. The vein mineralization assays up to 7.12% copper over 5 meters from samples collected in 1999 from a trench and grab samples collected in 2011 of redbed mineralization assay up to 5.41% copper.

Analysis of topographical map contours superimposed on air photographs shows that the Jurassic redbed copper mineralization which formed in Nicola Group Triassic volcanic rocks is cut by herringbone faults related to Cretaceous magnetism and hydrothermal mineralization and later the intersections of the herringbone faults were copper mineralized during hydrothermal activity in the Eocene. The sample evidence for the Eocene hydrothermal activity is located immediately east of the J Property. The Cretaceous herringbone faults form northwesterly and northeasterly conjugate pairs where the northeasterly faults have dextral offsets which offset the northwesterly faults. Intruding up the northwesterly fault in the middle of the J#1 mineral claim is a Cretaceous feldspar porphyry intrusion of roughly 15 meters by 50 meters in size as determined from a ground magnetometer survey performed in 1999.

The ages of the mineralization and hydrothermal activity were determined from the mathematical relationship of the concentration of U (ppm) divided by Th (ppm) of the rock sample and the quotient divided by 3 and the result multiplied by 960 million years - the rough age of the mantle beneath Quesnellia. This calculated age is the rough age of the rising and cooling magma as it rises and cools below 1300°C - the temperature when UO_3 decrepitates to UO_{2T} and fixes the age by loss of the U daughter products. This age can only be used if there has been no addition of U by hydrothermal fluids between 177°C and 399°C when U chlorides are mobile. If there has been alteration by later fluids which do not add U the ratio of U/Th may gradually change to the age of the hydrothermal alteration.

The redbed copper mineralization and the high grade copper veins on the J Property should be surveyed with an IP geophysical survey and later diamond drilled with shallow holes if the geophysical results warrant it.

A geological mapping program was conducted in late July 2018 in the northern part of the property (Appendix 1) and discovered jasper and epidote veined oxidized Upper Triassic Nicola Group basalt breccia in the vicinity of a possible Cretaceous potassic granite dyke with extensional – shear calcite veinlets in basalt breccia in similar orientation to the north-westerly faults.

TABLE OF CONTENTS

Summary	2
Introduction	4
Location and Access	4
General Setting, Climate and Infrastructure	5
History and Previous Work	6
Regional Geology	7
2017 Work Program	8
Interpretation of Results	8
Conclusions and Recommendations	9
References	10
Cost Statement	11
Statement of Qualifications	12
Appendix 1: July 2018 Geological Mapping program	13
 Figures and Tables	
Figure 1. Location Map	4
Figure 2. Location and Access map and Index map	5
Figure 3. Claim map	6
Figure 4. Regional Geology map	7
Figure 5. J Property air photo interpretation and geological stations	in pocket
Table 1. Mineral Claim data	6
Table 2. Actual and calculated ages of select rocks and mineralization on the J Property	9

Introduction:

This report has been commissioned by Jedway Enterprises Ltd. for the purposes of filing an assessment report on the J Property. The Air photo interpretation which is subject of this report was conducted by David Bridge P. Geo for during partial days in June, 2017 and a geological mapping program was conducted in late July 2018 after the work was filed on the property.

Location and Access

The J Property is located on BCGS map 093A052 and the area of interest is situated at Latitude $52^{\circ}32'17''$ North and Longitude $121^{\circ}44'13''$ West. The Property is located in the Cariboo Mining Division and is approximately 50 kilometers north of 150 Mile House in the Cariboo Plateau (Figure 1).



Figure 1. Location Map

Access to the property is by the Jacobie Lake Forest Service road for approximately 4 km from the Likely Road in central British Columbia approximately 50 kilometers north of 150 Mile House on Highway 97.




Figure 2. Topographical and Index map of J Property excerpted MapPlace2.gov.bc.ca.

General Setting, Climate and Local Infrastructure:

The J Property is located on the Cariboo Plateau roughly 50 km north of 150 Mile House at the eastern edge of the Jacobie Lake. The property is located roughly 1200 meters and is composed of rolling hills and shallow creeks. This area has been nearly completely logged in 1993 with second growth of pine trees growing on it. The J property receives an estimated up to 0.75 meters of snow and is thought to be generally snow free from May to the end of October.

The property is located 50 kilometers north of the 150 Mile House which is a small business area in the region besides the larger Williams Lake - 15 kilometers further along Highway 97 to the west.

The J Property consists of 2 mineral claim totaling 452.12 hectares and the geological air photo interpretation was conducted on tenure 1047973 (Figure 3, Table 1). Tenure 1056597 was staked after recording work on the J Property.

Table 1: Mineral claim data

Title Number	Claim Name	Good To Date	Area (ha)
1047973	J#1	2019/Sep/21	393.19
1056597	J#3	2018/Nov/22	58.93

The new expiry dates of the mineral claims are subject to the approval of the work contained in this report.

J Property (Tenures 1047973 and 1056597)




Figure 3. J Property mineral claim map


History and Previous Work

The area of the J Property was first explored in 1966 by Chataway Explorations who completed soil sampling in the area (Assessment Report 885). In 1972, the area was explored by Sunshine Valley Minerals who completed a program of geological mapping and rock sampling as the B claims (Assessment Report 4683). In 1984 the area was explored by Hennessy Resource Corp. who completed a program of rock and soil sampling and geological mapping as the Jacobe 2 claim (Assessment Report 12589). In 1990 the area was explored by Pamicon Developments who prospected and conducted rock sampling and obtained up to 7.29% copper and 35.9 g/t silver in grab samples (Assessment Report 20792). In 1997 the area was explored by Navarre Resources who completed 17.5 kilometers ground

electromagnetic (BEEP) survey and an 11.6 line-kilometer ground magnetic survey (Assessment Report 21516). In 1999 the area was explored by Globex Mining Enterprises Inc. and they conducted rock sampling, trenching, geological mapping and a 5.6 line-kilometer ground magnetic survey and they obtained a 5 meter interval of 7.12% copper from a trench which had a massive 0.2 meter thick chrysocolla and chalcocite vein in it and a magnetic low where a fessite intrusive dyke occurs on the property (Assessment Report 25960). In 2001 the area was explored by Phelps Dodge Corporation of Canada and they completed a program of geological mapping and rock and soil sampling (Assessment Report 26614). In 2011 the area was explored by Eagle Peak Resources who completed a program of geological mapping and rock and soil sampling and obtained up to 5.41% copper from grab samples of redbed copper mineralization from an old trench (Assessment Report 32283).

Regional Geology

The J Property redbed copper mineralization is hosted by Upper Triassic Nicola Group volcanoclastic rocks which lie above Upper Triassic Nicola Group basaltic volcanic rocks that are in part in fault contact with the volcanoclastic rocks (www.MapPlace2.gov.bc.ca). These rocks are locally intruded by a Cretaceous feldspar porphyry dyke which intruded along local northeasterly block faults which form a conjugate pair with dextral northwesterly striking faults. The age of the intrusive was determined by its concentration of Uranium (ppm) divided by its Thorium (ppm) and then divided by 3 and multiplied by 960 Ma from hydrothermally altered samples from the edge of the intrusive body. There is evidence from immediately east of the J Property of hydrothermal alteration coming up the intersections of the northwesterly and northeasterly faults during the Eocene.



- uTvolcl Nicola Group, Upper Triassic volcanic clastic rocks
- uTvol Nicola Group, Upper Triassic basaltic volcanic rocks

Figure 4: Regional Geology map of J Property excerpted from [www. MapPlace2.gov.bc.ca](http://www.MapPlace2.gov.bc.ca)

2017 Work Program:

A couple of partial days were spent by the author completing air photograph interpretation of air photos from Google Earth coupled with the analysis of a trim map from www.MapPlace2.gov.bc.ca covering the J Property from June 19 to June 21, 2017 and a one day geological mapping program was conducted in late July 2018 after the work was filed on the property. (Figure 5, Appendix 1).

Interpretation of Results:

Analysis of the air photograph coupled with the trim map from www.MapPlace2.gov.bc.ca shows that there are several linear features which might be brittle faults cutting across the landscape which form a herringbone pattern (Figure 5). Analysis of the location of mineralization discovered in trenches and the location of linear features show that the northwesterly linear features are dextral offsetting northeasterly faults which locally host a magnetic low approximately 15 meters wide by 50 meters long which has outcrops of feldspar porphyry of possible Cretaceous age in it. These faults making the

herringbone pattern on the J Property may be related to the Cretaceous intrusive activity on the property.

The age of the redbed copper mineralization which is composed of chalcocite- covellite- cuprite – malachite – native copper filled fractures and amygdules locally assays up to 5.41% in grab samples (Assessment Report 32283) and has been determined to be Jurassic in age from the concentration of U (ppm) divided by Th (ppm) and divided by 3 and multiplied by 960Ma (Table 2) (Assessment Report 26614) and the age of the intrusive intruding the Jurassic mineralization has been determined to be Cretaceous using the similar method and the late stage high grade copper veins composed of chrysocolla and chalcocite to be Eocene in age and these assay up to 7.12% copper over 5 meters. In all of these calculations an age of 960 Ma was used which is the rough age of the mantle beneath terranes offshore to the North American Craton. This age was determined by the oldest rocks in the Slide Mountain terrane in the East and North Barriere Lake area southeast of Clearwater, British Columbia (www.MapPlace2.gov.bc.ca). The resulting ages are in fact the age of the hydrothermal fluid and magma as it rises above the mantle and cools below 1300°C when UO_3 decrepitates to UO_2 and loses its daughter elements (Speight, 2005). This method can only be used if U is not added to the altered rock from fluids transporting Uranium chloride complexes between 177°C and 399°C (Speight, 2005). Hydrothermal fluids can later alter the U/Th ratio of host rocks to the age of the hydrothermal alteration by constantly flushing the rock with the hot water with the new U/Th ratio of the alteration.

Table 2

Actual and calculated ages of select rocks and mineralization on the J Property of Jedway Enterprises Ltd. (Samples from Ass Report 26614A)

Sample ID	Description	Actual age from mapping	UTM E	UTM N	U (ppm)	Th (ppm)	Ratio (U/TH)/3	Calculated age (Ma)
75572	Cu-Ag amygdules in Maroon Basaltic tuff	Jurassic	585256	5821807	0.6	1.1	0.181666667	174.40
75575	Re-opened NW Cretaceous structure with Chalcocite/Malachite vein in fragmental basalt (Sample is located immediately east of J Property)	Eocene	586406	5821987	0.3	1.7	0.058823529	56.47
75576	Cretaceous porphyry intrusive - Feldspar porphyry andesite with malachite staining	Cretaceous	584612	5821987	0.5	1.7	0.098039216	94.12

Conclusion and Recommendations

More prospecting, sampling and geological mapping needs to be done on the J Property to follow up on the air photo interpretation and to investigate the potential of the redbed copper mineralization and high grade copper veins and this should be followed by IP surveying and if the results warrant it diamond drilling with shallow drill holes.

References:

Assessment Report 885: Wright, S.W., 1967. Geochemical Survey, Jacobie Group of Mineral Claims, 10 pages.

Assessment Report 4683: Campbell, J. I., 1972. Reconnaissance of B Property, 27 pages

Assessment Report 12589: Simpson, R.G., 1984. Geological Mapping and Geochemical Survey Report on the Jacobe 2 Mineral Claim, 29 pages.

Assessment Report 20792: Montgomery, A., 1991. 1990 Assessment Report on a Prospecting and Geological Work Program, 55 pages

Assessment Report 25261: Kikauka, A, 1997. Geophysical Report on the JC1 claim, Jacobie Lake, Likely B.C., 47 pages

Assessment Report 25960: Kikauka, A., 1999. Geological and Geophysical report on the J 1-4, Claim Group, Jacobie Lake, Likely, B.C., 23 pages

Assessment Report 26614: Kulla, G., 2001. Geophysical and Geochemical Report on the Morehead Property, 28 pages

Assessment Report 32283: Fox, P.E, 2011. Geological, Geochemical Report Jacobie Lake Prospect, 38 pages.

Speight, J.G., 2005. Lange's Handbook of Chemistry, Sixteenth Edition, McGraw-Hill.

Software and Websites used

Corel Draw

MS Windows, MS Word, MS Excel

www.MapPlace2.gov.bc.ca; www.MtOnline.bc.ca and www.MapPlace.gov.bc.ca

Cost Statement:

J Property Cost Statement

Air Photograph and geophysical interpretation (Partial days during the period June 19 and June 21, 2017)

\$500.00

Report and post-processing of data

\$2200.00

Geological Mapping program (July 21, 2018)

Eric Mackenzie (\$400/day)

\$400

David Bridge (\$308/day)

\$308

4x4 Vehicle (654 km at \$0.40/km)

\$261.60

Food (\$20.85)

\$20.85

Total

\$3690.45

STATEMENT OF QUALIFICATIONS FOR David Bridge, P.Geo

I, David Bridge, hereby certify that:

I am a geologist residing at 1580-132B Street, Surrey, British Columbia, Canada.

I am a graduate of the University of British Columbia with a Bachelors degree in Geological Engineering (1990) and a Masters in geological engineering in (1994).

I am registered as a Professional Geoscientist with the Association of Professional Engineers and Geoscientists of British Columbia (APEGBC number 24944).

I completed the air photo interpretation on the J Property during partial days on June 19 and June 21, 2017 and it is subject of this report and conducted the geological mapping program July 21, 2018.

Dated at Surrey, BC

July 30, 2018

Respectfully submitted

"David J. Bridge"

David J. Bridge, P. Geo, MASc


APPENDIX 1

Geological Mapping Program July 21, 2018

A one day geological mapping program was conducted in the northern part of the J Property in late July 2018 and discovered a weakly carbonate altered Upper Triassic Nicola Group basalt breccia which grades into oxidized Upper Triassic Nicola Group basalt breccia with jasper – epidote veins and possible yellow magnesite filled vesicles away from veins in the vicinity of a possible Cretaceous potassic granite dyke which was trending approximately north-westerly in similar orientation to north-westerly herringbone faults in the southern part of the J Property (see geological station descriptions listed below). The oxidized basalt breccia has also extensional – shear calcite veins in similar orientation to the north-westerly faults.

Geological stations on J Property completed on July 21, 2018 for assessment report by Jedway Enterprises Ltd.

Station	Easting	Northing	Description
111	585145	5823278	Oxidized reddish brown outcrop in roadway of fine grained basalt breccia with planer calcite veinlets up to 2 mm thick striking N 70 E and northerly. There was an oxidised basalt breccia float boulder near the outcrop which has epidote and jasper vein across it and possible yellow magnesite filled vesicles approximately 12 cm away from vein measured on a broken surface of up to 2-3 mm in diameter.
112	585140	5823263	Vesicular dark green basalt with cream calcite stain and carbonate and hematite filled vesicles. Outcrop in road is roughly 1 by 1 meter. The contact between oxidized and unoxidized basalt is between stations 111 and 112 and it is not exposed in the roadway but is roughly half way between outcrops and trends east-west.



NAD 83 UTM Zone 10 Mapsheet 093A052

— Airphoto linears which might be conjugate faults that developed due to Cretaceous magmatism.

○ Cretaceous Feldspar porphyry intrusion

× Rock Samples described in text from Assessment Report 26614

1.0 Kilomteres

Jedway Enterprises Ltd.
J Property
Photogrammetric and
Geological Study
150 Mile House Area, BC

Figure 5 Drawn by: DJB Feb. 2018
and revised July 2018

Geological stations on J Property completed on July 21, 2018 for assessment report by Jedway Enterprises Ltd.			
Station	Easting	Northing	Description
111	585145	5823278	Oxidized reddish brown outcrop in roadway of fine grained basalt breccia with planer calcite veinlets up to 2 mm thick striking N 70 E and northerly. There was an oxidized basalt breccia float angular boulder near the outcrop which has epidote and jasper vein across it and possible yellow magnesite filled vesicles approximately 12 cm away from vein measured on a broken surface of up to 2-3 mm in diameter.
112	585140	5823263	Vesicular dark green basalt with cream calcite stain and carbonate and hematite filled vesicles. Outcrop in road is roughly 1 by 1 meter. The contact between oxidized and unoxidized basalt is between stations 111 and 112 and it is not exposed in the roadway but is roughly half way between outcrops and trends approximately east-west.