



Ministry of Energy, Mines & Petroleum Resources
Mining & Minerals Division
BC Geological Survey

Assessment Report
Title Page and Summary

McGILL

TYPE OF REPORT [type of survey(s)]: Geochemical

TOTAL COST: \$40,000.00

AUTHOR(S): J. T. Shearer, M.Sc. P.Geo.

SIGNATURE(S):

NOTICE OF WORK PERMIT NUMBER(S)/DATE(S): _____

YEAR OF WORK: 2020

STATEMENT OF WORK - CASH PAYMENTS EVENT NUMBER(S)/DATE(S): 5818069

PROPERTY NAME: McGillivray

CLAIM NAME(S) (on which the work was done): _____

COMMODITIES SOUGHT: Au/Cu/Ag

MINERAL INVENTORY MINFILE NUMBER(S), IF KNOWN: _____

MINING DIVISION: Kamloops

NTS/BCGS: 921/12 (921.042+-052)

LATITUDE: 50 ° 29 ' 45 " LONGITUDE: 121 ° 40 ' 30 " (at centre of work)

OWNER(S):

1) J. T. Shearer

2) _____

MAILING ADDRESS:

Unit 5 - 2330 Tyner Street

Port Coquitlam, BC V3C 2Z1

OPERATOR(S) [who paid for the work]:

1) Same as above

2) _____

MAILING ADDRESS:

Same as above

PROPERTY GEOLOGY KEYWORDS (lithology, age, stratigraphy, structure, alteration, mineralization, size and attitude):

The claims are underlain by large zones of alteration (pyritization+kaolin) and anomalous geochemistry typical of an
epithermal Au/Ag system, perhaps adjacent to porphyry intrusions, major faults cut the area juxtaposing Cretaceous Spences
Bridge Group with Lytton Complex Metamorphics

REFERENCES TO PREVIOUS ASSESSMENT WORK AND ASSESSMENT REPORT NUMBERS: _____

Assessment Reports 3154, 11371, 7027, 4455, 28182, 3154, 16352A+B, 3938, 3937, 12948, 12944, 8762, 7027

TYPE OF WORK IN THIS REPORT	EXTENT OF WORK (IN METRIC UNITS)	ON WHICH CLAIMS	PROJECT COSTS APPORTIONED (incl. support)
GEOLOGICAL (scale, area)			
Ground, mapping			
Photo interpretation			
GEOPHYSICAL (line-kilometres)			
Ground			
Magnetic			
Electromagnetic			
Induced Polarization			
Radiometric			
Seismic			
Other			
Airborne			
GEOCHEMICAL (number of samples analysed for...)			
Soil			20,000
Silt			
Rock			5,000
Other			
DRILLING (total metres; number of holes, size)			
Core			
Non-core			
RELATED TECHNICAL			
Sampling/assaying			
Petrographic			
Mineralographic			
Metallurgic			
PROSPECTING (scale, area)			15,000
PREPARATORY / PHYSICAL			
Line/grid (kilometres)			
Topographic/Photogrammetric (scale, area)			
Legal surveys (scale, area)			
Road, local access (kilometres)/trail			
Trench (metres)			
Underground dev. (metres)			
Other			
TOTAL COST:			\$40,000.00

GEOCHEMICAL ASSESSMENT REPORT
on the
McGILLIVRAY PROJECT
- A Porphyry Copper-Gold Project -

Lytton-Lillooet Area of British Columbia

NTS 92I/12 (92I.042+052)
Latitude 50°29'45"N/Longitude 121°40'30"W
Permit MX-4-480
Event #5818069

For

Homegold Resources Ltd.
#5-2330 Tyner St.
Port Coquitlam, B.C.
V3C 2Z1
Phone: 604-970-6402
Fax: 604-944-6102

Prepared by

J. T. SHEARER, M.Sc., P.Geo. (BC & Ontario) FSEG
E-mail: jo@HomegoldResourcesLtd.com

November 17, 2020

Fieldwork Completed between March 15, 2020 and November 17, 2020

TABLE OF CONTENTS

	<u>Page</u>
LIST of FIGURES	ii
LIST of PHOTOS	ii
SUMMARY	iii
INTRODUCTION	1
PROPERTY CLAIM STATUS.....	4
LOCATION and ACCESS	7
HISTORY	8
FIELD PROCEDURES	29
GEOLOGICAL SETTING	
Regional Geology	31
Local Geology.....	33
Petrology	35
EXPLORATION 2020	36
CONCLUSIONS and RECOMMENDATIONS	50
Estimate of Future Costs.....	51
REFERENCES	52
APPENDICES	
APPENDIX I: Statement of Qualifications	55
APPENDIX II: Statement of Costs	56
APPENDIX III: Sample Description	57
APPENDIX IV: Assay Results.....	60

TABLES

	<u>Page</u>
TABLE I Claim Status.....	4

FIGURES

	<u>Page</u>
FIGURE 1	Location Map..... 1
FIGURE 2	Claim Group Location and Access Map 2
FIGURE 3	Detail Access Map..... 3
FIGURE 4	Claim Map Overview 1-50,000..... 6
FIGURE 5	2009 Soils Grid 11
FIGURE 6	2006 Soil Samples 12
FIGURE 7	McGillivray Cliff Exposure 13
FIGURE 8	Airphoto Interpretation 14
FIGURE 9	Area of Previous High Au in Soils 15
FIGURE 10	Garmin Rock Sample Locations and Results 18
FIGURE 11	2019 Magnetometer Traverses..... 19
FIGURE 12a-12q	Magnetometer Maps 1-16 20-28
FIGURE 13	Regional Geology 30
FIGURE 14	Regional Geology 32
FIGURE 15	Detail Geology 34
FIGURE 16	Sample Locations and Results Bob Creek 37
FIGURE 17	Sample Locations and Results Bob Zone 38
FIGURE 18	Sample Locations and Results Lower Bob Creek..... 39
FIGURE 19	Google Image Showing Eastern Samples by S.Butler 40
FIGURE 20	Google Image Showing Western Samples by S.Butler 41
FIGURE 21	Alice Hill Stratigraphic Section..... 49

PHOTOS

	<u>Page</u>
PHOTO 1	Exposed Bedrock Along Alice Hill 41
PHOTO 2	Looking Northwest Showing Compositional Change 42
PHOTO 3	Old Adit at 666 elev. 43
PHOTO 4	Old open cut/pit 43
PHOTO 5	Structurally Controlled Vertical Dipping Cu/Ag Vein 44
PHOTO 6	Malachite-Copper Stained Iron Breccia 44
PHOTO 7	Close-up of Typical Mineralized Iron Carbonate..... 45
PHOTO 8	Sample Collected Adjacent to Adit Entrance..... 46

SUMMARY

The McGillivray property consists of 20 claims, acquired to cover an historical copper-porphyry target, with a large gossanous alteration zone to explore for its precious metal potential. The McGillivray property is near the epithermal precious metal Skoonka Creek property. The claims cover ground originally staked in the 1940's. Previous work in the area covered by the property outlined large zones of hydrothermal alteration and copper anomaly in soil geochemistry at the time when the focus of much exploration was toward porphyry copper targets.

The property is centered on McGillivray Creek and is located 34 kilometres east-southeast of Lillooet, British Columbia and is well served by roads and power. The claims are about midway between Lytton and the Blustry Mountain Gold Showings, on the east side of the Fraser River.

The McGillivray Property geology consists of fault bounded slices of dioritic and granodioritic intrusives of the Permian to Triassic age Mount Lytton Complex, with highly altered Lower Cretaceous andesitic volcanics of the Pimainus Formation of the Spences Bridge group. The western side of the McGillivray Property is next to the regional Fraser River fault. Within the fault bounded slices of altered volcanics are strong alteration zones with evidence of drusy quartz and anomalous silver soil samples.

However, the McGillivray property does demonstrate many features of classic epithermal deposits: the vein mineralogy and textures, with generations of carbonate, silica and chalcedony, the tendency for mineralization to occur in flat vein structures, the presence of brecciated quartz veins, and the suite of geochemical indicator elements Mo, As, and Ag.

Previous work in 2009 consisting of trenching, follow-up soil sampling, prospecting and geology which has confirmed the potential for an epithermal gold-silver and porphyry copper style mineralized systems.


Current work in 2020 focussed on continued rock geochemistry and soil samples in the lower elevations.

Assay results are plotted on Figure 13 and XRF results are contained in Appendix III. Values plotted on Figure 13 are Si%, Fe% and Cu ppm.

A total of 22 rocks and 51 soils were collected in the area of the "Bob" zone on claim 1075219. Results are plotted on Figures 17, 18 and 19 and Figures 20 and 21 on Claim 1074022 and soils were anomalous up to 407ppm Cu (Sample M20073). Rock samples returned very high results up to 8.09% Cu (Sample MR20023)

Soil sampling lines were run across prominent linears (fault zones/contact zones) as shown on Figures 20 and 21. The results indicate a marked change on either side of the linears.

Respectfully submitted,



J. T. Shearer, M.Sc., P.Geo. (BC & Ontario) FSEG

INTRODUCTION

The purpose of this report is to document the 2020 exploration program on the McGillivray Creek property.

This report is largely based on fieldwork conducted between October 19 and November 1, 2020, the historical reports of previous operators and government geological mapping. The author also discussed ongoing activities with the field exploration crew and Dan G. Cardinal, P.Geol. during the program. The documents reviewed are listed in the References near the end of this report.

Attention has focussed on a new belt of newly discovered gold showings nearby on the Skoonka Creek gold property, which represents a new gold discovery in southwestern BC. An initial drilling program completed in October 2005 on the JJ prospect returned high grade gold values including 20.2 g/t gold over 12.8 metres, 26.8 g/t Au over 3.31 metres and 7.5 g/t Au over 4.1 metres. Mineralization has been traced over a strike length of 350m and remains open to the east and west as well as to depth.

The Skoonka Creek property is about 12km southeast of the McGillivray Claims along the regional trend.




Figure 1 Location Map




Figure 2 Claim Group Location/General Access Map




Figure 3 Detail Access Map

PROPERTY CLAIM STATUS

Property Status (List of Claims)

The property consists of the following 11 mineral claims as tabulated in Table 1 and illustrated on Figure 3. The claims are in the Kamloops and Lillooet Mining Divisions.

Claim Status

The staked claims are recorded as follows:

Tenure Number	Claim Name	Size (ha)	Date Located	* Current Anniversary Date	Registered Owner
1057727	Alice/McGill	20.55	January 17, 2018	January 14, 2024	J. T. Shearer
1061539	McGill	41.12	July 3, 2018	December 3, 2022	J. T. Shearer
1061586	McGill 30	41.12	July 6, 2018	December 6, 2022	J. T. Shearer
1061587	McGill 31	164.48	July 6, 2018	December 6, 2022	J. T. Shearer
1061589	McGill 32	20.55	July 6, 2018	December 6, 2022	J. T. Shearer
1061863	Alice 7	164.43	July 18, 2018	December 18, 2022	J. T. Shearer
1061864	McGill 7	287.77	July 18, 2018	December 18, 2022	J. T. Shearer
1061879	McGill 44	123.37	July 20, 2018	December 20, 2022	J. T. Shearer
1065227	McGill 9	185.02	December 21, 2018	December 21, 2023	J. T. Shearer
1065848	McGill 8	61.67	January 18, 2019	January 18, 2023	J. T. Shearer
1065949	McGill South	102.82	January 22, 2019	December 22, 2022	J. T. Shearer
1073427	McGill 10	246.57	December 22, 2019	December 22, 2023	J. T. Shearer
1073432	McGill 11	41.11	December 22, 2019	December 22, 2022	J. T. Shearer
1074019	McGill 20	41.14	January 21, 2020	December 21, 2022	J. T. Shearer
1074020	McGill 21	411.4	January 21, 2020	December 21, 2022	J. T. Shearer
1074021	McGill 22	699.62	January 21, 2020	December 21, 2022	J. T. Shearer
1074022	McGill 23	205.84	January 21, 2020	December 21, 2022	J. T. Shearer
1074023	McGill 24	82.28	January 21, 2020	December 21, 2022	J. T. Shearer
1074547	McGill 25	205.66	February 12, 2020	December 12, 2022	J. T. Shearer
1075219	McGill 26	617.66	March 14, 2021	December 14, 2022	J. T. Shearer

Total 3,764.18 hectares

* Subject to approval of work documented in the Assessment Report

Following revisions to the Mineral Tenures Act on July 1, 2012, claims bear the burden of \$5 per hectare for the initial two years, \$10 per hectare for year three and four, \$15 per hectare for year five and six and \$20 per hectare each year thereafter.

The claims are located in mapsheets 92I-042 and 92I-052. The latitude 50°29'45"N and longitude 121°40'30"W are near the center of the area that work was done in 2006.


Most of these claims are located on Crown Land and have no surface rights attached to the claims. There is crown land available for use by permit application through a permit for development of a mill and tailings if the project moves to this level.

There is a small adit, with a tennantite showing, above McGillivray Creek, on the north side, described in the BC Ministry of Mines, Geological Fieldwork (White, 1980). There are reports of several small pits on the property near the highway described by Chisholm (1971). There are several filled in bulldozer or excavator trenches at the end of the logging road that likely date from 1972 or 1973, near the centre of the previous 2009 fieldwork area.

The property is within the territorial land of the Lytton First Nations band (NNTC).

There are no known new parks planned for any area contained within the McGillivray Property. No First Nations reserves are indicated on the claims maps within the boundaries of the McGillivray claims.

There is a network of logging roads and several clear-cut openings from previous logging operations. The environmental liabilities of this will be the responsibility of the logging companies. The creeks are often steep and the semi-arid environment limits the amount of water in creeks. It is not known whether there are any fish in the claim area. Wildlife throughout the area is sparse and primarily comprises deer and rare, itinerant black bears. Hill slopes are seasonal range for cattle.



McGillivray Claim Map
 Figure 4 Claim Map

Universal Transverse Mercator - Zone 10 (N)
 1:35000
 Printed at: 2020-10-25

LOCATION and ACCESS

The property is located on the eastern side of the Fraser Canyon. Highway 12, which follows the river, crosses the western side of the property. There is a network of logging roads over the property, accessing the highway on the north side of the McGillivray Creek valley. The Luluwissin Creek valley is to the south end of the property and has a deactivated logging road in its lower valley leading to the Fraser River. The upper valley of the Luluwissin Creek road, along the eastern edge of the claims, is accessed from Highway 12 in the Izman Creek valley to the south. This road connects up to a network of logging roads from Spences Bridge and Cache Creek on Highway 1 and near Pavilion on Highway 99. Most of the property is accessible by foot with cliff and landslide exposures limiting foot travel in a few areas.

Elevations range from 450m (1,500ft.) in the valley of the Fraser River to 1,800m (5,900ft.) on McGillivray Mountain. The area is mountainous with steep slopes. There are a couple of large natural slides and cliffs on the property, especially on the south facing slopes.

The area lies in the rain shadow of the Coast Mountains, therefore the climate is relatively dry; Lytton receives less than 40 cm precipitation per annum, of which 25% falls as snow during the winter months. Mean temperatures vary from -4°C in °winter to 30°C in the summer. At lower elevations, the vegetation is open pine forest. The north facing slopes have locally thick forests of pine and fir. The area is grazed by cattle during the summer months. Work can continue throughout the year although snow will likely be present on the ground throughout the winter limiting the activities and slowing access.

There are electrical power lines following Highway 12 on the western side of the property. McGillivray Creek and its tributaries had water sufficient for drilling in October after a long dry spell and should be sufficient for year round exploration. There is abundant water in the Fraser River for any need on the property.

Lillooet and Lytton are the major towns in the area, both on Highway 12. Lillooet the regional source of most required supplies, heavy equipment and services for exploration is 34 kilometres by highway to the north-north west. The regional population is about 50,000. The major industries include logging, ranching and hospitality. The distance to Lytton is about 15 kilometres to the south. There are major railroads, with access to the continental railroad networks, in both Lytton and Lillooet. Both communities would be the source of personnel for exploration or operations.

The property is underlain by crown land. It is used by the local rancher for grazing cattle in the summer. It has been logged in the past for timber. This has left a network of logging roads to access most areas of the property. The land is steep, but there are several areas locally that should be sufficient for a mill site and tailings impound.

J. T. Shearer has initiated informal discussions with First Nations Bands resident near the property. These are the Lytton Bands (NNTC) and Kanaka Bar Band.

HISTORY

In 1941 the Victory Claim was staked on the ridge between Lалуwissin Creek and McGillivray Creek within the existing claims, according to Duffel and McTaggart, 1952. This is described to be located over the ridge located in the area where the previous work program of 2006 was done. It describes a northwesterly trending zone of faulting. There is a description of “inclusions” that are consistent with the body or bodies of altered volcanics seen in the 2006 mapping. It also mentions fine grained pyrite in a rusty fault zone.

In 1971 Cuda Resources, (Chisholm, 1972) did a copper soil geochemical and magnetometer study in the area of Lалуwissin Creek and Highway 12 and south. This is about a kilometre southwest of the grid of the 2006 work. Geological mapping of these areas was completed in August of 1972 by Asano (1972) for Colt Resources Ltd. (renamed from Cuda Resources). He has mapped generally northerly trending bands of altered Nicola volcanics in Mount Lytton Complex diorite. The volcanics show varying levels of epidote and chlorite alteration. He correlates the magnetic highs to patches of gossan. There are several zones of copper mineralization described. The copper geochemistry and magnetometer survey were contoured in a general northerly trend. There is a special correlation between copper in soils and magnetometer highs.

D.C. Malcolm undertook geological mapping of the McGillivray Creek basin in 1972 to 1976 for Acacia Mineral Development Corporation. Copies of his reports were not available to the author. The following is credited to Malcolm’s report dated March 14, 1980 as recounted in the report of Pezzot and White (1986):

“The main deposits occur at the summit of a ridge and along its flanks between elevations 4,500 and 5,000 feet. On the north side of the ridge a number of small hand trenches expose sheared and brecciated feldspar porphyry and altered limy volcanics. Five samples over an area 200 feet by 200 feet, averaged 0.42% copper.

A road has been built from McGillivray Creek to the lower part of the deposit on the north slope of the ridge. Trenches have been roughed out partly across the deposit at elevations 4,650 and 4,800 feet. On the south side of the ridge, 1,500 feet south of these trenches, chalcopyrite occurs with magnetite in old trenches and malachite stained feldspar porphyry forms a slide in a dry gulch. One picked sample assayed 0.37 oz. silver, per ton and 7.16% copper.

On the road, at elevation 3,300 feet, a porphyry dike was exposed. Chalcopyrite bearing limestone breccia float occurs near it.

Pyritic deposits occur over a large area east of the porphyry dikes and extend across the claims. Two outcrops have been sampled and assayed 0.095% and 0.15% copper.”

The area described by Malcolm is consistent with the area that was the focus of the 2006 study.

A geochemical program was completed in 1978 (White, 1978) for Acacia Minerals. This is centered in the same basin as the 2006 work program of Atocha. His conclusions read:

“The limonite gossans exposed in the southern portion of the survey area have a strong copper zinc geochemical expression which indicates they are part of a northerly trending mineralized zone. They are heavily pyritized appear to be associated with a series of andesites, dacites, limestone breccias and tuffs. A strong copper, lead, silver and zinc anomaly occurs at 9 / 60s - OE at the head of a small stream which is seeping an alumina-rich white powder.”

In 1983 Ryan Energy undertook an 80 line kilometre VLF – EM and Magnetics airborne survey (Pezzot and White, 1983) over the ACE 1 to 8 claims in the McGillivray Creek basin that was the area of focus of the 2006 work. The resulting magnetic lows were interpreted as:

“Two northwest-southeast trending magnetic lows are evident across the survey area. One follows a geologically defined fault across the southwest corner of the claims area. The second follows

McGillivray Creek. Terrain clearance effects across the valley formed by McGillivray Creek are not influencing the magnetic field intensity in this area and it is likely that another fault is present. A north-south trending magnetic high correlates with a mountain ridge on the east side of McGillivray Creek. No geological evidence of a lithology change is reported in this area. The magnetic data may be reflecting an unmapped facies change within the volcanic unit; possibly a dioritic phase or simply an increased content of higher magnetic susceptibility materials. A closed magnetic high located on line 20 immediately west of this ridge is likely an outlier of the same rock unit.”

The VLF EM from the 1983 report is reported as:

“The VLF-EM data is presented in profile form over the same topographic and geological base map used to illustrate the magnetic contours. The Seattle frequency data ... shows a subtle shift in the field intensity which correlates with the G.S.C. defined fault crossing the southwest corner of the survey area. In addition, the northwest-southeast trending belt of limestone is reflected as a slight conductivity increase. This response extends further south than the unit as indicated by D.C. Malcolm.”

In 1978 to 1984 a geochemical survey was initiated by Ryan Exploration, a division of U.S. Borax, and designed to provide geochemical data over the area considered to be the best target (Richards, 1984b and Malcolm, 1978). Results indicated several areas of anomalous values in copper and zinc.

Western Aero Data completed 80 line km of VLF-EM and Magnetics airborne survey.

To the north on Blustry Mountain, in 1987 Aerodat Ltd. of Mississauga, Ontario was commissioned by Kanged Resources to conduct an airborne geophysical survey over the property. This survey consisted of a low level, helicopter supported programme which included a frequency VLF electromagnetic system, a high sensitivity caesium vapour magnetometer. Results of this survey were used to control the grid placement for a 1987 soil sampling programme conducted by Mark Management Ltd. (Gonzalez and Lechow, 1987).

In 1987 Mark Management Ltd. on the Blustry Mountain Property under the direction of Archean Engineering conducted a soil geochemical survey over a grid area of 900m x 100m in size. A total of 349 samples were collected and analyzed by Chemex Labs Ltd. using an ICP geochemical analytical technique. In general, anomalous values for Au, Ag, As, Cu, Hg, Mo, Sb, Pb and Zn outlined an open ended zone 650m long by 220m wide (Gonzalez and Lechow, 1987).

In 2003, Wyn Development completed geological mapping, prospecting geochemistry and detailed Induced Polarization (IP) on the nearby Blustry Mountain Property. Numerous drill targets have been selected based on the geology and IP results. Wyn was not able to negotiate with the Fountain Indian Band to address First Nation concerns on the Blustry Property.

Previous Geophysics

Several different airborne geophysical surveys were flown by the Geological Survey of Canada during the late 1960's and early 1970's, over ground which includes the McGillivray Property. The line spacings were somewhat broad and the instrumentation (non-digital) not as refined or precise as those currently available, but the data is, nonetheless, of very good quality.

Some very distinct patterns are apparent in the reprocessed data. Most obvious are the linear trends between positive and negative magnetic anomalies, which reflect the pattern of northwesterly and north-easterly trending faults in this area of the Cordillera. In addition it is clear that regional geochemical anomalies in pathfinder elements are often found in drainages which have their source in areas of moderate, negative magnetic relief. It is possible that ground geophysical surveys, properly managed, would be a useful exploration tool.

The 1983 Aeroborne Survey (Pezzot and White) document several magnetic lows correlated with major fault zones.

To this end a detail 3D IP survey was completed in the spring of 2004 and 2005, the results of which are documented in separate reports, Pezzot (2004) and S. J. Visser, 2005 on the nearby Blustry Mountain Property.

The survey was configured as a 3-D array with current and potential electrodes located on adjacent survey lines, spaced at 100 metre intervals. This configuration allows for the application of 3-D interpretation techniques, including 3-D inversion algorithms.

Combinations of resistivity and chargeability characteristics have outlined 3 distinct geological regimes across the survey area. A large portion of the northeastern corner of the grid (Lines 1600N – 2400N) is covered by a thin (50m thick) cap of highly resistive material. This overlies a 100m thick layer of highly variable material that includes several pods of extremely conductive and chargeable material. Basement rocks in this area appear to be relatively uniform, exhibiting low resistivity and elevated chargeability. The second regime is mapped from 1500N to 900N. It is also characterized with a resistive cap which often occurs as two or more thin layers. The underlying rocks exhibit low resistivity and low chargeability and contain a few isolated anomalies. The third regime covers the southwest corner of the grid. It is characterized by scattered zones of variable chargeability and resistivity in the top 75 metres. At depth the geophysical responses become more uniform and reveal two structural trends: N15°W and N45°E.

There are several lineations and trends that are mapped as abrupt discontinuities of particular geophysical parameter. These are likely representing sharp geological contacts or fault zones. There are several pods of extremely high resistivity that can be interpreted as areas of silica flooding. Several pods of anomalously high chargeability have been identified that could represent disseminated sulphide mineralization.

A program of prospecting and sample collection (and XRF assaying) was completed in 2014. Eleven representative samples were collected along the main access road (see locations on Figure 10). Samples M-1, M-3, M-4 and M-7 are examples of the highly leached volcanic often exhibiting box-works limonite textures. Assays suggest that leaching is variable. M-7 has the highest silica.

Exploration 2014

Assays were conducted by using an XRF Unit factory calibrated (Cert No. 0154-0557-1) on October 30, 2013, Instrument #540557 Type Olympus DPO-2000 Delta Premium. The instrument was calibrated using Alloy Certified reference materials by ARM1 and NIS5 standards. Only certified operators were employed and that were experienced in XRF assay procedures. Read times were 120 seconds or greater.

Samples M-5, M-6, M-9 and m-10 show the highly altered but less leached characteristics of the original andesitic and dacitic host rocks. Assays suggest, due to relatively low silica values, that the original rocks are not dacitic.

Samples M-2 and M-8 are less altered host rocks. Assays suggest that the rocks are very phosphate-rich with abundant iron and sulfur, M-2 has the highest Aluminum.

Sample M-11 is a chloritized and kaolinitic but otherwise relatively fresh feldspar-quartz porphyry. Assays suggest low iron content, aluminum lower than expected but the silica is higher.





Figure 5 2009 Soils Grid (Au/Ag/Cu)



Previous (2006) Soil Samples Location of 2009 Work.

Figure 6 2006 Soil Samples

Mineralization




Figure 7
McGillivray Cliff exposure looking northwesterly. Highly pyritized volcanic rocks consisting predominately of siliceous andesite with subordinate dacitic to rhyolitic lenses. Interpreted by author as part of the Cadwallader Island arc volcanic terrane.
D.G. Cardinal, P.Geo.

The mineralization is largely disseminated and shear related copper and silver - lead with some, zinc reported. There is gold reported as a possible metal in the copper porphyry deposits described in the BC Minfile property descriptions on this property near the ridge. Strong lineations were seen on the ground during the property visit and are also visible in the contour maps. These are probable fault boundaries to the altered volcanic units with the Mount Lytton Complex intrusives as described in several historical reports. The high level and large surface extent of alteration seen indicates strong hydrothermal alteration. This alteration was evident as the author walked the property as well as seen in the large landslide visible from a distance near the highway.

Styles of Mineralization

Several types of mineralization were identified and described by Richards (1984b). Quartz breccias with quartz crystal lined vugs and intense silicification of included wallrock have been noted in float. Sulphide content is generally less than 1% or 2% but tetrahedrite, galena and other silver coloured sulphides have been recognized with fine grained pyrite.

A second type of silica flood occurs as dark grey quartz veins in parallel bands, commonly 2mm wide but in places attaining a width of several centimetres. These compose as much as 70%, but on average 10%, of rock volume. This mineralization is developed in an area 50 to 100m wide and 200 to 300m long.

A third type of silicification occurs in rhyolite breccia with moderate clay alteration and less than 3% void space. The rhyolite breccia contains local zones with silicified fragments and with grey quartz partly filling the vugs. Silica flooding also occurs within the rhyolite and is accompanied by intense clay alteration.




Figure 8

Airphoto depicts interpretation, both from preliminary filed surveys and photos, a NW trending structural trace of accretionary collision of Cadwallader and Spences Bridge terranes with related first-second order structures and potential epithermal signatures. Silica-healed breccia-fragmented alkalic volcanic rocks outlined above are interpreted to be tectonic-accretion related.




Figure 9

Area interpreted as underlain by Spences Bridge terrane containing alkalic compositional rocks. Yellow dashed line indicates zone of carbonate enrichment as showing by large calcite veining possibly suggesting cooling fringe of epithermal event and deposition of carbonatized-rich fluids. Rock north of this zone shows little alteration and no tectonic breccia fragmentation. Gill 4-8 are bulk sampled sites collected by prospector for panning. Gill 5 is in the approximate location of previous soil sample with elevated Au of 390 ppb. Here, panned concentrate contain at least one very fine crystalline gold with a fine grain of electrum/telluride (?). Zone of silica-healed fragment/breccia volcanics is interpreted to be result of tectonic-accretionary collision with subsequent introduction of epithermal silica into the structural system.

Geochemistry and Trenching 2009

The geological examination of the ridge section show that the rocks are predominately composed of underlying, mildly altered siliceous andesite carrying 2-4% disseminated pyrite. No other sulphides were observed. The andesite is cut by series of roughly east-west trending second and third order faults. Within some of these structures are well silicified, bleached, carbonatized and, what appears to be alunite alteration. Trenching exposed a mineral assemblage associated with epithermal environments. The anomalous Ag and Cu values previously found are further defined by soil and rock samples exposed by trenches into fresh rock. Additional trenching is warranted to follow the mineralized structure on the ridge saddle.

A mapped thrust fault may have also acted as a channel way to ascending mineral-bearing solutions altering the andesitic rocks observed along the escarpment, with the cross-cutting, east-west trending second and third order

faults hosting epithermal, calcite-silica-alunite-bearing minerals. The ubiquitous pyrite associated with the andesite and concentrated mainly between the ridge escarpment and McGillivray creek to the east may also be spatially reflecting some distal epithermal system. As demonstrated by the highly iron-oxidized escarpment, that the disseminated pyrite, anomalous copper and silver and alteration minerals observed along the ridge are structurally controlled.

To the northeast, a new area of anomalous gold-in-soil results was found. Panned concentrate of soil collected near the site of the anomalous gold value contained at least one very fine crystalline gold flake along with a silvery grain believed to be electrum or telluride.

Bedrock observed along this area is composed of purplish coloured, alkali composition volcanic rocks associated with fine grain, creamy feldspathic phenocrysts. In some sections the volcanic rock appears as trachytic texture. In the area of the elevated gold-in-soil values the volcanic rocks are highly fragmented which is interpreted to be result of tectonic action. The fragments have been subsequently healed by banded white and pearl-white quartz veinlets, fracture-filling colliform silica and large bands of massive, dark, siliceous incipient-like chalcedony.

Trenching and follow-up soils sample results are plotted on Figure 8 (following page 13). A sample of sheared and rehealed volcanic assayed 1.0 ppm Ag and 2029 ppm Copper.

On the west end of the west trench there are anomalous silver and copper values (samples MG-West 1+2 and MG-W 1+2+3). This area requires further follow-up work.

Previous Exploration

The 2006 work program consisted of prospecting and soil/rock sampling. A total of 453 soils and 40 rock samples were collected in 2006. Silver appears to be anomalous in two sub-parallel zones with a central area low in silver content.

Reconnaissance soil sampling along the ridge shows highly anomalous silver in soils with values up to 42.0 ppm Ag. Anomalous values in Pb, Cu, Mo, and As.

Banded silicified zones were discovered at lower elevations which may be related to through-going fault zones.

Past producing deposits in the area are generally restricted to the Highland Valley porphyry deposits associated with granodioritic intrusive rocks of the Late Triassic to Early Jurassic Guichon Creek Batholith at the southeastern edge of the area.

The only other past producer in the general area is the Blackdome low-sulphidation epithermal gold deposit 96 km to the northwest. From start of production in April 1986, until the end of July 1990, the mine processed a total of 305,614 tonnes of ore yielding 6303 km Au and 19,518 kg Ag. This deposit is hosted by Eocene volcanic rocks of the type reported on the Blustry property; This deposit type is therefore to be targeted in the proposed exploration.

The abundance of regional geochemical data for the Ashcroft map sheet (0921) and for adjoining sheets to the north and west (BCGSB RGS 35, 36, 40, 41) permits a regional assessment for tracer elements appropriate to high and low sulphidation epithermal environments. The locations of regional stream sediment samples, including those which returned values in the top ten, five and two percent for the area's sample population in Au, Ag, As, Sb, Hg, and Mo. All are tracer elements for epithermal mineralization, among other types. All elements show an increase in anomalous samples in the vicinity of the McGillivray property, suggesting that the drainages samples cross rocks with elevated values of the elements. More comprehensive sampling in the vicinity of the property is necessary.

The work program in 2006 (Butler, 2007) included field grid development and soil and rock geochemistry. There was some geological mapping of the north facing bowl area that was the focus of the program. Another area of focus was around the rim of a large gossanous landslide that faces southwest. This landmark is clearly visible from a distance and was one of the reasons this property was located by Mr. Shearer.

The work included development of systematic lines of geochemical soil sampling along the ridge line and other geographic landmarks. A total of 453 soils and 40 rock samples were collected in 2006. A line along the ridge includes two samples with anomalous silver of 26 and 42 ppm. There are elevated values in lead in these two samples. The samples are located near a linear structure seen in the contour map. Several of the nearby samples are also elevated or anomalous in silver.

Prospecting of several other areas was completed to assess the outlying areas of the property.




Figure 10 2019 Rock Sample Locations and Results




Figure 11 2019 Magnetometer Traverses

Exploration 2019

Exploration at McGillivray in 2019 focussed on rock geochemistry at lower elevations and ground magnetometer along the main access road at mid elevations.

A total of 11 diverse rock samples were collected to characterize the variability of the lithology on the property.

Assays were conducted by using an XRF Unit factory calibrated (Cert No. 0154-0557-1) on October 30, 2013, Instrument #540557 Type Olympus DPO-2000 Delta Premium. The instrument was calibrated using Alloy Certified reference materials by ARM1 and NIS5 standards. Only certified operators were employed and that were experienced in XRF assay procedures. Read times were 120 seconds or greater.

Assay results are plotted on Figure 10. Values plotted on Figure 10 are Si%, Fe% and Cu ppm.

The magnetometer survey (Figures 16a to 16q) and values are contained in Appendix IV. The magnetic pattern varies from 2500 gammas in the lower elevations increasing to Figure 16L (Map #11) up to 2850 gammas, then decreasing in the upper area (southeast) back to 2550 gammas. The Figure 16L (Map #11) magnetometer anomaly likely reflects an increase in magnetite and sulfides and may be a good drill target.




Figure 12a Compilation of Survey Maps for 2019 McGillivray Magnetometer Survey




Figure 12b McGillivray Magnetometer Survey




Figure 12c McGillivray Magnetometer Survey




Figure 12d McGillivray Magnetometer Survey




Figure 12e McGillivray Magnetometer Survey




Figure 12f McGillivray Magnetometer Survey




Figure 12g McGillivray Magnetometer Survey




Figure 12h McGillivray Magnetometer Survey




Figure 12i McGillivray Magnetometer Survey




Figure 12j McGillivray Magnetometer Survey




Figure 12k McGillivray Magnetometer Survey




Figure 12L McGillivray Magnetometer Survey




Figure 12m McGillivray Magnetometer Survey




Figure 12n McGillivray Magnetometer Survey




Figure 12o McGillivray Magnetometer Survey




Figure 12p McGillivray Magnetometer Survey




Figure 12q McGillivray Magnetometer Survey

FIELD PROCEDURES

Sample locations (see Appendix III) were established using a topochain and Garmin GPS Unit at 10m and 25m spacing. The field data was downloaded to the Garmin MapSource program for plotting. The magnetometer used was a Sharpe MF1 Fluxgate instrument and diurnal variation was corrected by repeated readings at a base station.

Assays were conducted by using an XRF Unit factory calibrated (Cert No. 0154-0557-1) on October 30, 2013, Instrument #540557 Type Olympus DPO-2000 Delta Premium. The instrument was calibrated using Alloy Certified reference materials by ARM1 and NIS5 standards. Only certified operators were employed and that were experienced in XRF assay procedures. Read times were 120 seconds or greater.




Figure 13 Regional Geology

GEOLOGICAL SETTING

Regional Geology

The major rock formations on the property are dioritic and granodioritic intrusives of the Permian to Triassic age Mount Lytton Complex. The other major unit is the altered Lower Cretaceous andesitic volcanics of the Pimainus Formation of the Spences Bridge group. The Spences Bridge Group outcrops on the eastern side of the claims as well as fault controlled bands as inliers or roof pendants in the diorite on the western side. The 2006 program reportedly found sedimentary rocks on the eastern side of the property (Shearer pers. comm.). This is likely the interbedded volcanoclastic rocks of the Pimainus Formation.

There are bands of fault bounded northerly trending altered volcanics that have been mapped as gneisses and schists (Duffell and McTaggart, 1952). There are gneisses and schists defined to the south of this property on the geological map from the MapPlace, used in this report. Locally the alteration was observed to be argillic to kaolinitic. These bands extend over the ridge and were mapped near Highway 12 (Asano, 1972) as well in the basin to the east (Shearer, 2006). The intensity of alteration varies greatly on a local basis. These are likely part of the Pimainus Formation of volcanics of the Spences Bridge Group. The geological map reproduced from the BC MEMPR MapPlace reproduced for this report (Figure 4) does not show these bands of altered volcanics, but were observed during the field visit and reported in many property scale reports.

The regional Fraser Fault, a major north-north westerly trending structure, is located on the western boundary of the McGillivray property. This strike slip fault may have 135 to 160 kilometres of dextral strike slip. This was determined by the correlation of Late Permian intrusives of the Mount Lytton Complex in the area of McGillivray Creek with the Farwell Pluton in the area of the mouth of the Chilcotin River as noted in Read (2000) crediting a GSC paper by Friedman and van der Heyden. The rocks to the west of this structure, the Fraser Fault, are not related geologically to the units found on the McGillivray property and the geology and mineral deposit types are not reported by the author.

The close spatial relation to this fault has likely influenced the units on the McGillivray property. The strong northerly trending faulting that separates the Mount Lytton intrusives and the altered volcanics, sub parallel to this fault is likely related to this fault. As well, deep faults like the Fraser Fault have acted as conduits of deep hydrothermal fluids in other regions.

At this early stage of mapping there is field evidence to suggest to the author that a tectonic plate collision between 2 accreted terranes may occur in the McGillivray Property and that McGillivray creek valley may part of a surface expression to such a structural suture zone (Plate I).


Evidence to suggest a possible terrane collision proposal includes the following:

(i) Regional GSC map shows the McGillivray Property and area to be underlain mostly by the Lytton Complex. However, field mapping shows the Property, at least within the McGillivray watershed, to be predominately underlain by 2 different types of compositional volcanic rocks - andesitic and alkalic.

(ii) Regional geology along the Fraser River fault system shows faulted sections of the bedrock that has been part of the Cadwallader Group, an island arc terrane of Upper Triassic to Lower Cretaceous age that includes mafic to intermediate volcanic flows and younger fine clastic sediments. Sections of the Cadwallader sediments can be observed on Highway 12 consisting of mudstone, shale, and siltstone – along an area of the highway that is precariously unstable directly overlooking the Fraser River. To the south and on the Property - the ridge overlooking the highway, the rocks here are composed of, what the author believes, as part of the Cadwallader terrane, composed predominately andesitic and minor intermediate rocks.

(iii) North and east of the McGillivray creek are alkalic compositional volcanic rocks. These rocks are believed to part of the Spences Bridge terrane.

(iv) Fragmented alkalic volcanic rocks discussed above are believed to be result of tectonic activity related to an accretionary collision between terrane represented by the andesitic rocks to southwest and the alkalic volcanic rocks to the northeast.



McGillivray Creek Claims overlain on the geology, roads (red) and creeks/ivers (blue).

Claim numbers locate the claims (purple boundaries inside black property boundary)

FIGURE 14 Regional Geology

A geological map of the McGillivray Creek and surrounding areas is shown in Figure 5. It is based upon mapping carried out by Duffell and McTaggart (1952) and Trettin (1961); smaller studies by Mortimer (1987) and Read (1988a, 1988b, 1990) have augmented the broader regional mapping. The area was compiled as part of the Geological Survey of Canada's Terrane Assemblage Map by Monger and Journeay (1994).

The McGillivray property lies on the east side of the Fraser Fault, which experienced Eocene strike-slip movement of approximately 80km and which forms a geological boundary to the west. The basement to the area comprises rocks of the Permo-Triassic Cache Creek Complex, which are bounded to the southwest by granodioritic intrusive rocks of the Permo-Triassic Mount Lytton Complex. To the north of the study area, the Cache Creek assemblage is intruded by Late Jurassic granodioritic intrusive rocks associated with the Mount Martley and Tiffin Creek Stocks.

The McGillivray property is shown on Figure 3 to be underlain by calc-alkaline volcanic rocks of the Lower Cretaceous Spences Bridge Group in fault contact to the west with Lytton metamorphic complex. Outliers of the Eocene volcanic rocks assigned to the Kamloops Group occur to the east.

The Spences Bridge Group was previously not considered prospective for epithermal or other deposits, until the successful drilling in late 2005 by Strongbow discovered a promising intersection of 12.8m averaging 20.02g/tonne gold.

Regional structural geology in the area is not well defined. Brittle faults cross the property, with two prominent strike direction, parallel (northwesterly) and crudely perpendicular (north-easterly) to the structural grain of the Canadian Cordillera. Normal movement is apparent on several of the faults by the lateral juxtaposition of the Cretaceous volcanic rocks against older rocks.


Local Geology

The Company has received results for samples collected from the initial trenching program. Geological examination of the ridge section shows that the rocks are predominately composed of underlying, mildly altered siliceous andesite carrying 2-4% disseminated pyrite. Minor chalcopyrite was observed. The andesite is cut by series of roughly east-west trending second and third order faults. Within some of these structures are well silicified, bleached, carbonatized and appears to be alunite alteration. Trenching found associated with epithermal environments.

A thrust fault may have also acted as a channel way to ascending mineral-bearing solutions altering the andesitic rocks observed along the escarpment, with the cross-cutting, east-west trending second and third order faults hosting epithermal, calcite-silica-alunite-bearing minerals. The ubiquitous pyrite associated with the andesite and concentrated mainly between the ridge escarpment and McGillivray creek to the east may also be spatially reflecting some distal epithermal system. Nevertheless, it is obvious as noted by the highly iron oxidized escarpment (Photos 1-3), that the disseminated pyrite, anomalous copper and silver and alteration minerals observed along the ridge are structurally controlled.




To the northeast a new area of previously defined gold-in-soil results panned concentrate collected near the site of the anomalous gold value contained at least one (possibly 2) very fine crystalline gold flake along with a silvery grain believed to be electrum or telluride.

Bedrock observed along this area is composed of purplish coloured, alkali composition volcanic rocks associated with fine grain, creamy feldspathic phenocrysts. In some sections the volcanic rock appears as trachytic texture. In the area of the elevated gold value the volcanic rocks are highly fragmented which the author interprets to be result of tectonic action. The fragments have been subsequently healed by banded white and pearl-white quartz veinlets, fracture-filling colliform silica and large bands of massive, dark, siliceous incipient-like chalcedony.



Geology of the McGillivray Creek Property

PTrMdr	Permian to Triassic Mount Lytton Complex diorite
PTrMgd	Permian to Triassic Mount Lytton Complex granodiorite
PTrMml	Permian to Triassic Mount Lytton Complex metamorphic rocks
IKSBPva	lower Cretaceous Spences Bridge Group—Pimainus Formation volcanics

	Native Reserve
	Property Boundary
	Fraser Fault

Source BC MEMPR MapPlace

Figure 15 Detail Geology

A summary of general property geology (Richards, 1984b) is as follows:

Geological mapping is just starting to be done on a property scale for the area now covered by the McGillivray property. As noted above, regional mapping by the Geological Survey of Canada (Duffell and McTaggart, 1952) is over 50 years old and subsequent mapping by the British Columbia Geological Survey Branch (Mortimer, 1987; Read, 1988a, 1988b, 1990) did not cover the entire area.

Previous authors have noted that the McGillivray mineral claims are underlain by volcanic rocks of the lower Cretaceous Spences Bridge Group. This Group is composed mainly of an accumulation of lavas and pyroclastic rocks. Most of the lavas are porphyritic and are fine to coarse grained rocks of various colours. The colours are red, green mauve, purple, brown, grey, white and black.

In the vicinity of McGillivray Creek, dacites and minor rhyolites form part of the Spences Bridge Group and are intruded by a north-easterly trending dyke swarm of creamy pink, weakly feldspar hornblende phyric andesite. Gabbroic rocks intrude the volcanic sequence to the southwest of Blustry Mountain (Richards, 1984a, b) and a small plug of syenite, possibly a coarser grained equivalent of the pink feldspar-phyric dykes has been observed south of Cairn Peak.

The gossanous rocks south of McGillivray Creek shows a strong altered zone characterized by alunite with intense silica-kaolin alteration. Areas of vuggy porosity in silica matrix with kaolin are cut by fine stringers of translucent quartz. The vugs are normally lined with fine glassy quartz crystals. Some late stage quartz veins were also noted associated with occasional fine metallic lustre minerals – possible specularite. On the north side of the ridge hand trenches expose sheared and brecciated feldspar porphyry. Five samples over a 60m x 60m area averaged 0.42% Copper.

This section of the zone appears to have undergone a higher degree of silicification as evident by the quartz veining, suggesting several stages of silica flooding.

The alteration zone appears in part to represent a silica-clay cap of an epithermal system. The multi precious-base metal soil geochemical anomalies over the zone also support such an environment.

The coincidental geochemical anomalies and the intense silica-clay alteration zone may be pointing to near a surface precious metal-polymetallic epithermal deposit.

Basaltic volcanic rocks of the Kamloops Group are found to the east of the property, near Hat Creek. In Hat Creek valley, a thick section of sedimentary rocks is preserved in a graben that is floored by Eocene volcanic rocks.

Petrology

Zones of alteration are strongly controlled by structure. The most prominent structural trend is north-easterly while north-northwesterly trends also appear to have influenced the localization of alteration. These structural trends are thought to reflect Lower Tertiary translation and extensional tectonics that are well developed within this area.

The north-easterly trending dyke swarm is associated with a clay-sulphide zone that is developed over an area 4500 metres long and as wide as 1500 metres. Within the clay-sulphide zone are areas of silicification (silica flooding) which host precious metal and minor base metal mineralization.

Altered rocks from the Blustry Mountain area to the north of McGillivray Creek are dominated by vuggy silica/quartz alteration ± adularia ± Kaolinite ± possible alunite. The vuggy silica may be largely derived as a residual product of acid leaching. Quartz/silica forms a dense mosaic texture. Vuggy quartz alteration forms by reaction of extremely low-pH aqueous fluids or vapours with the host rocks. These fluids effectively remove all components in the rock apart from SiO₂ and TiO₂ leaving residual vuggy quartz. On the margins of this type of alteration zone, vuggy quartz may grade into quartz-alunite and quartz-kaolinite (or pyrophyllite) alteration. This change reflects the partial neutralization of the low-pH fluids during wall rock interaction. Low-pH fluids are commonly magmatic in origin and vuggy quartz alterations often form the cores of high-sulfidation precious metal systems. Sutured grain boundaries are common and suggest variable stress perhaps along nearby faults.

Kaolinite and dickite, (Al₂Si₂O₅(OH)₄), which are polymorphs occur in several specimens. The Kaolinite/dickite is mainly very fine grained anhedral, platy flakes. This mineral is indicative of formation at a pH of around 3 to 4 in the marginal argillic zone of high sulfidation systems (kaolinite forms under low-temperature conditions <150-200°C, whereas dickite at higher temperatures <200-250°C transitional to those for pyrophyllite formation). Sericite is commonly associated with kaolinite.

Possible fine grained alunite, (Na,K)Al₃(SO₄)₂(OH)₆, was tentatively identified in one sample, closely associated with fine grained kaolinite. Further work with a “PIMA” short wave infrared (SWIR) spectroscopy analyzer may be useful to define the presence of both kaolinite/dickite and alunite. Alunite is indicative of advanced argillic alteration and is often found in high-sulfidation epithermal precious metal systems. In this environment, magmatic SO₂ in the presence of water generates H₂S and H₂SO₄ which together with HCl react with host rocks to form zones of alunite-bearing advanced argillic alteration.

EXPLORATION 2020

Exploration at McGillivray in 2020 focussed on rock geochemistry at lower elevations and soil geochemistry along the main access road at mid elevations.

Assays were conducted by using an XRF Unit factory calibrated (Cert No. 0154-0557-1) on October 30, 2013, Instrument #540557 Type Olympus DPO-2000 Delta Premium. The instrument was calibrated using Alloy Certified reference materials by ARM1 and NIS5 standards. Only certified operators were employed and that were experienced in XRF assay procedures. Read times were 120 seconds or greater.

Assay results are plotted on Figures 16, 17 and 18 and XRF results are contained in Appendix III. Values plotted on Figure 16, 17 and 18 are Si%, and Cu%.

A total of 22 rocks and 51 soils were collected in the area of the “Bob” zone on claim 1075219. Results are plotted on Figures 17, 18 and 19 and Figures 20 and 21 on Claim 1074022 and soils were anomalous up to 407ppm Cu (Sample M20073). Rock samples returned very high results up to 8.09% Cu (Sample MR20023)

Soil sampling lines were run across prominent linears (fault zones/contact zones) as shown on Figures 20 and 21. The results indicate a marked change on either side of the linears.




Figure 16 Sample Location and Results Bob Zone




Figure 17 Sample Locations and Results Bob Creek




Figure 18 Sample Locations and Results Lower Bob Creek




Figure 19 Google Showing Eastern Samples by S. Butler




Figure 20 Google Showing Western Samples by S.Butler

In light of recent encouraging gold exploration activities along the northwestern extension of the Spences Bridge Gold Belt adjacent to the McGillivray claim group. The copper mineralization and alteration zones covered by the McGillivray claims - mainly focusing in the area of the Alice copper mineralization and historical workings and associated geological host rocks were re-examined in 2020.

In recent years the Spences Bridge volcanic belt – also referred to as the ‘Spences Bridge Gold Belt’, has been the focus of epithermal-related gold mineral exploration. Exploration surveys have identified epithermal type hydrothermal and alteration signatures along parts of the belt hosting gold-silver and copper mineralization.

Objective of this follow-up examination is to attempt to better constrain the lithological host rock units, possible genesis of mineralization, associated alteration characteristics and structural control associated with the Alice copper zone, and based on empirical field observations suggest a mineral exploration model as a target for future exploration surveys.

Based on the re-examination of the Alice copper zone it was observed that the alteration and structural characteristics associated with the zone do not appear to fit the epithermal-porphyry models being applied to other adjacent discoveries such as those found on the adjacent Bob, Spin, B&B and Cobra copper-gold zones.

The mineralized host rock, alteration and structural features associated with the copper zone may be somewhat more unique. The Alice copper zone requires re-interpretation and consideration taken to develop an exploration mineral model that will best fit more an ‘Alice type’, possibly a copper (silver)-bearing, iron carbonate(+quartz-calcite-kaolinitic) breccia (diatreme-like), volcanic hosted stratabound-related exploration model.

ALICE CLAIM:

The northern portion of the McGillivray claim group covered by: Alice (1061863), Alice McGill (1057727) and McGill (1061589) was re-examined. The Alice claim is one of the key mineral claims on the property covering an important copper-bearing, iron carbonate breccia zone herein referred to also as the ‘Alice copper zone’.

The Alice claim can easily be reached on Highway 12 some 30km south-southeast of the town Lillooet, BC. It is situated about half way between the towns of Lillooet and Lytton. The claim, as well as remaining claim group, is located east of and overlooks the Fraser River and highway. The Alice copper zone and historical workings occur just east of a Hydro powerline r/w, which parallels the highway, associated with an iron oxidized reddish coloured, well exposed rocky bluff. The author was able to easily access the copper showings and old workings by walking up an older exploration road which leads from the powerline r/w part way up the hill crossing some 50-60m below the Alice copper zone. The road switches back and forth across the hill cutting across bedrock affording detail examination but does not intersect the Alice copper zone due the steep rocky bluffs.

BEDROCK LITHOLOGY – ALICE HILL ROAD–SECTION:

Starting near the base of the hill adjacent to powerline r/w and leading up to the Alice copper zone, sections of well exposed bedrock were encountered along the exploration road noted above, which the author had the opportunity to examine in some detail. The dominate rock type observed is a variable: medium grain, grey to dark-grey diorite sections of which phase to lesser quartz diorite and localized feldspar porphyritic phases. In places it appears to be locally bleached, it characteristically displays pervasive choritic overprint forming weak to mild propylitic alteration consisting mainly of calcite-chlorite and minor epidote with iron oxidation along fractures and joints. Occasionally the diorite hosts minor, small blebs of disseminated malachite. Also commonly observed were steeply dipping, east-west striking, structurally controlled, hydrothermal quartz-calcite-iron carbonate (siderite-ankerite) veins cutting the dioritic rocks.

Along its southern section the exploration road cuts across a large talus slope, which forms part of Alice hill, comprised mainly of diorite-quartz diorite slide material with occasional quartz-calcite iron carbonate breccia talus carrying abundant disseminated malachite-azurite mineralization (see cover page). This mineralized talus is believed to be derived from the Alice copper zone further up slope. On the northern section of the hill road switch-backs abruptly terminate along an east-west trending, exposed bedrock of highly fractured oxidized dioritic rocks.



Photo 1: Exposed bedrock along Alice hill road-section showing one of several structurally controlled quartz-calcite iron carbonate veins observed striking about west-east cutting jointed and fractured, propylitic altered diorite –quartz diorite rocks (for scale handle of rock hammer is about 36cm long).

ALICE COPPER ZONE – IRON CARBONATE BRECCIA:

Further up slope overlying the diorite rocks discussed above is a highly altered rock unit that has undergone intense carbonitization, here the rocks significantly change. This altered, brecciated rock unit appears to be stratigraphically concordant and stratabound with the overlying andesitic volcanic rock unit and is host to the Alice copper zone. The contact between the diorite and the carbonitized stratabound, copper-bearing zone is masked by the talus material. Future follow-up mapping to the north and south along Alice hill may encounter an exposed section the contact horizon between the 2 rock units further constraining the Alice copper-bearing horizon.



Photo 2: Looking northwest showing compositional change in the andesitic strata and structural attitude of the folded volcanic rocks striking northwest dipping northeast, found near the old adit, part of the Alice copper zone workings located just to the right of the photo.

An andesitic flow layer was observed, which outcrops in at least 2 different areas on Alice hill, which appears to be the basement to the overlying iron carbonatized felsic breccia unit. The andesite forms a stratigraphic layer separating the diorite and the carbonate breccia and appears to directly overlie in contact the dioritic rocks. To the left of photo 2 (down slope) contact with the diorite is obscured by talus material. Up slope (right of photo) the upper section of the andesitic basement has experienced gradational alteration influenced by the overlying hydrothermally altered iron-rich carbonate felsic (rhyolitic) sequence, which hosts the Alice copper-bearing horizon. The folded andesite (photo 2) represents a deformational event (D1/D2), strikes 3000 dip 60-650 northeast. Snow covered valley upper right is part of the Fraser River represents a crustal break, part of the north trending Fraser Fault system.

Photo 3 below shows the old Alice adit at (GPS) elevation of about 666m, it occurs several metres east or right of photo 2. Photos 3 to 6 below depict part of the old Alice workings.



Photo 3: Old adit @ elev. 666 – part of Alice copper zone. Copper-bearing iron carbonate appears to be associated with altered felsic – rhyolitic unit. Dimensions of adit: 6m long x 1.25m wide (at the face) x 2.4m high. Black, broken lines bottom left trace altered volcanic-carbonate (dolomitic) strata, which strike and dip are similar to strata shown in photo 2.



Photo 4: Old open-cut/pit located approximately 20 metres south (left) of photo 3, cut into the iron carbonate breccia unit hosting malachite staining along exposed face.

The folded, altered volcanic strata in photo 3 above conforms with the andesitic strata in photo 2 and in the adit (photo 5) as part of the same carbonate altered stratigraphic sequence. This would suggest that the Alice copper- iron carbonate (dolomitic) breccia zone is part of copper-silver-bearing, hydrothermally altered felsic (rhyolitic) volcanic stratabound flow beccia layer, occurring between an andesitic volcanic formation. Also, approximately 100 metres up slope (east) from the old Alice copper workings, in 2011, the author during reconnaissance surveys mapped a thick sequence of mafic (andesitic-basaltic) volcanic rocks.




Photo 5: Structurally controlled, vertical dipping, copper-silver vein (yellow) associated with oxidized iron carbonate mineralization (red) ranging 10-40cm in width occurs on the face of the adit . A narrow (2-4cm), iron carbonate vein parallels the main mineralized vein both are about 40cm apart. The mineralized vein vertically cuts easterly dipping altered iron carbonate, quartz-calcite, kaolinitic volcanic strata .

Photo 6: Malachite-copper stained iron breccia boulders from old open-cut/pit (photo 4). Alteration minerals include brecciated quartz, calcite, iron carbonate –dolomitic (ankerite/siderite), includes kaolinitic and cryptocrystalline-like agate mineral fragments.





Photo 7: Close-up view of a typical mineralized, iron carbonate (dolomitic) breccia containing kaolinite-quartz-calcite and limonite fragments with occasional, smaller cryptocrystalline agate-like fragments.

MINERALIZATION – ALICE COPPER ZONE:

Based on the style of copper mineralization observed there appears to have been at least 2 mineralizing events. The earlier phase is associated with the iron carbonate breccia and appears the copper mineralization as indicated by the malachite breccia was introduced suggesting syn/pre-breccia hydrothermal mineral event. The latter phase is a post-brecciation mineral event, as indicated by the structurally controlled, epigenetic copper-silver sulphide mineralized vein found in the adit.

Samples selected from the mineralized vein and examined under binocular microscope display a metallic silver appearance (see photo 8 below). It is fairly soft and partly sectile under hardness test indicating tetrahedrite characteristics. BCGS (1980) briefly inspected the Alice claim and identified the mineral as 'Tennantite' a copper arsenic (zinc) sulfosalt. The current work suggests it may be closer to a tennantite-tetrahedrite composition with arsenopyrite. Other sulphides observed resemble secondary chalcite/enargite and peacock color probable bornite. This is also supported by sample collected by the author in May 2011 from the same vein and analyzed. The results include: 4.70% copper, 31.5ppm silver, 4,411ppm zinc, >1000ppm arsenic and 2,375ppm antimony .

ALTERATION – ALICE COPPER ZONE:

The copper (silver)-bearing iron carbonate-felsic breccia and associated rhyolitic breccia flows have undergone a high degree of hydrothermal alteration. Iron carbonatization is pervasive throughout the felsic-rhyolitic horizon and appears to be more intense where the felsic unit is highly brecciated. Alteration minerals include: iron carbonate (siderite/ ankerite), silica (quartz), carbonate (calcite), kaolin and smaller agate-like, white cryptocrystalline fragments (photo 9).



Photo 8: Sample collected adjacent to the adit entrance displaying intense iron carbonate-kaolinite altered breccia carrying malachite fragments.

STRATABOUND-LIKE COPPER (SILVER)-BEARING IRON CARBONATE BRECCIA:

Series of 3 photos below (10a,b&c) show a diatreme-like or chimney bedrock jutting upward mapped along the south-eastern slope of Alice hill exposing part of the stratabound iron carbonate breccia unit at (GPS) elevation 656m. Located about 75m southerly of the old adit workings discussed above.

ALICE HILL – STRATAGRAPHIC SECTION:




Figure 21 Alice Hill – Stratigraphic Section

CONCLUSIONS and RECOMMENDATIONS

The McGillivray Project, centred around McGillivray Creek in south-central British Columbia, represents a potentially large belt of underexplored, poorly understood volcanic rocks, of Cretaceous Spences Bridge Group and similar in structure, alteration and mineralization to those hosting the former producing Blackdome mine to the north and the Skoonka Creek Zones. Anomalous precious metal values are associated with later stage silica flooding/stockwork veinlets which cut felsic volcanic rocks. The altered volcanic system which contains this system extends over several kilometres. A methodical approach of detailed structural mapping and sampling, would define the geological controls on the existing anomalies. Prospecting and regional sampling of more remote areas with polymetallic anomalies in the regional geochemical survey might well define new areas of prospective mineralization.

A very strong through-going structure, possibly reflecting a major terrane boundary, trends approximately east-west touching mid McGillivray Creek. Strong parallel structures occur to the south along lower Luluwissan Creek and bounding the central crustal blocks and may control the emplacement of intrusive elements in the Lytton Metamorphic Complex.

A prominent splay to the southeast can be observed crossing from the McGillivray Valley into the mid Luluwissan Valley and beyond. A series of lesser linears oriented NE and NW are evident in the north fork of McGillivray Creek associated with normal faults in the upper Hat Creek Valley system.


A program of prospecting and sample collection (and XRF assaying) was completed in 2014. Eleven representative samples were collected along the main access road (see locations on Figure 10).

A total of 22 rocks and 51 soils were collected in the area of the “Bob” zone on claim 1075219. Results are plotted on Figures 17, 18 and 19 and Figures 20 and 21 on Claim 1074022 and soils were anomalous up to 407ppm Cu (Sample M20073). Rock samples returned very high results up to 8.09% Cu (Sample MR20023)

Soil sampling lines were run across prominent linears (fault zones/contact zones) as shown on Figures 20 and 21. The results indicate a marked change on either side of the linears.

An initial Phase I consisting of prospecting and soil sampling was carried out during the latter part of 2006 (Shearer, 2006). An additional period of 84 man days in the field is recommended, in addition to time expended in preparation and in report writing. The purpose of the fieldwork will be to re-establish a grid in the central area of the property and resample certain areas, predominantly those locations from which samples were anomalous as well as to expand the sampling to other mineralized zones. Silt sampling and prospecting of all drainages should be undertaken to aid in locating new or hidden targets. Coincident with the sampling, a programme of geological mapping will prioritize location of alteration, rock units and structures controlling or channelling the mineralizing fluids and upon establishing the limits of the gold-bearing mineralization. To this end, it is recommended that preparations for the field include facilities for staining to detect potassium in altered samples and also rental of a PIMA unit to expedite mapping of the alteration and mineralization. The budget for Phase I is estimated at \$210,000 as follows. (see next page)

Respectfully submitted,



J. T. Shearer, M.Sc., P.Geo. (BC & Ontario)

Cost Estimate of Future Work

Phase I

Phase I programme at \$210,000 should consist of more detailed mapping, sampling, and expansion of anomalous zones, and IP geophysics followed by contingent diamond drilling if warranted. Phase II budget is set at \$249,000 as follows.

Senior Geologist	42 days @ \$600/day	\$ 25,200.00
Geotechnician	42 days @ \$400/day	16,800.00
Geotechnician	42 days @ \$300/day	12,600.00
Labour	42 days @ \$250/day	10,500.00
Management Fee, WCB, Office and Overhead @ 10%		6,510.00
IP Geophysics		40,000.00
Equipment Rental		
(2) 4x4 Trucks	42 days @ \$75/day	3,150.00
(2) 4-Trax	42 days @ \$50/day	2,100.00
Camp @ \$3,000/month		4,500.00
(2) PIMA Geophysics Instrument @ \$500/month		4,000.00
GST 6%		7,521.00
Excavator Trail Building		19,119.00
Excavator Trenching		9,000.00
Petrographic Work		5,000.00
Food and Fuel, Mob/Demob		3,000.00
Assays	1600 samples @ \$15/sample	21,000.00
Field Supplies (pickets, tags, sample bags, flagging, etc.)		3,000.00
Preparation and Report Writing		8,000.00
Contingency @ 10%		9,000.00
TOTAL – Phase I		\$ 210,000.00
Phase II: Contingent Diamond Drilling		
Diamond drilling (1000m @ \$75/m all in)		\$ 150,000.00
Geological Mapping		30,000.00
Assays		14,000.00
Support, Camp, Supplies		30,000.00
Contingency		25,000.00
GRAND TOTAL – Phase II		\$ 249,000.00

REFERENCES

Allen Guy, 1988:

Prospecting Report on the Laurie Mineral Claim, Mines and Petroleum Resources Assessment Report 18133
1989:

Geological Report on the Laurie Mineral Claim (Rec, No, 7411) Kamloops Mining Division, Ministry of Energy, Mines and Petroleum Resources Assessment Report 18806

Asano, Edward, 1972

Geological Report on the Colt Resources Ltd. Property, Ministry of Energy, Mines and Petroleum Resources Assessment Report 04455

British Columbia Survey Branch, 1999-2003:

The Map Place: www.em.gov.bc.ca/Mining/Geosurv/MapPlace

Minfile: www.em.gov.bc.ca/Mining/Geosurv/minfile/

British Columbia Geological Survey Branch and the Geological Survey of Canada, 1980-1992:
British Columbia Regional Geochemical Survey Program, NTS 92O – Taseko Lakes, BC: RGS 35.

British Columbia Geological Survey Branch and the Geological Survey of Canada, 1980-1992:
British Columbia Regional Geochemical Survey Program, NTS 92P – Bonaparte Lake, BC: RGS 36.

British Columbia Geological Survey Branch and the Geological Survey of Canada, 1982-1994:
British Columbia Regional Geochemical Survey Program, NTS 92I – Ashcroft, BC: RGS 40.

British Columbia Geological Survey Branch and the Geological Survey of Canada, 1981-1994:
British Columbia Regional Geochemical Survey Program, NTS 92J – Pemberton, BC: RGS 41.

Butler, S.P., 2020

Technical Report on the McGill Property for Prisma Capital Inc., dated December 2020

Cardinal, D.G., 2020

Brief Summary Report Update on the McGillivray Claim Group, Alice Copper Mineralization and Alteration Zone, dated November 2, 2020

Chang, F and Gale, D., 2006

Report on Exploration Activities, Prospecting, Mapping, Geochemistry, Geophysics and Drilling, Skoonka Creek Property for Strongbow Exploration Inc., Ministry of Energy, Mines and Petroleum Resources Assessment Report 28182

Chisholm, Edward O, 1971

Qualifying Report on the Lillooet Property, for Cuda Resources, Ministry of Energy, Mines and Petroleum Resources Assessment Report 03154

Duffell, S. and McTaggart, K. C., 1952:

Ashcroft Map Area, British Columbia (92I/NW). Geological Survey of Canada Memoir 262.

Gonzalez, R. A. 1987:

Geochemical Assessment Report on the Top Hat Property, Kamloops Mining Division, Ministry of Energy, Mines and Petroleum Resources Assessment Report 16,352B

Jones, H.M., 1972a

Geochemical Report on the Spot No 1; COP Nos 1, 2,25 to 30; Ken 2; Kent 1, 2 and Alpine 1-8 Claims, Part of the East Group, Ministry of Energy, Mines and Petroleum Resources Assessment Report 03937

1972b

Geochemical Report on the CB Nos 1 – 9 and REB Nos 1 – 4 Claims, Part of the West Group Izman Creek Area, Ministry of Energy, Mines and Petroleum Resources Assessment Report 03938

Lechow, W., 1987:

Report on Combined Helicopter Borne Electromagnetic Magnetic and VLF EMI Survey, Top Hat Property, Lillooet Area, Ministry of Energy, Mines and Petroleum Resources Assessment Report 16,352A

Malcolm, D. C., 1978:

Soil Sampling and Geology on the Acecia Property, McGillivray Creek Area.

Monger, J. W. H., and Journeay, J. M., 1994:

Guide to the Geology and Tectonic Evolution of the Southern Coast mountains. Geological Survey of Canada Open File 2490, 77p.

Mortimer, N. 1987:

Geological Map of the Pavillion Map Area, British Columbia. Geological Map of Part of NTS92I/13. British Columbia Geological survey Branch Open File 1987-18.

Mortimer, N. 1987:

Geological Map of the Pavilion Map Area, British Columbia. Geological Map of Part of NTS92I/13. British Columbia Geological Survey Branch Open File 1987-18.

Panteleyev, A., 1988:

A Canadian Cordilleran Model for Epithermal Gold-Silver Deposits. *In*: Roberts, R. G. and Sheahan, P.A. (eds.) Ore Deposit Models. Geoscience Canada Reprint Series 3, pp. 31-43.

1992:

Copper-Gold-Silver Deposits Transitional, Between Subvolcanic Porphyry and Epithermal Environments. *In*: Geological Fieldwork, 1991, British Columbia Ministry of Energy, Mines and Petroleum Resources, Paper 1992-1, pp. 231-234.

Pezzot, E. T., 2004:

Assessment Report on a 3-D Induced Polarization and Resistivity Survey, July 2004, 21pp and maps.

Pezzot, E. T. and White, G. E., 1983:

Airborne VLF-EM and Magnetometer Survey on the Ace 1-8 Claims, Assessment Report 11,371

Read, P. B., 1988a:

Tertiary Stratigraphy and Industrial Minerals; Fraser River, Lytton to Gang Ranch. British Columbia Geological Survey Branch Open File 1988-29a.

1988b:

Tertiary Stratigraphy and Industrial Minerals, Fraser River, Lytton to Gang Ranch. British Columbia Geological Survey branch Open file 1933-2b.

1990:

Cretaceous and Tertiary Stratigraphy and Industrial Minerals, Hat Creek, Southern B.C. British Columbia Geological Survey branch Open File 1990-23.

2000:

Geology and Industrial Minerals of the Tertiary Basins, South-Central British Columbia, British Columbia Geological Survey branch, Geofile 2000-3.

Richards, G. G., 1984a:

Geochemical Report on the Top Hat Property, Top Hat 1-4 Mineral Claims, Kamloops Mining Division. British Columbia Ministry of Energy, Mines and Petroleum Resources Assessment Report 12,948.

1984b:

Geochemical Report on the Rusty Property, Rusty 1-2 Mineral Claims, Lillooet Mining Division. British Columbia Ministry of Energy, Mines and Petroleum Resources Assessment Report 12,944.

Shearer, J.T. 2006:

Geochemical, Geological and Prospecting Assessment Report on the McGillivray Project, Lillooet, Kamloops Mining Division, November 30, 2006

2013:

Airphoto Assessment Report on the McGillivray Property, dated February 1, 2013 for Atocha Resources Ltd.

2014:

Geological and Geochemical Assessment Report on the McGillivray Project, dated July 12, 2014 for Homegold Resources Ltd.

Trettin, H. P., 1961:

Geology of the Fraser River Valley Between Lillooet and Big Bar Creek. British Columbia Geological Survey Branch Bulletin B44.

Weymark, William J., 1980

Geological – Geochemical – Geophysical Surveys of the Laurie Mineral Claims Group, British Columbia Ministry of Energy, Mines and Petroleum Resources Assessment Report 8762

White, Glen E., 1978

Geochemical Report Acacia Mineral Dev. Corp., Kamloops Mining Division, British Columbia Ministry of Energy, Mines and Petroleum Resources Assessment Report 07027

White, G.P.E., 1980

Geological Fieldwork: A Summary of Field Activities and Current Research, 1980, British Columbia Ministry of Energy, Mines and Petroleum Resources, Pg 115-116 A Tennantite Occurrence McGillivray Creek -Lytton Lillooet Area.

APPENDIX I

STATEMENT of QUALIFICATIONS

November 17, 2020


STATEMENT OF QUALIFICATIONS

I J. T. (Jo) Shearer do hereby certify that:

1. I am a consulting geologist and principal of Homegold Resources Ltd.
2. My academic qualifications are:
 - Bachelor of Science, Honours Geology from the University of British Columbia, 1973
 - Associate of the Royal School of Mines (ARSM) from the Imperial College of Science and Technology in London, England in 1977 in Mineral Exploration
 - Master of Science from the University of London, 1977
3. My professional associations are:
 - Member of the Association of Professional Engineers and Geoscientists in the Province of British Columbia, Canada, Member #19,279
 - Fellow of the Geological Association of Canada, Fellow #F439
 - Fellow of the Geological Society of London
 - Fellow of the Canadian Institute of Mining and Metallurgy, Fellow # 97316
 - Fellow of the Society of Economic Geologists (SEG), Fellow #723766
4. I have been professionally active in the mining industry continuously for over 45 years since initial graduation from university.
5. I am responsible for the preparation of all sections of the technical report entitled "Geological and Geochemical Assessment Report on the McGillivray Property" dated November 17, 2020. I have most recently visited the Property between June 15 and October 30, 2020 and May 11 and 12, 2019, but also in the past in May 18 and 19, 2012, June 29 and 30, 2014 and July 10 and 11, 2014. General geological parameters were also examined.

Signed and dated in Vancouver B.C.

November 17, 2020
Date



J.T. (Jo) Shearer, M.Sc., P.Geo. (BC & Ontario) FSEG

APPENDIX II

STATEMENT of COSTS

November 17, 2020

Statement of Costs McGillivray Project 2020

Wages	Total without GST
Senior Geologist J.T. Shearer, M.Sc., P.Ge., 12 days @ \$800/day, Oct. 19-23 + Oct. 26-Nov. 1, 2020	\$ 9,600.00
Senior Geologist D. Cardinal, P.Ge., 6 days @ \$800/day, Oct. 21-26, 2020	4,800.00
Geologist, Sean Butler, P.Ge., 8 days @ \$700/day, Oct. 19-27, 2020	5,600.00
Field Assistant John Grabavac, 12 days @ \$350/day, Oct. 19-23 + Oct. 26-Nov. 1, 2020	4,200.00
Subtotal	\$ 24,200.00
Expenses	
Transportation:	
Truck 1 Rental, fully equipped, 8 truck days @ \$125/day	1,000.00
Truck 2 Rental, fully equipped, 12 truck days @ \$125/day	1,500.00
Fuel	800.00
Side-by-side & trailer, 12 days @ \$150/day	1,800.00
Food	600.00
Meals	1,500.00
Hotel	2,500.00
XRF Operator	800.00
XRF Rental and Assays	600.00
Assays – ALS Canada Invoice # 5320675 & 5320759	2,285.06
Mapping and Computer Compilation	1,200.00
Report Preparation	1,800.00
Word Processing	400.00
Subtotal	\$ 16,785.06
Total	\$ 40,985.06

Event #	5818069
Date	November 17, 2020
File	\$ 40,000.00
PAC Debit	\$ 7,863.94
Total	\$ 47,863.94

APPENDIX III

SAMPLE DESCRIPTIONS

November 17, 2020

McGillivray Sample Locations 2020

UTM NAD83 Zone 10

Sample Number	Type	Waypoint	UTM E	UTM N	Elev from GPS	Description
M-20-001	Soil	093	599830	5589985	1235	
M-20-002	Soil	094	599835	5590017	1238	
M-20-003	Soil	095	599869	5590046	1244	
M-20-004	Soil	096	599899	5590079	1245	
M-20-005	Soil	097	599919	5590113	1259	
M-20-006	Soil	098	599933	5590126	1236	
M-20-007	Soil	099	599966	5590165	1246	
M-20-008	Soil	103	597604	5591307	1272	
M-20-009	Soil	104	597644	5591310	1307	
M-20-010	Soil	105	597691	5591306	1331	
M-20-011	Soil	106	597741	5591298	1349	
M-20-012	Soil	107	597790	5591281	1379	
M-20-013	Soil	108	597854	5591278	1400	
M-20-014	Soil	109	597888	5591265	1410	
M-20-015	Soil	110	597764	5591008	1396	
M-20-016	Soil	111	597679	5590938	1362	
MR-20-001	Rock	098	599933	5590126	1236	fg, volcanic, dark green
MR-20-002	Rock	111	597679	5590938	1362	fg, volcanic, dark green, fg sulfides
MR-20-003	Rock	111	597679	5590938	1362	fg, volcanic, dark green, rusty vugs after sulfides
MR-20-004	Rock	111	597679	5590938	1362	fg, volcanic, dark green, fg Py on fractures

Grid UTM
Datum WGS 84

Waypoint	Date	Position	Altitude	Cu ppm	Si%	Description
BOB	30/10/2020	10 U 598890 5587200	1612 m			Pyrite mineralized outcrop at top of hill.
BOBADIT	31/10/2020	10 U 598791 5586974	1668 m			Adit covered with a lattice of logs. 40cm wide mineralized quartz vein with chalcopyrite and malachite exposed by BOBADIT.
BOBTRUCK	30/10/2020	10 U 599163 5587152	1613 m			Open field. Motorcycle trail 100m southwest along roadway goes right up to BOB.
M20030	29/10/2020	10 U 600216 5586417	1433 m	49	14.56	Soil
M20031	29/10/2020	10 U 600166 5586459	1434 m	95	13.48	Soil
M20032	29/10/2020	10 U 600119 5586487	1426 m	52	11.45	Soil
M20033	29/10/2020	10 U 600055 5586461	1415 m	4	16.06	Soil

Waypoint	Date	Position	Altitude	Cu ppm	Si%	Description
M20034	29/10/2020	10 U 600022 5586450	1408 m	85	14.44	Soil
M20035	29/10/2020	10 U 599968 5586476	1396 m	41	13.34	Soil
M20036	29/10/2020	10 U 599915 5586475	1388 m	66	12.89	Soil
M20037	29/10/2020	10 U 599887 5586466	1381 m	65	16.81	Soil
M20038	29/10/2020	10 U 599839 5586455	1374 m	47	14.03	Soil
M20039	29/10/2020	10 U 599801 5586439	1367 m	29	15.66	Soil
M20040	29/10/2020	10 U 599762 5586424	1364 m	41	16.69	Soil
M20041	29/10/2020	10 U 599718 5586417	1358 m			Soil-no results
M20042	29/10/2020	10 U 599672 5586367	1352 m			Soil-no results
M20043	29/10/2020	10 U 599631 5586333	1345 m			Soil-no results
M20044	29/10/2020	10 U 599593 5586295	1337 m	47	18.84	Soil
M20045	29/10/2020	10 U 599662 5586277	1341 m	131	15.74	Soil
M20046	29/10/2020	10 U 599564 5586273	1318 m	4	18.34	Soil
M20047	29/10/2020	10 U 599516 5586289	1313 m	142	14.86	Soil
M20048	29/10/2020	10 U 599457 5586246	1306 m	112	16.32	Soil
M20049	29/10/2020	10 U 599413 5586259	1298 m			Soil-no results
M20050	29/10/2020	10 U 599372 5586257	1294 m	118	15.41	Soil
M20051	29/10/2020	10 U 599330 5586269	1285 m	152	15.84	Soil
M20052	29/10/2020	10 U 599280 5586255	1286 m	206	14.68	Soil
M20053	29/10/2020	10 U 599250 5586242	1278 m	103	16.47	Soil
M20053	29/10/2020	10 U 599250 5586242	1278 m	97	16.42	
M20054	29/10/2020	10 U 599191 5586257	1270 m	106	14.5	Soil
M20054	29/10/2020	10 U 599191 5586257	1270 m	91	16.28	
M20055	29/10/2020	10 U 599141 5586254	1263 m	73	17.35	Soil
M20056	29/10/2020	10 U 599091 5586248	1251 m	196	11.55	Soil
M20057	29/10/2020	10 U 599037 5586240	1243 m	174	13.31	Soil
M20058	30/10/2020	10 U 600244 5586392	1417 m	55	17.04	Soil
M20059	30/10/2020	10 U 599025 5587187	1621 m	51	21.46	Soil
M20060	30/10/2020	10 U 598965 5587253	1630 m	85	12.39	Soil
M20061	30/10/2020	10 U 598912 5587207	1646 m	234	9.96	Soil
M20062	30/10/2020	10 U 598913 5587093	1669 m	251	10.89	Soil
M20063	30/10/2020	10 U 598980 5587055	1663 m	65	15.09	Soil
M20064	30/10/2020	10 U 598996 5587027	1660 m	15	13.25	Soil
M20065	30/10/2020	10 U 598978 5586990	1655 m	196	14.2	Soil
M20066	30/10/2020	10 U 598913 5587065	1666 m	222	12.25	Soil
M20067	30/10/2020	10 U 599055 5586985	1634 m	115	14.38	Soil
M20068	31/10/2020	10 U 600146 5587229	1627 m	67	14.3	Soil
M20069	31/10/2020	10 U 599135 5587115	1618 m	72	15.21	Soil
M20070	31/10/2020	10 U 599134 5587057	1621 m	66	15.5	Soil
M20071	31/10/2020	10 U 599086 5587017	1638 m	157	13.68	Soil
M20072	31/10/2020	10 U 599054 5587033	1649 m	147	12.47	Soil

Waypoint	Date	Position	Altitude	Cu ppm	Si%	Description
M20073	31/10/2020	10 U 598901 5586986	1668 m	370	14.83	Soil
M20074	31/10/2020	10 U 598778 5587004	1664 m	765	12.62	Soil
M20075	31/10/2020	10 U 598779 5586975	1663 m	765	4.6187	Soil
MR20010	29/10/2020	10 U 597872 5585937	1047 m	10.68%	11.28	Float rock. Felsic quartz diorite with epidote and malachite.
MR20010	29/10/2020	10 U 597872 5585937	1047 m	479	29.95	
MR20011	30/10/2020	10 U 598227 5586048	1106 m	49	27.42	Float rock. Felsic quartz diorite with epidote, gniessic.
MR20012	30/10/2020	10 U 599085 5587251	1613 m	49	12.74	Subcrop rock, quartz fp gniess
MR20013	30/10/2020	10 U 598910 5587203	1650 m	91	13.92	Black, biotie rich hornfels Outcrop rock .
MR20014	30/10/2020	10 U 598929 5587049	1664 m	130	9.98	Outcrop rock with quartz vein. Some pyrite, black biotite hornfels.
MR20015	30/10/2020	10 U 599003 5587024	1659 m	231	11.98	Outcrop rock. Green altered andesite
MR20016	31/10/2020	10 U 599032 5587039	1657 m	88	15.43	Subcrop rock. Some disseminated pyrite, Green altered andesite.
MR20017	31/10/2020	10 U 599017 5587045	1663 m	264	10.27	Subcrop rock. Dark greenish Diorite
MR20018	31/10/2020	10 U 598993 5587054	1668 m	49	25.38	Subcrop rock. Light grey fine-grained felsite
MR20019	31/10/2020	10 U 598922 5586993	1667 m	247	11.42	Subcrop rock.
MR20020	31/10/2020	10 U 598779 5586971	1663 m	4.41%	11.72	Float rock. Some quartz with bright green mineralization. Very rusty
MR20021	31/10/2020	10 U 598790 5586973	1668 m	798	22.89	Float rock. Likely blasted from BOBADIT. 40cm wide mineralized quartz vein with chalcopyrite and malachite exposed by BOBADIT. Qtz-fp gneiss, malachite
MR20022	31/10/2020	10 U 598812 5586955	1673 m	7.68%	11.91	Outcrop rock. Vuggy, Quartz vein with malachite. Lots of malachite
MR20023	31/10/2020	10 U 598825 5586953	1674 m	8.09%	9.16	Outcrop rock. Vuggy, Quartz vein with abundant malachite.
MR20023	31/10/2020	10 U 598825 5586953	1674 m	7.62%	8.69	

APPENDIX IV

ASSAY RESULTS

November 17, 2020



ALS Canada Ltd.
 2103 Dollarton Hwy
 North Vancouver BC V7H 0A7
 Phone: +1 604 984 0221 Fax: +1 604 984 0218
 www.alsglobal.com/geochemistry

To: **HOMEGOLD RESOURCES LTD.**
UNIT 5, 2330 TYNER ST.
PORT COQUITLAM BC V3C 2Z1

Page: 1
 Total # Pages: 2 (A - D)
 Plus Appendix Pages
 Finalized Date: 9-NOV-2020
 Account: MWE

CERTIFICATE VA20246194

Project: McGill

This report is for 16 Soil samples submitted to our lab in Vancouver, BC, Canada on 26-OCT-2020.

The following have access to data associated with this certificate:


SEAN BUTLER	JO SHEARER
-------------	------------

SAMPLE PREPARATION	
ALS CODE	DESCRIPTION
WEI-21	Received Sample Weight
LOG-22	Sample login - Rcd w/o BarCode
DISP-01	Disposal of all sample fractions
SCR-41	Screen to -180um and save both

ANALYTICAL PROCEDURES		
ALS CODE	DESCRIPTION	INSTRUMENT
Au-ICP21	Au 30g FA ICP-AES Finish	ICP-AES
ME-MS41	Ultra Trace Aqua Regia ICP-MS	

This is the Final Report and supersedes any preliminary report with this certificate number. Results apply to samples as submitted. All pages of this report have been checked and approved for release.

***** See Appendix Page for comments regarding this certificate *****

Signature: 
 Saa Traxler, General Manager, North Vancouver



ALS Canada Ltd.
 2103 Dollarton Hwy
 North Vancouver BC V7H 0A7
 Phone: +1 604 984 0221 Fax: +1 604 984 0218
 www.alsglobal.com/geochemistry

To: HOMEGOLD RESOURCES LTD.
 UNIT 5, 2330 TYNER ST.
 PORT COQUITLAM BC V3C 2Z1

Page: 2 - A
 Total # Pages: 2 (A - D)
 Plus Appendix Pages
 Finalized Date: 9-NOV-2020
 Account: MWE

Project: McGill

CERTIFICATE OF ANALYSIS VA20246194

Sample Description	Method Analyte Units LOD	WEI-21 Recvd Wt. kg	Au-ICP21 Au ppm	ME-MS41 Ag ppm	ME-MS41 Al %	ME-MS41 As ppm	ME-MS41 Au ppm	ME-MS41 B ppm	ME-MS41 Ba ppm	ME-MS41 Be ppm	ME-MS41 Bi ppm	ME-MS41 Ca %	ME-MS41 Cd ppm	ME-MS41 Ce ppm	ME-MS41 Co ppm	ME-MS41 Cr ppm
M-20-001		0.32	0.003	0.12	1.92	6.0	<0.02	<10	110	0.36	0.19	0.57	0.22	15.25	11.7	23
M-20-002		0.24	<0.001	0.07	2.43	3.4	<0.02	<10	210	0.39	0.14	0.56	0.21	13.80	9.4	29
M-20-003		0.22	0.001	0.16	2.35	4.8	<0.02	<10	130	0.41	0.12	0.60	0.18	15.70	9.8	29
M-20-004		0.22	0.001	0.16	2.51	7.9	<0.02	<10	90	0.51	0.14	0.81	0.17	21.5	13.0	32
M-20-005		0.24	0.003	0.26	3.69	12.0	<0.02	<10	90	0.71	0.19	1.12	0.17	23.3	16.8	41
M-20-006		0.28	0.002	0.16	3.97	11.2	<0.02	<10	90	0.57	0.30	1.07	0.38	19.50	34.8	36
M-20-007		0.34	0.021	0.30	3.38	8.2	0.02	<10	60	0.55	0.41	1.00	0.63	13.80	33.1	23
M-20-008		0.24	0.002	0.05	2.45	2.7	<0.02	<10	140	0.43	0.16	0.77	0.07	13.25	14.9	19
M-20-009		0.30	0.006	0.07	3.64	3.4	<0.02	<10	250	0.48	0.10	1.37	0.07	12.70	16.5	17
M-20-010		0.24	<0.001	0.04	2.87	3.4	<0.02	<10	200	0.48	0.14	0.68	0.08	16.60	15.3	30
M-20-011		0.24	<0.001	0.04	2.31	1.9	<0.02	<10	220	0.42	0.11	0.53	0.07	13.00	10.2	24
M-20-012		0.26	0.002	0.06	1.92	2.2	<0.02	<10	190	0.37	0.09	0.45	0.07	14.20	10.6	19
M-20-013		0.22	<0.001	0.02	2.45	2.7	<0.02	<10	280	0.47	0.16	0.48	0.06	17.80	12.0	30
M-20-014		0.20	<0.001	0.03	1.39	2.2	<0.02	10	140	0.16	0.09	0.39	0.08	5.67	5.2	17
M-20-015		0.24	0.001	0.03	2.20	3.1	<0.02	<10	230	0.37	0.44	0.70	0.08	15.15	11.9	34
M-20-016		0.24	0.001	0.34	2.80	11.4	<0.02	10	80	0.37	0.15	0.82	1.19	10.30	31.5	40

***** See Appendix Page for comments regarding this certificate *****



ALS Canada Ltd.
 2103 Dollarton Hwy
 North Vancouver BC V7H 0A7
 Phone: +1 604 984 0221 Fax: +1 604 984 0218
 www.alsglobal.com/geochemistry

To: **HOMEGOLD RESOURCES LTD.**
UNIT 5, 2330 TYNER ST.
PORT COQUITLAM BC V3C 2Z1

Page: 2 - B
 Total # Pages: 2 (A - D)
 Plus Appendix Pages
 Finalized Date: 9-NOV-2020
 Account: MWE

Project: McGill

CERTIFICATE OF ANALYSIS VA20246194

Sample Description	Method Analyte Units LOD	ME-MS41	ME-MS41	ME-MS41	ME-MS41	ME-MS41	ME-MS41	ME-MS41	ME-MS41	ME-MS41	ME-MS41	ME-MS41	ME-MS41	ME-MS41	ME-MS41	
		Cs ppm 0.05	Cu ppm 0.2	Fe % 0.01	Ga ppm 0.05	Ge ppm 0.05	Hf ppm 0.02	Hg ppm 0.01	In ppm 0.005	K % 0.01	La ppm 0.2	Li ppm 0.1	Mg % 0.01	Mn ppm 5	Mo ppm 0.05	Na % 0.01
M-20-001		0.78	38.3	3.04	5.95	<0.05	0.08	0.03	0.026	0.14	6.6	8.5	0.75	460	1.63	0.02
M-20-002		0.98	24.4	2.62	6.56	<0.05	0.14	0.02	0.025	0.16	6.3	12.0	0.63	594	0.58	0.02
M-20-003		1.01	26.5	2.99	6.64	0.06	0.29	0.02	0.029	0.27	7.9	10.4	0.66	490	0.74	0.02
M-20-004		0.96	30.5	3.73	7.51	0.08	0.29	0.02	0.032	0.30	9.9	9.9	0.84	481	1.09	0.03
M-20-005		0.96	55.4	4.64	9.88	0.10	0.30	0.02	0.038	0.29	13.9	12.2	1.26	627	1.15	0.03
M-20-006		2.11	76.7	5.37	9.67	0.09	0.17	0.11	0.042	0.28	10.2	14.8	1.09	933	1.99	0.02
M-20-007		1.15	131.0	6.05	8.21	0.14	0.15	0.32	0.043	0.18	7.7	17.7	1.39	1020	2.61	0.02
M-20-008		1.49	44.5	3.60	9.40	0.08	0.14	0.07	0.025	0.14	6.2	14.9	0.95	592	0.60	0.02
M-20-009		2.82	38.5	4.44	12.55	0.12	0.13	0.05	0.028	0.30	6.7	20.1	1.24	764	0.42	0.02
M-20-010		1.58	45.1	3.62	8.58	0.09	0.27	0.02	0.026	0.32	8.2	14.2	0.89	431	0.58	0.02
M-20-011		1.63	24.3	2.82	6.88	0.06	0.19	0.02	0.021	0.25	6.1	12.0	0.64	485	0.37	0.02
M-20-012		2.00	25.7	2.60	6.41	0.05	0.15	0.02	0.019	0.31	6.1	11.1	0.66	459	0.37	0.02
M-20-013		1.14	24.2	2.86	6.80	0.05	0.25	0.02	0.026	0.14	7.8	13.4	0.65	581	0.55	0.02
M-20-014		1.08	8.9	1.58	4.72	<0.05	0.06	0.02	0.011	0.21	2.7	9.8	0.37	407	0.30	0.02
M-20-015		0.68	23.0	2.49	6.78	<0.05	0.11	0.08	0.019	0.19	6.5	11.4	0.78	527	0.34	0.02
M-20-016		0.85	171.0	5.12	7.96	0.08	0.10	0.04	0.027	0.22	5.0	12.5	1.11	782	1.37	0.02

***** See Appendix Page for comments regarding this certificate *****



ALS Canada Ltd.
 2103 Dollarton Hwy
 North Vancouver BC V7H 0A7
 Phone: +1 604 984 0221 Fax: +1 604 984 0218
 www.alsglobal.com/geochemistry

To: HOMEGOLD RESOURCES LTD.
 UNIT 5, 2330 TYNER ST.
 PORT COQUITLAM BC V3C 2Z1

Page: 2 - C
 Total # Pages: 2 (A - D)
 Plus Appendix Pages
 Finalized Date: 9-NOV-2020
 Account: MWE

Project: McGill

CERTIFICATE OF ANALYSIS VA20246194

Sample Description	Method Analyte Units LOD	ME-MS41	ME-MS41	ME-MS41	ME-MS41	ME-MS41	ME-MS41	ME-MS41	ME-MS41	ME-MS41	ME-MS41	ME-MS41	ME-MS41	ME-MS41	ME-MS41	
		Nb ppm 0.05	Ni ppm 0.2	P ppm 10	Pb ppm 0.2	Rb ppm 0.1	Re ppm 0.001	S % 0.01	Sb ppm 0.05	Sc ppm 0.1	Se ppm 0.2	Sn ppm 0.2	Sr ppm 0.2	Ta ppm 0.01	Te ppm 0.01	Th ppm 0.2
M-20-001		0.81	20.3	690	4.1	6.8	<0.001	0.02	0.20	6.4	0.2	0.4	61.5	<0.01	0.18	0.9
M-20-002		0.84	23.5	750	4.4	10.0	<0.001	0.01	0.13	5.2	<0.2	0.5	49.7	<0.01	0.06	1.3
M-20-003		1.00	24.3	520	4.5	10.5	<0.001	0.01	0.19	7.0	0.4	0.5	51.9	<0.01	0.05	1.2
M-20-004		1.40	26.9	470	5.7	10.0	<0.001	0.02	0.21	9.3	0.3	0.7	65.3	<0.01	0.10	1.2
M-20-005		1.32	40.1	830	7.0	8.5	<0.001	0.03	0.69	12.6	0.4	0.6	92.4	<0.01	0.16	1.2
M-20-006		0.88	35.1	1080	5.8	9.7	<0.001	0.05	0.31	16.5	0.6	0.6	98.9	0.01	0.35	1.2
M-20-007		0.57	27.6	970	4.0	7.5	<0.001	0.04	0.30	20.7	1.0	0.5	108.0	<0.01	1.98	0.7
M-20-008		0.51	16.5	350	4.0	6.3	<0.001	0.01	0.21	8.8	0.3	0.6	59.8	<0.01	0.10	1.4
M-20-009		0.51	15.5	770	3.3	12.3	<0.001	0.01	0.14	9.1	<0.2	0.6	90.5	<0.01	0.09	1.4
M-20-010		0.63	23.8	310	4.3	11.7	<0.001	0.01	0.20	9.3	0.3	0.5	55.4	<0.01	0.06	1.7
M-20-011		0.63	17.1	280	3.9	12.6	<0.001	0.01	0.14	6.6	0.2	0.4	42.5	<0.01	0.03	1.3
M-20-012		0.67	13.6	300	3.5	12.8	<0.001	0.01	0.16	5.5	0.2	0.4	35.5	<0.01	0.07	1.3
M-20-013		0.85	21.6	340	5.4	11.4	<0.001	0.01	0.16	6.3	0.2	0.5	35.6	<0.01	0.04	1.7
M-20-014		0.80	11.5	580	3.7	11.0	<0.001	0.01	0.08	2.5	<0.2	0.3	26.2	<0.01	0.01	0.6
M-20-015		0.46	35.2	630	5.3	6.0	<0.001	0.02	0.10	5.5	0.2	0.4	47.6	<0.01	0.24	1.4
M-20-016		0.59	25.2	910	13.5	4.5	<0.001	0.05	0.22	13.2	0.7	0.3	63.9	<0.01	0.19	0.8



ALS Canada Ltd.
 2103 Dollarton Hwy
 North Vancouver BC V7H 0A7
 Phone: +1 604 984 0221 Fax: +1 604 984 0218
 www.alsglobal.com/geochemistry

To: HOMEGOLD RESOURCES LTD.
 UNIT 5, 2330 TYNER ST.
 PORT COQUITLAM BC V3C 2Z1

Page: 2 - D
 Total # Pages: 2 (A - D)
 Plus Appendix Pages
 Finalized Date: 9-NOV-2020
 Account: MWE

Project: McGill

CERTIFICATE OF ANALYSIS VA20246194

Sample Description	Method Analyte Units LOD	ME-MS41	ME-MS41	ME-MS41	ME-MS41	ME-MS41	ME-MS41	ME-MS41	ME-MS41
		Ti %	Ti ppm	U ppm	V ppm	W ppm	Y ppm	Zn ppm	Zr ppm
M-20-001		0.107	0.05	0.42	67	0.09	6.06	80	3.4
M-20-002		0.113	0.07	0.37	53	0.07	5.13	151	6.3
M-20-003		0.146	0.08	0.41	62	0.09	7.97	123	13.3
M-20-004		0.219	0.09	0.61	87	0.10	10.45	84	15.9
M-20-005		0.231	0.08	0.71	107	0.12	18.80	99	16.1
M-20-006		0.206	0.11	1.36	113	0.12	23.3	146	8.6
M-20-007		0.272	0.06	1.15	155	0.32	27.7	231	6.5
M-20-008		0.182	0.05	0.39	83	0.11	10.70	76	4.5
M-20-009		0.234	0.11	0.54	102	0.10	12.95	84	4.9
M-20-010		0.191	0.08	0.46	83	0.10	11.15	71	10.6
M-20-011		0.156	0.08	0.34	62	0.08	6.61	73	7.6
M-20-012		0.114	0.07	0.32	56	0.13	5.21	71	5.2
M-20-013		0.145	0.09	0.46	64	0.10	6.19	89	10.6
M-20-014		0.093	0.05	0.12	35	0.08	1.62	99	2.0
M-20-015		0.050	0.05	0.33	49	0.06	5.51	80	2.8
M-20-016		0.138	0.05	0.44	135	0.10	9.70	263	4.2

***** See Appendix Page for comments regarding this certificate *****



ALS Canada Ltd.
2103 Dollarton Hwy
North Vancouver BC V7H 0A7
Phone: +1 604 984 0221 Fax: +1 604 984 0218
www.alsglobal.com/geochemistry

To: **HOMEGOLD RESOURCES LTD.**
UNIT 5, 2330 TYNER ST.
PORT COQUITLAM BC V3C 2Z1

Page: 1
Total # Pages: 2 (A - D)
Plus Appendix Pages
Finalized Date: 12-NOV-2020
Account: MWE

CERTIFICATE VA20246211

Project: McGill

This report is for 4 Rock samples submitted to our lab in Vancouver, BC, Canada on 26-OCT-2020.

The following have access to data associated with this certificate:

SEAN BUTLER

JO SHEARER

SAMPLE PREPARATION

ALS CODE	DESCRIPTION
WEI-21	Received Sample Weight
LOG-22	Sample login - Rcd w/o BarCode
DISP-01	Disposal of all sample fractions
CRU-31	Fine crushing - 70% <2mm
SPL-21	Split sample - riffle splitter
PUL-31	Pulverize up to 250g 85% <75 um

ANALYTICAL PROCEDURES

ALS CODE	DESCRIPTION
ME-MS61	48 element four acid ICP-MS
Au-ICP21	Au 30g FA ICP-AES Finish ICP-AES

This is the Final Report and supersedes any preliminary report with this certificate number. Results apply to samples as submitted. All pages of this report have been checked and approved for release.

***** See Appendix Page for comments regarding this certificate *****

Signature:

Saa Traxler, General Manager, North Vancouver



ALS Canada Ltd.
 2103 Dollarton Hwy
 North Vancouver BC V7H 0A7
 Phone: +1 604 984 0221 Fax: +1 604 984 0218
 www.alsglobal.com/geochemistry

To: HOMEGOLD RESOURCES LTD.
 UNIT 5, 2330 TYNER ST.
 PORT COQUITLAM BC V3C 2Z1

Page: 2 - A
 Total # Pages: 2 (A - D)
 Plus Appendix Pages
 Finalized Date: 12-NOV-2020
 Account: MWE

Project: McGill

CERTIFICATE OF ANALYSIS VA20246211

Sample Description	Method Analyte Units LOD	WEI-21 Recvd Wt. kg	Au-ICP21 Au ppm	ME-MS61 Ag ppm	ME-MS61 Al %	ME-MS61 As ppm	ME-MS61 Ba ppm	ME-MS61 Be ppm	ME-MS61 Bi ppm	ME-MS61 Ca %	ME-MS61 Cd ppm	ME-MS61 Ce ppm	ME-MS61 Co ppm	ME-MS61 Cr ppm	ME-MS61 Cs ppm	ME-MS61 Cu ppm
MR20-001		1.76	0.001	0.17	8.01	1.4	120	0.37	0.15	3.79	0.04	8.90	23.5	3	1.01	33.2
MR20-002		0.54	<0.001	0.08	8.39	0.3	190	0.31	0.06	6.60	0.30	6.31	43.5	88	0.38	39.7
MR20-003		0.50	<0.001	0.07	7.69	1.9	220	0.63	0.52	0.41	0.56	13.15	5.9	5	0.09	24.0
MR20-004		0.60	<0.001	0.09	8.63	1.0	80	0.31	0.07	4.76	0.38	9.07	23.6	46	0.30	62.3

4 Rocks



ALS Canada Ltd.
 2103 Dollarton Hwy
 North Vancouver BC V7H 0A7
 Phone: +1 604 984 0221 Fax: +1 604 984 0218
 www.alsglobal.com/geochemistry

To: **HOMEGOLD RESOURCES LTD.**
UNIT 5, 2330 TYNER ST.
PORT COQUITLAM BC V3C 2Z1

Page: 2 - B
 Total # Pages: 2 (A - D)
 Plus Appendix Pages
 Finalized Date: 12-NOV-2020
 Account: MWE

Project: McGill

CERTIFICATE OF ANALYSIS VA20246211

Sample Description	Method Analyte Units LOD	ME-MS61	ME-MS61	ME-MS61	ME-MS61	ME-MS61	ME-MS61	ME-MS61	ME-MS61	ME-MS61	ME-MS61	ME-MS61	ME-MS61	ME-MS61	ME-MS61	
		Fe %	Ca ppm	Ge ppm	Hf ppm	In ppm	K %	La ppm	Li ppm	Mg %	Mn ppm	Mo ppm	Na %	Nb ppm	Ni ppm	P ppm
		0.01	0.05	0.05	0.1	0.005	0.01	0.5	0.2	0.01	5	0.05	0.01	0.1	0.2	
MR20-001		8.83	18.90	0.09	1.0	0.075	0.84	2.7	12.8	2.98	1900	0.59	2.46	0.7	4.8	1300
MR20-002		7.93	17.90	0.08	0.8	0.061	0.45	2.2	6.0	3.84	2040	0.16	2.52	0.4	32.0	450
MR20-003		2.28	13.80	0.11	1.6	0.007	0.23	5.6	1.6	0.06	163	4.36	5.82	0.6	2.7	170
MR20-004		7.96	20.1	0.09	0.6	0.062	0.20	3.2	4.8	3.58	2780	0.44	3.49	0.8	17.9	620

***** See Appendix Page for comments regarding this certificate *****



ALS Canada Ltd.
 2103 Dollarton Hwy
 North Vancouver BC V7H 0A7
 Phone: +1 604 984 0221 Fax: +1 604 984 0218
 www.alsglobal.com/geochemistry

To: **HOMEGOLD RESOURCES LTD.**
UNIT 5, 2330 TYNER ST.
PORT COQUITLAM BC V3C 2Z1

Page: 2 - C
 Total # Pages: 2 (A - D)
 Plus Appendix Pages
 Finalized Date: 12-NOV-2020
 Account: MWE

Project: McGill

CERTIFICATE OF ANALYSIS VA20246211

Sample Description	Method Analyte Units LOD	ME-MS61	ME-MS61	ME-MS61	ME-MS61	ME-MS61	ME-MS61	ME-MS61	ME-MS61	ME-MS61	ME-MS61	ME-MS61	ME-MS61	ME-MS61	ME-MS61	
		Pb	Rb	Re	S	Sb	Sc	Se	Sn	Sr	Ta	Te	Th	Ti	Tl	U
		ppm	ppm	ppm	%	ppm	ppm	ppm	ppm	ppm	ppm	ppm	ppm	%	ppm	ppm
		0.5	0.1	0.002	0.01	0.05	1	0.2	0.2	0.05	0.05	0.01	0.005	0.02	0.1	
MR20-001		3.6	12.6	0.101	3.61	0.40	2	0.6	226	<0.05	0.55	0.31	0.851	0.14	1.0	
MR20-002		11.6	3.3	0.002	0.03	0.17	<1	0.4	188.0	<0.05	<0.05	0.16	0.540	0.08	0.6	
MR20-003		13.6	2.8	<0.002	0.04	0.17	1	<0.2	91.2	0.10	0.42	3.48	0.067	0.02	0.6	
MR20-004		10.8	1.9	<0.002	0.26	0.18	<1	0.7	128.5	0.05	0.12	0.38	0.699	0.03	0.4	



ALS Canada Ltd.
 2103 Dollarton Hwy
 North Vancouver BC V7H 0A7
 Phone: +1 604 984 0221 Fax: +1 604 984 0218
 www.alsglobal.com/geochemistry

To: **HOMEGOLD RESOURCES LTD.**
UNIT 5, 2330 TYNER ST.
PORT COQUITLAM BC V3C 2Z1

Page: 2 - D
 Total # Pages: 2 (A - D)
 Plus Appendix Pages
 Finalized Date: 12-NOV-2020
 Account: MWE

Project: McGill

CERTIFICATE OF ANALYSIS VA20246211

Sample Description	Method Analyte Units LOD	ME-MS61	ME-MS61	ME-MS61	ME-MS61	ME-MS61
		V	W	Y	Zn	Zr
		ppm	ppm	ppm	ppm	ppm
		1	0.1	0.1	2	0.5
MR20-001		273	0.3	24.6	135	31.1
MR20-002		371	0.1	16.5	179	19.4
MR20-003		23	0.4	3.6	131	32.7
MR20-004		366	0.2	19.3	168	11.3

***** See Appendix Page for comments regarding this certificate *****



ALS Canada Ltd.
2103 Dollarton Hwy
North Vancouver BC V7H 0A7
Phone: +1 604 984 0221 Fax: +1 604 984 0218
www.alsglobal.com/geochemistry

To: **HOMEGOLD RESOURCES LTD.**
UNIT 5, 2330 TYNER ST.
PORT COQUITLAM BC V3C 2Z1

Page: **Appendix 1**
Total # Appendix Pages: **1**
Finalized Date: **12-NOV-2020**
Account: **MWE**

Project: McGill

CERTIFICATE OF ANALYSIS VA20246211

CERTIFICATE COMMENTS

ANALYTICAL COMMENTS

Applies to Method: REEs may not be totally soluble in this method.
ME-MS61

LABORATORY ADDRESSES

Applies to Method: Processed at ALS Vancouver located at 2103 Dollarton Hwy, North Vancouver, BC, Canada.
Au-ICP21 CRU-31 DISP-01 LOG-22
ME-MS61 PUL-31 SPL-21 WEI-21

McGillivray 2020 XRF Assays

All Results in %

Sample #	Mg	Mg +/-	Al	Al +/-	Si	Si +/-	P	P +/-	S	S +/-	Cl	Cl +/-	K	K +/-	Ca	Ca +/-	Ti
MR20010	1.64	0.49	2.94	0.08	11.28	0.14	1.9479	0.0412	0.6555	0.0098	ND		0.5622	0.0082	1.3739	0.0175	ND
MR20010	ND		1.24	0.0466	29.95	0.17	0.9607	0.0261	0.1147	0.0036	ND		0.8327	0.0066	0.194	0.0049	0.0451
MR20011	ND		1.743	0.0497	27.42	0.16	0.9688	0.0273	0.1295	0.0038	ND		0.1088	0.0036	1.7118	0.0118	0.064
MR20012	ND		2.98	0.07	12.74	0.1	1.7848	0.0321	1.396	0.0124	ND		1.2805	0.0109	1.0254	0.0095	ND
MR20013	1.74	0.32	4.35	0.07	13.92	0.12	1.0204	0.0281	0.12	0.0034	ND		0.2421	0.0039	8.6	0.07	0.3858
MR20014	1	0.32	4.13	0.08	9.98	0.09	0.89	0.0227	0.1134	0.0031	ND		2.8207	0.0252	2.2739	0.0209	0.8016
MR20015	ND		3.46	0.07	11.98	0.11	1.0199	0.0279	0.1006	0.0033	ND		0.0665	0.003	8.09	0.07	0.6339
MR20016	ND		5.4	0.07	15.43	0.11	1.4172	0.0275	0.1914	0.0036	ND		1.3782	0.0106	4.4789	0.0316	0.6067
MR20017	ND		3.46	0.07	10.27	0.09	0.9367	0.0278	0.1034	0.0032	ND		0.0866	0.003	12.22	0.1	0.5068
MR20018	ND		3.35	0.06	25.38	0.16	1.2699	0.0294	0.2502	0.0045	ND		1.3132	0.0098	0.8338	0.008	0.2597
MR20019	ND		5.49	0.09	11.42	0.1	1.2661	0.0358	0.1456	0.0042	ND		ND		17.98	0.15	0.0695
MR20020	ND		2.71	0.07	11.72	0.12	1.5536	0.0343	0.7251	0.0091	ND		0.161	0.0041	1.1279	0.0122	ND
MR20021	ND		1.3672	0.0484	22.89	0.15	1.3008	0.0299	0.1677	0.0039	ND		0.028	0.0032	3.6333	0.024	0.0508
MR20022	ND		1.5243	0.0413	11.91	0.1	0.7639	0.0174	1.0673	0.0094	ND		0.0077	0.0019	0.2735	0.0034	ND
MR20023	ND		1.8498	0.0445	9.16	0.08	0.8163	0.0176	0.3366	0.0041	ND		0.0443	0.0019	0.322	0.0038	ND
MR20023	ND		3.51	0.09	8.69	0.1	1.7307	0.04	0.3529	0.0063	ND		0.261	0.0051	8.3	0.09	0.1333
M20030	ND		4.88	0.06	14.56	0.1	1.8783	0.0279	0.1043	0.0028	ND		0.6387	0.0054	2.032	0.0139	0.3086
M20031	ND		4.52	0.07	13.48	0.1	1.0312	0.0246	0.1297	0.0033	ND		0.6143	0.0059	2.103	0.0161	0.2795
M20032	ND		3.98	0.07	11.45	0.09	0.8746	0.0221	0.1051	0.003	ND		0.5608	0.0053	1.6997	0.0132	0.2402
M20033	ND		4.71	0.06	16.06	0.11	1.1314	0.0236	0.0949	0.0028	ND		0.7607	0.0062	2.3441	0.016	0.2776
M20034	0.89	0.24	4.63	0.07	14.44	0.1	0.7656	0.0204	0.0873	0.0027	ND		0.6641	0.0058	2.1797	0.016	0.3009
M20035	ND		3.95	0.06	13.34	0.09	0.9214	0.0219	0.0987	0.0029	ND		0.5496	0.0051	1.3777	0.0104	0.2197
M20036	ND		4.57	0.06	12.89	0.09	1.9159	0.0281	0.0871	0.0027	ND		0.6487	0.0055	2.1557	0.0152	0.3198
M20037	ND		4.92	0.07	16.81	0.11	0.9746	0.0226	0.0864	0.0028	ND		0.8478	0.0066	2.6333	0.0176	0.3275
M20038	ND		4.65	0.06	14.03	0.09	1.1873	0.0227	0.0816	0.0026	ND		0.683	0.0056	1.9206	0.0133	0.2993
M20039	ND		3.75	0.07	15.66	0.11	0.9978	0.0252	0.1142	0.0034	ND		0.6397	0.006	1.8239	0.0139	0.2419
M20040	ND		4.83	0.07	16.69	0.11	0.9753	0.0227	0.0852	0.0028	ND		0.7071	0.0059	2.3448	0.0158	0.2927
M20044	ND		4.89	0.07	18.84	0.12	0.8288	0.0225	0.1009	0.0031	ND		0.6408	0.0056	1.5862	0.0114	0.2508
M20045	ND		5.43	0.07	15.74	0.11	1.0432	0.0251	0.0993	0.0031	ND		0.7005	0.0063	3.1609	0.0229	0.3601
M20046	ND		5.82	0.08	18.34	0.12	1.0321	0.0256	0.1026	0.0033	ND		1.2158	0.0092	2.4041	0.0168	0.2355

Date	Mg	Mg +/-	Al	Al +/-	Si	Si +/-	P	P +/-	S	S +/-	Cl	Cl +/-	K	K +/-	Ca	Ca +/-	Ti
M20047	ND		4.03	0.06	14.86	0.1	0.8251	0.0219	0.1109	0.0029	ND		0.6592	0.0057	4.4555	0.0305	0.2977
M20048	ND		4.27	0.06	16.32	0.11	0.8162	0.0223	0.1071	0.003	ND		0.7695	0.0065	3.6017	0.0249	0.4057
M20050	ND		4.47	0.08	15.41	0.13	1.0337	0.0281	0.1335	0.0038	ND		0.8076	0.0079	3.2947	0.0267	0.3556
M20051	0.82	0.24	4.79	0.07	15.84	0.12	0.8393	0.0224	0.0945	0.0029	ND		0.5455	0.0053	3.2495	0.0236	0.277
M20052	ND		5.4	0.07	14.68	0.1	0.8236	0.0219	0.0881	0.0028	ND		0.4982	0.0048	3.615	0.025	0.3542
M20053	ND		4.86	0.07	16.47	0.11	0.7967	0.022	0.0832	0.0029	ND		0.6091	0.0055	2.5631	0.0179	0.3937
M20053	0.98	0.24	4.85	0.07	16.42	0.12	0.7721	0.0217	0.086	0.0029	ND		0.6291	0.0058	2.6482	0.0193	0.3527
M20054	ND		4.35	0.07	14.5	0.1	1.011	0.0232	0.1026	0.0029	ND		0.6004	0.0055	2.7677	0.0199	0.3238
M20054	ND		4.89	0.07	16.28	0.11	0.8889	0.0229	0.1044	0.003	ND		0.8254	0.0069	2.9514	0.0208	0.3673
M20055	ND		4.57	0.07	17.35	0.12	1.0877	0.0255	0.0961	0.0031	ND		0.9046	0.0076	3.1578	0.0228	0.4153
M20056	ND		3.8	0.07	11.55	0.1	0.819	0.0233	0.1632	0.0035	ND		0.6084	0.0062	3.2752	0.0268	0.3866
M20057	ND		4.13	0.07	13.31	0.11	0.8816	0.0248	0.1172	0.0034	ND		0.5695	0.0059	3.13	0.0253	0.3223
M20058	ND		5.11	0.07	17.04	0.12	0.8935	0.0231	0.1018	0.0031	ND		0.7248	0.0063	1.5591	0.0118	0.3567
M20059	ND		3.37	0.05	21.46	0.13	0.8438	0.0215	0.0983	0.0028	ND		0.6943	0.0056	1.5747	0.0107	0.2592
M20060	ND		4.03	0.06	12.39	0.09	0.9208	0.0205	0.09	0.0025	ND		0.5447	0.0049	2.382	0.017	0.3485
M20061	ND		3.68	0.06	9.96	0.08	0.8018	0.0199	0.0902	0.0026	ND		0.3439	0.0039	1.6599	0.0133	0.4038
M20062	ND		4.88	0.07	10.89	0.08	0.827	0.0195	0.1014	0.0025	ND		0.509	0.0047	2.4798	0.0182	0.403
M20063	ND		4.19	0.06	15.09	0.1	1.0246	0.0212	0.0898	0.0026	ND		0.7112	0.0056	1.923	0.013	0.2954
M20064	ND		4.59	0.07	13.25	0.1	1.055	0.0238	0.1127	0.0031	ND		0.5438	0.0053	2.0694	0.0159	0.3508
M20065	ND		4.79	0.06	14.2	0.09	0.9789	0.0207	0.0946	0.0025	ND		0.6062	0.005	2.3764	0.0157	0.345
M20066	1.2	0.26	4.89	0.07	12.25	0.1	0.9447	0.0221	0.0953	0.0028	ND		0.5524	0.0054	2.4592	0.0191	0.3813
M20067	ND		3.94	0.06	14.38	0.1	0.9092	0.0216	0.1199	0.0029	ND		0.6137	0.0054	2.3253	0.0164	0.2765
M20068	ND		5.12	0.07	14.3	0.1	0.9718	0.0213	0.1161	0.0028	ND		0.6782	0.0057	1.6515	0.0119	0.374
M20069	ND		5.12	0.07	15.21	0.11	0.9009	0.0221	0.0979	0.0029	ND		0.5319	0.005	2.1174	0.0151	0.3215
M20070	ND		4.83	0.07	15.5	0.11	0.9057	0.023	0.0913	0.003	ND		0.5805	0.0055	2.3058	0.0169	0.3909
M20071	ND		4.31	0.06	13.68	0.1	0.9203	0.0214	0.1108	0.0028	ND		0.5868	0.0053	2.1096	0.0153	0.3266
M20072	ND		4.64	0.07	12.47	0.1	0.9373	0.0242	0.1109	0.0033	ND		0.5046	0.0053	2.243	0.0178	0.2959
M20073	1.11	0.3	5.31	0.08	14.83	0.13	0.8716	0.0261	0.1051	0.0035	ND		0.6992	0.0072	3.676	0.0306	0.3522
M20074	0.95	0.28	5.16	0.08	12.62	0.11	0.9115	0.0241	0.0921	0.0031	ND		0.5682	0.0059	3.3753	0.0276	0.368
M20075	2.83	0.58	2.43	0.08	4.619	0.04	0.9631	0.0265	0.1366	0.0039	ND		0.9963	0.0081	0.8737	0.0076	ND

Ti +/-	V	V +/-	Cr	Cr +/-	Mn	Mn +/-	Fe	Fe +/-	Co	Co +/-	Ni	Ni +/-	Cu	Cu +/-	Zn	Zn +/-	As
	ND		ND		0.0214	0.0045	18.47	0.22	ND		ND		10.68	0.13	0.0749	0.0067	0.0034
0.0143	0.0392	0.008	ND		0.0114	0.0027	0.3576	0.0076	ND		ND		0.0479	0.0015	0.0021	0.0004	ND
0.0154	ND		ND		0.0129	0.0028	0.1568	0.0053	ND		ND		0.0049	0.0007	ND		ND
	ND		ND		0.0774	0.0047	2.486	0.0265	ND		ND		0.0049	0.0008	0.0031	0.0005	ND
0.0214	0.0342	0.0083	0.0177	0.0041	0.2056	0.0074	9.32	0.08	ND		0.0088	0.0014	0.0091	0.0012	0.0161	0.001	ND
0.0248	0.0643	0.0086	ND		0.3856	0.0093	11.65	0.11	ND		ND		0.013	0.0013	0.0636	0.0019	ND
0.0248	ND		ND		0.312	0.009	12.34	0.11	ND		ND		0.0231	0.0017	0.0359	0.0016	ND
0.0231	ND		ND		0.193	0.0065	6.0989	0.0495	ND		ND		0.0088	0.001	0.0159	0.0009	ND
0.0245	ND		ND		0.1739	0.007	6.15	0.06	ND		0.0043	0.0012	0.0264	0.0016	0.0083	0.0008	ND
0.0202	0.0677	0.0096	ND		0.0197	0.0032	0.7188	0.0117	ND		ND		0.0049	0.0008	0.0029	0.0005	ND
0.0196	ND		ND		0.0983	0.0067	8.08	0.08	ND		ND		0.0247	0.0018	0.0173	0.0012	ND
	ND		ND		0.2301	0.0072	0.689	0.0123	ND		0.0037	0.0012	4.409	0.0441	0.0399	0.0027	ND
0.0151	ND		ND		0.0727	0.0046	0.3352	0.0079	ND		ND		0.0798	0.002	0.0024	0.0005	ND
	ND		ND		0.4986	0.0078	0.3522	0.0063	ND		ND		7.68	0.06	ND		ND
	ND		ND		0.8557	0.0114	2.8088	0.0272	ND		ND		8.09	0.07	0.0168	0.0028	ND
0.0181	ND		ND		0.7812	0.0165	6.97	0.08	ND		ND		7.62	0.09	0.1772	0.0059	0.002
0.0169	0.0289	0.0067	ND		0.0803	0.0041	3.0301	0.026	ND		ND		0.0049	0.0007	0.01	0.0006	0.0011
0.0178	ND		ND		0.1164	0.0051	3.4138	0.0316	ND		ND		0.0095	0.0009	0.0172	0.0008	ND
0.0162	0.0202	0.0065	ND		0.0768	0.0041	3.2005	0.0295	ND		0.0034	0.0008	0.0052	0.0008	0.0123	0.0007	0.001
0.0168	0.0325	0.007	ND		0.1092	0.0047	3.2689	0.028	ND		ND		0.004	0.0007	0.0119	0.0007	ND
0.0168	ND		ND		0.0717	0.0039	3.9537	0.0337	ND		ND		0.0085	0.0008	0.0085	0.0006	0.0007
0.0159	0.0324	0.0069	ND		0.0501	0.0035	2.3056	0.0222	ND		ND		0.0041	0.0007	0.006	0.0005	ND
0.0169	0.0392	0.0069	ND		0.0852	0.0041	3.7468	0.0314	ND		0.0025	0.0008	0.0066	0.0008	0.015	0.0007	0.0016
0.0177	ND		ND		0.0674	0.004	3.6136	0.03	ND		ND		0.0065	0.0008	0.0151	0.0007	0.0009
0.0164	0.0288	0.0065	0.0088	0.0029	0.0442	0.0033	3.4402	0.0286	ND		ND		0.0047	0.0007	0.0145	0.0007	ND
0.018	ND		ND		0.0657	0.0042	1.9003	0.0207	ND		ND		0.0029	0.0007	0.009	0.0006	ND
0.0174	ND		0.0099	0.0031	0.0571	0.0037	2.8113	0.025	ND		ND		0.0041	0.0007	0.0083	0.0006	ND
0.0174	ND		ND		0.0357	0.0033	2.4068	0.0226	ND		ND		0.0047	0.0007	0.0049	0.0005	ND
0.0196	0.0267	0.0075	ND		0.0981	0.0049	4.9518	0.0419	ND		ND		0.0131	0.0011	0.0123	0.0008	ND
0.0182	0.0293	0.0078	ND		0.0571	0.004	2.353	0.0232	ND		ND		0.004	0.0007	0.0041	0.0005	0.0008

Ti +/-	V	V +/-	Cr	Cr +/-	Mn	Mn +/-	Fe	Fe +/-	Co	Co +/-	Ni	Ni +/-	Cu	Cu +/-	Zn	Zn +/-	As
0.0173	0.0236	0.0068	ND		0.1994	0.0062	5.4746	0.0437	ND		0.0027	0.0009	0.0142	0.001	0.0122	0.0007	ND
0.0195	0.0225	0.0072	ND		0.2458	0.007	5.0324	0.041	ND		ND		0.0112	0.001	0.0135	0.0008	0.0011
0.0216	0.0259	0.0084	ND		0.161	0.0066	4.7743	0.0455	ND		ND		0.0118	0.0011	0.0151	0.0009	0.0013
0.0174	0.0256	0.007	ND		0.1094	0.0049	4.4664	0.0384	ND		0.003	0.0009	0.0152	0.0011	0.0086	0.0006	ND
0.0183	0.0359	0.0072	ND		0.1145	0.0049	5.219	0.0419	ND		ND		0.0206	0.0012	0.0092	0.0007	ND
0.0194	0.0252	0.0073	ND		0.0742	0.0042	4.0127	0.0338	ND		0.0025	0.0008	0.0103	0.0009	0.006	0.0006	ND
0.0186	0.0246	0.0071	ND		0.1019	0.0047	4.3183	0.0369	ND		ND		0.0097	0.0009	0.009	0.0006	ND
0.0178	ND		ND		0.1435	0.0054	4.6481	0.0388	ND		0.0042	0.0009	0.0106	0.0009	0.0272	0.001	ND
0.0192	0.0272	0.0073	ND		0.1241	0.0052	4.6844	0.0389	ND		ND		0.0091	0.0009	0.0098	0.0007	ND
0.0206	ND		ND		0.146	0.0058	5.1007	0.0428	ND		0.004	0.001	0.0073	0.0009	0.0221	0.001	ND
0.0199	0.0325	0.0076	ND		0.1452	0.0059	6.42	0.06	ND		ND		0.0196	0.0013	0.0124	0.0008	ND
0.0195	0.0245	0.0075	ND		0.1599	0.0062	5.67	0.05	ND		ND		0.0174	0.0013	0.0108	0.0008	ND
0.0192	0.0263	0.0074	ND		0.0758	0.0043	3.6261	0.0317	ND		ND		0.0055	0.0008	0.0115	0.0007	ND
0.0166	0.0252	0.0068	ND		0.2169	0.0062	2.731	0.0235	ND		ND		0.0051	0.0007	0.0159	0.0007	ND
0.0166	0.0273	0.0063	ND		0.307	0.0071	5.628	0.0441	ND		0.0033	0.0008	0.0085	0.0009	0.0219	0.0009	ND
0.0174	ND		ND		0.151	0.0052	6.84	0.06	ND		0.004	0.0009	0.0234	0.0013	0.019	0.0009	0.0012
0.017	0.0275	0.0062	ND		0.141	0.005	7.89	0.06	ND		ND		0.0251	0.0013	0.0199	0.0009	ND
0.016	ND		ND		0.2107	0.0058	3.7624	0.0298	ND		0.003	0.0008	0.0065	0.0007	0.0148	0.0007	0.0007
0.0183	ND		ND		0.1995	0.0063	5.0243	0.0427	ND		0.003	0.0009	0.015	0.0011	0.0183	0.0009	ND
0.0166	0.0306	0.0064	ND		0.1259	0.0046	4.303	0.0331	ND		0.0032	0.0008	0.0196	0.001	0.0166	0.0007	ND
0.018	0.0307	0.0068	ND		0.1192	0.0049	6.21	0.05	ND		0.0028	0.0009	0.0222	0.0012	0.0142	0.0008	ND
0.0164	0.0218	0.0065	ND		0.218	0.0062	4.2232	0.0348	ND		0.0024	0.0008	0.0115	0.0009	0.0207	0.0009	ND
0.0176	0.0361	0.0068	ND		0.1182	0.0047	4.5407	0.0361	ND		0.0044	0.0009	0.0067	0.0008	0.0107	0.0007	0.0007
0.0177	0.0248	0.0069	ND		0.1188	0.0049	4.236	0.035	ND		0.0047	0.0009	0.0072	0.0008	0.0181	0.0008	ND
0.0192	0.0232	0.0071	ND		0.1063	0.0048	4.8042	0.0401	ND		0.0029	0.0009	0.0066	0.0009	0.0155	0.0008	ND
0.017	0.024	0.0065	0.009	0.0029	0.232	0.0064	4.885	0.0395	ND		0.0033	0.0008	0.0157	0.001	0.0234	0.0009	0.0009
0.0184	ND		ND		0.1275	0.0054	4.3141	0.0394	ND		0.0036	0.0009	0.0147	0.0011	0.0134	0.0008	ND
0.0209	0.0311	0.0082	ND		0.1032	0.0054	5.57	0.05	ND		ND		0.037	0.0017	0.0152	0.0009	ND
0.0195	0.0285	0.0074	ND		0.1313	0.0055	6.11	0.06	ND		0.0032	0.001	0.0765	0.0023	0.0176	0.001	ND
	ND		ND		ND		5.7199	0.043	0	0.007	5.0768	0.0371	54.12	0.33	22.46	0.14	ND

As +/-	Se	Se +/-	Rb	Rb +/-	Sr	Sr +/-	Y	Y +/-	Zr	Zr +/-	Mo	Mo +/-	Ag	Ag +/-	Cd	Cd +/-	Sn	Sn +/-
0.0009	ND		0.002	0.0004	0.0229	0.0008	0.0022	0.0004	0.0021	0.0005	0.0105	0.0005	ND		ND		ND	
	ND		0.0013	0.0001	0.0043	0.0002	ND		ND		ND		ND		ND		ND	
	ND		ND		0.0003	0.0001	ND		ND		ND		ND		ND		ND	
	ND		0.0043	0.0002	0.0034	0.0002	0.0006	0.0002	0.0026	0.0002	0.001	0.0002	ND		ND		ND	
	ND		0.0014	0.0002	0.0176	0.0004	0.0021	0.0002	0.0041	0.0003	0.001	0.0002	ND		ND		ND	
	ND		0.0166	0.0004	0.0276	0.0005	0.0015	0.0003	0.0087	0.0004	0.0016	0.0002	ND		ND		ND	
	ND		ND		0.0454	0.0008	0.0013	0.0002	0.0015	0.0003	0.0022	0.0003	ND		ND		ND	
	ND		0.0037	0.0002	0.0223	0.0004	0.004	0.0002	0.0122	0.0003	ND		ND		ND		ND	
	ND		ND		0.0518	0.0007	0.003	0.0002	0.0017	0.0003	ND		ND		ND		ND	
	ND		0.002	0.0001	0.0102	0.0002	0.0007	0.0001	0.0159	0.0003	ND		ND		ND		ND	
	ND		ND		0.0611	0.0009	0.003	0.0003	0.0012	0.0004	0.0016	0.0003	ND		ND		ND	
	ND		ND		ND		ND		ND		0.003	0.0003	ND		ND		ND	
	ND		ND		ND		ND		ND		ND		ND		ND		ND	
	ND		ND		ND		ND		ND		0.0062	0.0002	ND		ND		ND	
	ND		ND		ND		0.0005	0.0002	ND		0.0024	0.0002	ND		ND		ND	
0.0006	ND		ND		0.0335	0.0008	0.0021	0.0003	0.0028	0.0005	0.0028	0.0003	ND		ND		ND	
0.0002	ND		0.0023	0.0001	0.0252	0.0004	0.0009	0.0001	0.0084	0.0003	ND		ND		ND		0.0058	0.0016
	ND		0.0026	0.0002	0.0213	0.0004	0.0015	0.0002	0.0097	0.0003	ND		ND		ND		ND	
0.0002	ND		0.0026	0.0002	0.0221	0.0004	0.0005	0.0001	0.0068	0.0003	ND		ND		ND		ND	
	ND		0.0032	0.0002	0.026	0.0004	0.0008	0.0001	0.0081	0.0003	ND		ND		ND		ND	
0.0002	ND		0.0028	0.0002	0.0273	0.0004	0.0013	0.0002	0.0089	0.0003	ND		ND		ND		ND	
	ND		0.0018	0.0001	0.0197	0.0003	ND		0.006	0.0002	ND		ND		ND		ND	
0.0003	ND		0.0024	0.0002	0.0279	0.0004	0.0011	0.0001	0.0136	0.0003	ND		ND		ND		ND	
0.0002	ND		0.0029	0.0002	0.026	0.0004	0.001	0.0001	0.0106	0.0003	ND		ND		ND		ND	
	ND		0.0024	0.0001	0.0217	0.0003	0.0008	0.0001	0.0078	0.0002	ND		ND		ND		ND	
	ND		0.0023	0.0001	0.0202	0.0003	0.0005	0.0001	0.0067	0.0003	ND		ND		ND		ND	
	ND		0.0021	0.0001	0.0256	0.0004	0.0006	0.0001	0.0071	0.0002	ND		ND		ND		ND	
	ND		0.0013	0.0001	0.016	0.0003	ND		0.0067	0.0002	ND		ND		ND		ND	
	ND		0.003	0.0002	0.032	0.0005	0.0011	0.0002	0.0055	0.0003	ND		ND		ND		ND	
0.0002	ND		0.0037	0.0002	0.0251	0.0004	ND		0.0056	0.0003	ND		ND		ND		ND	

As +/-	Se	Se +/-	Rb	Rb +/-	Sr	Sr +/-	Y	Y +/-	Zr	Zr +/-	Mo	Mo +/-	Ag	Ag +/-	Cd	Cd +/-	Sn	Sn +/-
	ND		0.0028	0.0002	0.0269	0.0004	0.0013	0.0002	0.0057	0.0003	ND		ND		ND		ND	
0.0003	ND		0.0032	0.0002	0.0242	0.0004	0.0011	0.0002	0.0053	0.0003	ND		ND		ND		ND	
0.0003	ND		0.0032	0.0002	0.0265	0.0005	0.0016	0.0002	0.0079	0.0003	ND		ND		ND		ND	
	ND		0.0019	0.0001	0.0227	0.0004	0.0016	0.0002	0.0065	0.0003	ND		ND		ND		ND	
	ND		0.0024	0.0002	0.0244	0.0004	0.0019	0.0002	0.0059	0.0003	ND		ND		ND		ND	
	ND		0.0027	0.0002	0.0238	0.0004	0.0011	0.0002	0.0081	0.0003	ND		ND		ND		ND	
	ND		0.0029	0.0002	0.023	0.0004	0.0018	0.0002	0.0092	0.0003	ND		ND		ND		ND	
	ND		0.0031	0.0002	0.0249	0.0004	0.0017	0.0002	0.0088	0.0003	ND		ND		ND		ND	
	ND		0.0042	0.0002	0.0355	0.0005	0.0014	0.0002	0.0055	0.0003	ND		ND		ND		ND	
	ND		0.0038	0.0002	0.0362	0.0005	0.0009	0.0002	0.0067	0.0003	ND		ND		ND		ND	
	ND		0.0035	0.0002	0.0317	0.0005	0.0015	0.0002	0.0059	0.0003	0.001	0.0002	ND		ND		ND	
	ND		0.0029	0.0002	0.0269	0.0005	0.0009	0.0002	0.0045	0.0003	ND		ND		ND		0.0066	0.002
	ND		0.0017	0.0001	0.0218	0.0004	0.002	0.0002	0.0221	0.0004	ND		ND		ND		ND	
	ND		0.0029	0.0002	0.0165	0.0003	0.002	0.0002	0.0078	0.0002	ND		ND		ND		ND	
	ND		0.0054	0.0002	0.0176	0.0003	0.0011	0.0002	0.0069	0.0003	ND		ND		ND		ND	
0.0003	ND		0.0025	0.0002	0.0209	0.0004	0.0021	0.0002	0.0056	0.0003	ND		ND		ND		ND	
	ND		0.0042	0.0002	0.0236	0.0004	0.0017	0.0002	0.0068	0.0003	ND		ND		ND		ND	
0.0002	ND		0.004	0.0002	0.0209	0.0003	0.0006	0.0001	0.0091	0.0003	ND		ND		ND		ND	
	ND		0.0042	0.0002	0.0207	0.0004	0.0015	0.0002	0.0091	0.0003	ND		ND		ND		ND	
	ND		0.0031	0.0002	0.0167	0.0003	0.0008	0.0001	0.008	0.0002	ND		ND		ND		ND	
	ND		0.0043	0.0002	0.0197	0.0004	0.0013	0.0002	0.0086	0.0003	ND		ND		ND		ND	
	ND		0.0037	0.0002	0.0195	0.0003	0.0008	0.0002	0.0077	0.0003	ND		ND		ND		ND	
0.0002	ND		0.0032	0.0002	0.0159	0.0003	0.002	0.0002	0.012	0.0003	ND		ND		ND		ND	
	ND		0.0025	0.0002	0.0319	0.0005	0.001	0.0002	0.0085	0.0003	ND		ND		ND		ND	
	ND		0.0027	0.0002	0.0234	0.0004	0.0016	0.0002	0.0105	0.0003	ND		ND		ND		ND	
0.0003	ND		0.0035	0.0002	0.0225	0.0004	0.0013	0.0002	0.0099	0.0003	ND		ND		ND		ND	
	ND		0.0023	0.0002	0.0269	0.0004	0.0007	0.0002	0.0069	0.0003	ND		ND		ND		ND	
	ND		0.0051	0.0002	0.0255	0.0005	0.001	0.0002	0.0065	0.0003	ND		ND		ND		ND	
	ND		0.0035	0.0002	0.0316	0.0005	0.0012	0.0002	0.0086	0.0003	ND		ND		ND		ND	
	ND	ND			0.0066	0.0006	0.0223	0.0011	0.0097	0.0017	0.0769	0.0012	ND		ND		ND	

Sb	Sb +/-	W	W +/-	Hg	Hg +/-	Pb	Pb +/-	Bi	Bi +/-	Th	Th +/-	U	U +/-	LE	LE +/-	Unit
ND		ND		ND		0.0057	0.0011	ND		0.0097	0.0015	0.0032	0.0008	50.28	0.58	%
ND		ND		ND		0.0012	0.0003	ND		ND		ND		66.2	0.19	%
ND		ND		ND		0.0015	0.0003	ND		ND		ND		67.68	0.18	%
ND		ND		ND		0.0016	0.0003	ND		0.0024	0.0006	ND		76.21	0.18	%
ND		ND		ND		0.0025	0.0005	ND		0.0028	0.0008	ND		59.98	0.34	%
ND		ND		ND		0.0019	0.0005	ND		0.0025	0.0008	ND		65.76	0.35	%
ND		ND		ND		0.0045	0.0006	ND		0.0031	0.0009	ND		61.87	0.3	%
ND		ND		ND		0.0019	0.0004	ND		ND		ND		64.74	0.23	%
ND		ND		ND		0.0025	0.0004	ND		ND		ND		66	0.24	%
ND		ND		ND		0.0023	0.0003	ND		ND		ND		66.49	0.2	%
ND		ND		ND		0.0038	0.0006	ND		ND		ND		55.34	0.33	%
ND		ND		ND		0.0021	0.0004	ND		ND		ND		76.63	0.22	%
ND		ND		ND		0.0014	0.0003	ND		ND		ND		70.07	0.18	%
ND		ND		ND		0.001	0.0003	ND		ND		0.0014	0.0003	75.91	0.18	%
ND		ND		ND		0.0017	0.0004	ND		ND		0.0019	0.0004	75.69	0.2	%
ND		ND		ND		0.0042	0.0008	ND		0.0048	0.0012	0.0028	0.0007	61.4	0.39	%
ND		ND		ND		0.0019	0.0003	ND		ND		ND		72.4	0.17	%
ND		ND		ND		0.0022	0.0003	ND		ND		ND		74.24	0.18	%
ND		ND		ND		0.0013	0.0003	ND		ND		ND		77.74	0.16	%
ND		ND		ND		0.0017	0.0003	ND		ND		ND		71.15	0.18	%
ND		ND		ND		0.0012	0.0003	ND		ND		ND		71.96	0.25	%
ND		ND		ND		0.0014	0.0003	ND		ND		ND		77.12	0.15	%
ND		ND		ND		0.0018	0.0003	ND		ND		ND		73.47	0.17	%
ND		ND		ND		0.0017	0.0003	ND		ND		ND		69.66	0.19	%
ND		ND		ND		0.0019	0.0003	ND		ND		ND		73.57	0.17	%
ND		ND		ND		0.0011	0.0003	ND		ND		ND		74.77	0.17	%
ND		ND		ND		0.0015	0.0003	ND		ND		ND		71.15	0.18	%
ND		ND		ND		0.0014	0.0003	ND		ND		ND		70.39	0.18	%
ND		ND		ND		0.0022	0.0004	ND		ND		ND		68.32	0.21	%
ND		ND		ND		0.0013	0.0003	ND		ND		ND		68.37	0.2	%

Sb	Sb +/-	W	W +/-	Hg	Hg +/-	Pb	Pb +/-	Bi	Bi +/-	Th	Th +/-	U	U +/-	LE	LE +/-
ND		ND		ND		0.0025	0.0004	ND		ND		ND		68.99	0.2 %
ND		ND		ND		0.002	0.0003	ND		ND		ND		68.35	0.21 %
ND		ND		ND		0.0023	0.0004	ND		ND		ND		69.47	0.23 %
ND		ND		ND		0.0017	0.0003	ND		ND		ND		68.87	0.26 %
ND		ND		ND		0.0016	0.0003	ND		ND		ND		69.11	0.2 %
ND		ND		ND		0.0013	0.0003	ND		ND		ND		70.06	0.19 %
ND		ND		ND		0.0019	0.0003	ND		ND		ND		68.75	0.26 %
ND		ND		ND		0.0023	0.0003	ND		ND		ND		71.47	0.19 %
ND		ND		ND		0.0017	0.0003	ND		ND		ND		68.79	0.2 %
ND		ND		ND		0.0019	0.0004	ND		ND		ND		67.09	0.22 %
ND		ND		ND		0.0023	0.0004	ND		ND		ND		72.72	0.21 %
ND		ND		ND		0.0019	0.0004	ND		ND		ND		71.61	0.22 %
ND		ND		ND		0.0017	0.0003	ND		ND		ND		70.43	0.2 %
ND		ND		ND		0.0016	0.0003	ND		ND		ND		68.67	0.18 %
ND		ND		ND		0.0015	0.0003	ND		ND		ND		73.27	0.18 %
ND		ND		ND		0.0011	0.0003	ND		ND		ND		75.99	0.18 %
ND		ND		ND		0.0021	0.0004	ND		ND		ND		71.77	0.19 %
ND		ND		ND		0.0016	0.0003	ND		ND		ND		72.64	0.17
ND		ND		ND		0.0023	0.0004	ND		ND		ND		72.73	0.19
ND		ND		ND		0.0024	0.0003	ND		ND		ND		72.08	0.17
ND		ND		ND		0.0018	0.0003	ND		ND		ND		70.8	0.27
ND		ND		ND		0.0019	0.0003	ND		ND		ND		72.9	0.18
ND		ND		ND		0.0013	0.0003	ND		ND		ND		72.04	0.18
ND		ND		ND		0.002	0.0003	ND		ND		ND		71.24	0.19
ND		ND		ND		0.0017	0.0003	ND		ND		ND		70.39	0.2
ND		ND		ND		0.0018	0.0003	ND		ND		ND		72.73	0.18
ND		ND		ND		0.0017	0.0003	ND		ND		ND		74.28	0.19
ND		ND		ND		0.0018	0.0004	ND		ND		ND		67.24	0.32
ND		ND		ND		0.0027	0.0004	ND		ND		ND		69.53	0.3
ND		ND		ND		ND		ND		0.2447	0.0078	0.0156	0.002		